

Occurrence, detection and detoxification of mycotoxins

VISENUO AIKO and ALKA MEHTA*

*School of Biosciences and Technology, VIT University, Vellore 632 014, India***Corresponding author (Email, alkamehta@vit.ac.in)*

Mycotoxins have been identified as important toxins affecting animal species and humans ever since the discovery of aflatoxin B₁ in 1960. Mycotoxigenic fungi are ubiquitous in nature and are held responsible for economic loss as they decrease crop yield and quality of food. The presence of fungi and their mycotoxins are reported not only in food grains but also in medicinal herbs and processed foods. Since prevention is not always possible, detoxification of mycotoxins have been attempted using several means; however, only few have been accepted for practical use, e.g. ammonia in the corn industry. Organizations such as the World Health Organization, US Food and Drug Administration and European Union have set regulations and safety limits of important mycotoxins, viz. aflatoxins, fusarium toxins, ochratoxin, patulin zearalenone, etc., to ensure the safety of the consumers. This review article is a brief and up-to-date account of the occurrence, detection and detoxification of mycotoxins for those interested in and considering research in this area.

[Aiko V and Mehta A 2015 Occurrence, detection and detoxification of mycotoxins. *J. Biosci.*] DOI 10.1007/s12038-015-9569-6

1. Introduction

The word mycotoxin is derived from the Greek word 'mykes' meaning 'fungus' and the Latin word 'toxicum' meaning 'poison'. They are low molecular weight molecules produced as secondary metabolites by saprophytic fungi, especially *Aspergillus*, *Penicillium* and *Fusarium*. Mycotoxins have been known to mankind since the 1800s as 'St. Anthony's Fire' caused by ergot alkaloids, and as 'Alimentary Toxic Aleukia' caused by T2 toxins during World War II in Russia (Richard 2003). It was not until the 1960s, with the outbreak of 'Turkey X' disease in England, that mycotoxins were identified as important toxins. Turkey X disease refers to the death of 100,000 Turkey poultts due to consumption of peanut meal contaminated with fungi (Blount 1961). The responsible fungus was identified as *Aspergillus flavus* and the toxin as aflatoxin. Aflatoxins are acutely and chronically toxic to humans and animals. They cause liver and kidney damage, and induce mutagenic, carcinogenic and

immunosuppressive effects. Among the mycotoxins, aflatoxin B₁ is considered the most toxic and is classified as a group I carcinogen by the International Agency for Research on Cancer (IARC 2002). Some of the important mycotoxins of significant health hazards are listed in table 1.

There have been several outbreaks of mycotoxicosis in the human population. In 1974, an outbreak of hepatitis in India resulted in the death of 100 people due to consumption of contaminated maize (Krishnamachari *et al.* 1975). In another case in India, an outbreak of gastrointestinal disorder associated with consumption of bread made of contaminated wheat was reported in 1987. The contaminating moulds consisted of *Fusarium* sp. and *Aspergillus* sp., and the toxins were identified as deoxynivalenol, nivalenol, acetyl-deoxynivalenol and T2 toxin (Bhat *et al.* 1989). The Kenyan outbreak in 2004 was one of the largest outbreaks, where 125 people died due to liver failure caused by acute aflatoxicosis after consumption of contaminated maize (Muture and Oqana 2005).

Keywords. Aflatoxin; contamination; degradation; toxic effect

Table 1. Important mycotoxins affecting humans (CAST 2003)

Mycotoxins	Producer fungi	Commodities	Toxic effect
Aflatoxins (B ₁ , B ₂ , G ₁ , G ₂ , M ₁ , M ₂)	<i>A. flavus</i> , <i>A. parasiticus</i> , <i>A. nominus</i> , <i>A. tamari</i>	Peanuts, corn, wheat, rice, milk, cheese, figs, herbs	Mutagenic, carcinogenic, hepatotoxic, immunosuppressive
Citrinin	<i>P. citrinum</i> , <i>P. viridicatum</i>	Wheat, barley, corn, rice	Nephrotoxic, carcinogenic.
Deoxynivalenol (Trichothecenes)	<i>F. graminearum</i> , <i>F. culmorum</i>	Corn, wheat, barley, oats	Vomiting, diarrhea, immunosuppressive
Fumonisin	<i>F. verticillioides</i> , <i>F. proliferatum</i>	Corn, wheat	Tumors of the kidney and liver
Ochratoxin A	<i>A. ochraceus</i> , <i>A. flavus</i> , <i>P. viridicatum</i>	Cereals, beans, peanuts, cheese, coffee, dried fruits, grapes, wine	Nephrotoxic, hepatotoxic, teratogenic, carcinogenic,
Patulin	<i>P. patulum</i> , <i>P. expansum</i>	Apples, apple juice	Subcutaneous sarcomas, hemorrhage, carcinogenic
T-2 toxin (Trichothecenes)	<i>F. sporotrichoides</i> , <i>F. poae</i> , <i>F. roseum</i>	Corn, wheat, barley, oats	Emetic, cytotoxic, teratogenic
Zearalenone	<i>F. graminearum</i> , <i>F. tricinatum</i> , <i>F. Culmorum</i>	Corn, hay	Hyper-estrogenic, abortion

2. Occurrence of mycotoxins

The ubiquitous nature of fungi makes food crops vulnerable to fungal contamination during pre-harvest and post-harvest conditions. In the field, the contaminating fungi are airborne or transmitted by insects, and damaged kernels often become infected. Stress conditions like drought, floods, insect infestation and delayed harvest increase the level of contamination. Post-harvest conditions such as inadequate drying, warm humid environment during storage lead to mould formation. The extent of contamination by fungi depends on various factors like geographic location, processing and storing periods of the crops. Environmental factors like temperature, water activity or pH, damage of crop by insects, crop densities, etc., influence the growth of fungi and mycotoxin production (Magan and Olsen 2004).

2.1 Mycotoxins in food and feed

A survey of mycotoxin contamination in animal feeds in European and Mediterranean markets and Asia-Pacific regions was carried out by Binder *et al.* (2007). Their result showed the occurrence of deoxynivalenol, zearalenone and T2 toxin as the major contaminants in the European samples, while aflatoxins, fumonisins, deoxynivalenol and zearalenone were mainly found in the samples from Asia and Pacific regions. Herzallah (2009) has reported the presence of aflatoxin M₁ and M₂ in milk samples and aflatoxins B₁, B₂, G₁ and G₂ in meat samples collected from the local markets in Jordan. Rice is the staple food of India, consumed almost daily in every household, and if contaminated with aflatoxins, will have high impact on human health. A survey by Reddy *et al.* (2009) revealed the presence of several species of *Aspergillus* and aflatoxin B₁ in rice samples consisting of paddy and milled rice. Occurrence of *Aspergillus* and aflatoxins in rice grains were also reported from other countries like China, Nigeria and United Arab Emirates (Hussaini *et al.* 2007; Osman *et al.* 1999; Zuoxin *et al.* 2006). Maize samples in Vietnam intended for human and animal consumption were reported to be contaminated with high fungal load and aflatoxin B₁ and fumonisin B₁ (Trung *et al.* 2008). Alborch *et al.* (2012) isolated several species of *Aspergillus*, *Fusarium*, *Penicillium* and *Mucorales* from maize flour and popcorn kernels in Spain, and reported the natural contamination of aflatoxin B₁ and ochratoxin A in the samples. Mycotoxins such as aflatoxins, ochratoxin and fumonisins have been detected in processed foods which were sold in the market (Sugita-Konishi *et al.* 2006; Romagnoli *et al.* 2007; Mushtaq *et al.* 2012). Aflatoxins and fumonisins were also detected in maize-based products intended for human consumption in Glasgow, UK (Candlish *et al.* 2000).

2.2 Mycotoxins in medicinal herbs

Studies in India showed the natural occurrence of aflatoxin B₁ and citrinin in medicinal plants and herbal drugs (Roy and Kumari 1991). Aflatoxin B₁, citrinin, ochratoxin A and zearalenone were detected in several medicinal plants such as *Asparagus racemosus*, *Carum ajmoda*, *Cinnamomum zeylanicum*, *Cuminum cyminum*, *Elettaria cardamomum*, *Embllica officinalis*, *Piper longum*, *P. nigrum*, *Saraca indica* and *Zingiber officinale* (Chourasia 1995). Thirumala-Devi *et al.* (2001) reported the contamination of *Coriandrum sativum*, *Piper nigrum*, *Zingiber officinale* and *Curcuma longa* with ochratoxin A in India. The incidence of toxigenic fungi producing aflatoxins, ochratoxin A and fumonisin on medicinal herbs was reported from Argentina (Rizzo *et al.* 2004). The medicinal herb ginseng was reported to be contaminated with aflatoxin B₁, ochratoxin A and zearalenone (Gray *et al.* 2004; Trucksess *et al.* 2006). An investigation from South Africa showed the presence of fumonisin B₁ in dietary (*Rumex lanceolatus*, *Zantedeschia aethiopica*, *Raphanus raphanistrum*, *Solanum nigrum*) and medicinal (*Catha edulis*, *Dalbergia obovata*, *Brunsvigia* sp., *Datura stramonium*) wild plants (Sewram *et al.* 2006). Aflatoxins-, ochratoxin-A- and citrinin-producing *Aspergillus* and *Penicillium* were isolated from medicinal herbs in Brazil (Bugno *et al.* 2006). The presence of aflatoxin B₁ in *Pimpinella anisum*, *Piper nigrum*, *Mentha piperita* and *Origanum majorana* was reported by Bokhari (2007) in Saudi Arabia. In Korea, a survey was conducted on spices and processed spice products for aflatoxin contamination, where aflatoxin B₁ was detected in 13.6% of the spices (Cho *et al.* 2008). Multi-contamination of mycotoxins with T2 toxin, zearalenone, aflatoxins, ochratoxin A, deoxynivalenol, citrinin and fumonisin were detected in 84 medicinal herbs surveyed in Spain (Santos *et al.* 2009).

3. Mechanism of toxicity

3.1 Aflatoxin

Aflatoxin B₁ is a well-known human carcinogen. The World Health Organization has reported that hepatocellular carcinoma (HCC) or liver cancer is the third leading cause of cancer death worldwide and chronic aflatoxicosis leading to the development of HCC has been implicated (Wild and Gong 2010). Aflatoxin B₁ is activated by cytochrome P450 to form aflatoxin B₁-8,9-epoxide, which is responsible for the mutagenic activity of aflatoxin B₁ (McLean and Dutton 1995). Aflatoxin B₁-8,9-epoxide specifically binds to the N⁷ position of guanine of DNA and RNA to form aflatoxin B₁-N⁷-guanine

adduct (Croy *et al.* 1978). Aflatoxin B₁ inhibits DNA, RNA and protein synthesis, resulting in immuno-suppressive, hormonal and teratogenic effects (McLean and Dutton 1995).

3.2 Ochratoxins

Ochratoxin primarily affects the kidney of all animal species and at high concentration can affect the liver. The toxic effect has been attributed to inhibition of phenylalanine tRNA synthetase. It is also an immune suppressor, teratogen and a carcinogen (Kuiper-Goodman and Scott 1989) and its role in the Balkan endemic nephropathy has been implicated (Hult *et al.* 1982). Studies have shown the formation of ochratoxin A-DNA adducts in the kidney and bladder tissues of Bulgarian patients undergoing surgery for cancer (Pfohl-Leszkiwicz *et al.* 1993).

3.3 Citrinin

Citrinin is hepatotoxic and nephrotoxic to a number of animal species, and the possible role of citrinin and ochratoxin A in the Balkan endemic nephropathy was implicated (Vrabcheva *et al.* 2000). Citrinin has been shown to inhibit RNA, DNA and protein synthesis in porcine kidney at 0.01 mM concentration (Braunberg *et al.* 1992).

3.4 Fumonisin

Fumonisin B₁ and B₂ are known to cause leukoencephalomalacia in horse (Marasas *et al.* 1988), and pulmonary edema and hydrothorax in swine (Colvin and Harrison 1992). This toxin has been reported to interfere with sphingolipid metabolism by inhibiting ceramide kinase. The high incidence of esophageal cancer in South Africa, China and Italy has been correlated with fumonisin B₁ (Peraica *et al.* 1999). In a recent study, Wang *et al.* (2014) reported that fumonisin B₁ stimulated the proliferation of normal human esophageal epithelial cells increasing the protein expression of cyclin D1 and decreasing cyclin E, p21 and p27.

3.5 Trichothecenes

Deoxynivalenol, diacetoxyscirpenol and T2 are the most common trichothecenes. They are known to cause nausea, vomiting, diarrhea and suppress the immune system in animals. Trichothecenes are reported to be potent inhibitors of protein synthesis by direct inhibition of peptidyl transferase in the large ribosome subunit (Feinberg and McLaughlin 1989).

3.6 Zearalenone

Zearalenone, also known as a mycoestrogen or phytoestrogen, mainly affects swine causing hyperestrogenism (Kurtz and Mirocha 1978). The exact mechanism of toxicity is not known however it has been reported that zearalenone is metabolized into α - and β -zearalenol which conjugates with glucuronic acid (Olsen 1989).

4. Detection of mycotoxins

Mycotoxins occur naturally and frequently in food and feeds, as mentioned earlier. The toxic nature of mycotoxins makes their detection an absolute necessity. Several detection methods have been developed, and among them chromatographic techniques are widely used. The procedure for detecting mycotoxins involves extraction from sample material, purification, and qualitative and quantitative analysis. The most common methods currently used are described here.

4.1 Thin layer chromatography (TLC)

TLC is one of the traditional methods of detecting mycotoxins. This technique enables screening of large number of samples, easy identification and is cost-effective. A silica gel layer is most commonly used; however, phenyl non-polar bonded, silanized and polyamide are also been used (Lin *et al.* 1998). Mycotoxins are visualized on the TLC plate by observing under UV light or by spraying chemicals which react with mycotoxins and enhance the fluorescence or produce colour products (Betina 1985). Aflatoxins, citrinin and ochratoxin are naturally fluorescent compounds; hence, they are identified based on their fluorescent properties. For example, the B and G aflatoxins are differentiated by blue and green fluorescence, respectively, while citrinin is identified by yellow fluorescence (Betina 1985). Aflatoxins have been identified by chemical confirmation by spraying trifluoroacetic acid and sulphuric acid on the TLC plate (Stack and Pohland 1975). Serralheiro and Quinta (1985) have reported that spraying sulphuric acid improves the limit of detection of aflatoxin M₁ from 0.5 $\mu\text{g}/\text{kg}$ to 0.3 $\mu\text{g}/\text{kg}$. Semi-quantitative analysis has been carried out for mycotoxins by TLC; however, the method has low sensitivity.

The TLC method has been improved in high-performance thin layer chromatography (HPTLC) to enhance the resolution and accuracy. HPTLC has been used to determine aflatoxins in peanut products and was shown to be equivalent to liquid chromatography in precision, accuracy and sensitivity (Tosch *et al.* 1984).

4.2 High-performance liquid chromatography (HPLC)

HPLC provides higher accuracy and precision of mycotoxin determination. Normal and reversed-phase HPLC are used with a variety of detection systems. UV and fluorescence detectors are most commonly used. Pons and Franz (1978) reported accurate and sensitive detection of all aflatoxins at levels of 0.3–1 $\mu\text{g}/\text{kg}$, whereas aflatoxins B₁ and B₂ were detected by a UV detector at 360–365 nm, and G₁ and G₂ by fluorescence. Akiyama *et al.* (1998) detected non-fluorescent mycotoxins, fumonisins, by using *o*-phthalaldehyde post-column derivatization and then detection by fluorescence detector. The detection limit of fumonisin by this method was reported to be 10 $\mu\text{g}/\text{kg}$ of corn. Ochratoxin A in wine has been accurately detected by HPLC following immunoaffinity clean-up with a detection limit of 0.01 ng/mL (Visconti *et al.* 1999). HPLC with fluorescence detector has been used for detecting aflatoxin B₁, citrinin and ochratoxin in rice, and a detection limit of 0.07, 0.11 and 0.08 $\mu\text{g}/\text{kg}$, respectively, for these mycotoxins was reported by Nguyen *et al.* (2007).

4.3 Liquid chromatography-mass spectrometry (LC-MS)

Liquid chromatography coupled with mass spectrometry eliminates the need for sample derivatization for fluorescent activity. LC-MS is a very selective and sensitive method for identification and quantification of mycotoxins. Spanjer *et al.* (2008) had developed an LC-MS/MS method to detect 33 mycotoxins simultaneously in various food materials. The mycotoxins include aflatoxin B₁, B₂, G₁ and G₂, and ochratoxin A with a limit of quantification of 1 $\mu\text{g}/\text{kg}$ and 50 $\mu\text{g}/\text{kg}$ for deoxynivalenol. An ultra-high-performance liquid chromatography combined with electrospray ionization triple quadrupole tandem mass spectrometry (UHPLC-ESI-MS/MS) has been developed to determine aflatoxin M₁, ochratoxin A, zearalenone and α -zearalenol in milk. The limits of quantification of these toxins were reported to be in the range 0.003 to 0.015 $\mu\text{g}/\text{kg}$ (Huang *et al.* 2014). LC-MS/MS was used for detection and quantification of mycotoxins in blood and urine samples. In a recent study, 23 mycotoxins and their metabolites were monitored in human population of Bangladesh and Germany (Gerding *et al.* 2015).

4.4 Gas chromatography (GC)

This method is also used for detecting mycotoxins especially trichothecenes in food samples. Most mycotoxins are non-volatile and hence are derivatized for detection (Scott 1995). Electron capture detection (ECD), mass spectrometry (MS) and flame ionization are the common detectors used with

GC. Croteau *et al.* (1994) analysed trichothecenes in corn using GC. The trichothecene mycotoxins were derivatized using heptafluorobutyric anhydride and detected by ECD. The limit of quantification was reported in the range 50–200 µg/kg of corn. In another report, trichothecenes were determined in corn by MS detector with a detection limit of 10–40 µg/kg and limit of quantification of 70–200 µg/kg (Milanez and Valente-Soares 2006). The GC method has also been used for analysing multi-mycotoxins such as patulin, zearalenone and trichothecenes in wheat (Rodríguez-Carraso *et al.* 2012). There are a few disadvantages with this method such as the need for derivatization and thermal stability of mycotoxins, where heating degrades the samples.

4.5 Fluorescence spectrometry

Fluorescence property of several mycotoxins is an important characteristic for their detection. A novel biosensor based on surface plasmon-enhanced fluorescence spectroscopy has been developed for detecting aflatoxin M₁ in milk with a detection as low as 0.6 pg/mL (Wang *et al.* 2009).

4.6 Fourier transform infrared spectroscopy

The use of infrared spectroscopy has proved to be a promising technique for the fast and non-destructive detection of mycotoxins in food grains. Deoxynivalenol was detected in wheat kernel samples using near-infrared spectroscopy at concentrations above 400 µg/kg (Pettersson and Aberg 2003). In a similar study, Kos *et al.* (2003) used mid-infrared to detect deoxynivalenol at concentrations as low as 310 µg/kg in corn samples. Near-infrared spectroscopy technique has also been used to detect aflatoxin B₁ and ochratoxin A in red paprika in Spain (Hernandez-Hierro *et al.* 2008).

4.7 Enzyme-linked immunosorbent assay (ELISA)

ELISA technique has been used for determining aflatoxin in a large number of foods. This method is based on direct and indirect competitive assay. ELISA has been used for detecting deoxynivalenol and zearalenone in maize (Cavaliere *et al.* 2005). Reddy *et al.* (2009) have used indirect competitive ELISA for detecting aflatoxin B₁ in rice with a detection limit of 0.02 ng/kg. Other mycotoxins like fumonisin have also been detected using this method with a detection limit of 3 ng/mL of beer (Torres *et al.* 1998). This method has the advantage of screening bulk samples and is highly specific. The performance of various detection methods differs for different food materials. Dell *et al.* (1990) has reported the determination of aflatoxin in peanut butter using

high-performance TLC which gave precise and consistent data than HPLC and ELISA.

4.8 Radioimmunoassay (RIA)

The other immunological methods for mycotoxin detection include radioimmunoassay. Sun and Chu (1977) had developed a solid-phase RIA for detecting aflatoxin B₁ in corn and wheat. Commercial kit for the application of RIA has been validated for determining ochratoxin A in food and feed stuffs with a detectable limit of 1 µg/kg (Fukal 1990). RIA was also used for detecting nivalenol in barley and the result was reported to be consistent with gas chromatographic analysis (Teshima *et al.* 1990).

4.9 Lateral flow devices

Lateral flow or dipstick immunoassay, developed using the principle of ELISA, are being successfully used for detecting mycotoxins. This technique was used for screening aflatoxin B₁ and ochratoxin A simultaneously in chili samples, with limit of quantification of 2 and 10 µg/kg, respectively (Saha *et al.* 2007). Deoxynivalenol and zearalenone were detected in wheat samples using colloidal gold-based lateral flow immunoassay (Kolosova *et al.* 2007). They have reported cut-off levels of 1500 and 100 µg/kg for deoxynivalenol and zearalenone, respectively. Other studies include aflatoxin B₁ detection in pig feed (Delmulle *et al.* 2005), T2 toxin in wheat and oat (Molinelli *et al.* 2008) and aflatoxin M₁ in milk (Zhang *et al.* 2012). This method is very simple and rapid where the result is obtained within 10 min and offers a convenient on-site screening tool.

4.10 Signal amplification method

With the advancement of technology, very low concentration of mycotoxin can be detected using the signal amplification technique. Pal and Dhar (2004) had developed an analytical device for immunoassay using an improved catalysed reporter deposition method of signal amplification. Groundnut, corn, wheat, cheese and chili were analysed for aflatoxin B₁ using this method, and a detection limit of 0.1 ng/mL was reported. In another study, a gold nano-particle-enhanced surface plasmon resonance imaging chip was designed to detect multiple mycotoxins using a competitive immunoassay (Hu *et al.* 2014). They have reported high specific and sensitive detection of aflatoxin B₁, ochratoxin A and zearalenone with low detection limits of 8, 30 and 15 pg/mL, respectively.

5. Detoxification of mycotoxins

Removal or detoxification of mycotoxins has been studied using physical, chemical or biological methods. Efficient degradation of mycotoxins is a challenge since most mycotoxins are heat-stable and form toxic degradation products. Although several detoxification methods have been developed, only a few have been accepted for practical use. Some of the common methods are described here.

5.1 Physical treatment

Physical treatment includes cooking, boiling, roasting, microwave heating, extrusion, irradiation, etc. Food undergo heat treatment during the processing stage, and hence thermal inactivation of mycotoxins is practical. Mycotoxins are relatively heat-stable and so are not easily destroyed (Bullerman and Bianchini 2007). The level of mycotoxin degradation by thermal process depends on factors like temperature, moisture content and time period. In heat treatment, temperature and time period are important in determining the level of degradation. Higher level of aflatoxin degradation was achieved when heated at 200°C for longer exposure time (Levi 1980). The moisture content of a product also plays an important part in degrading aflatoxins. At high moisture content, degradation was found more efficient (Mann *et al.* 1967). Under dry conditions, citrinin was decomposed at 170°C, whereas under moist condition it was detoxified at 140°C (Kitabatake *et al.* 1991). Heating ochratoxin A in the presence of sodium hydroxide (NaOH) resulted in the detoxification of the toxin (Trivedi *et al.* 1992). Roasting fumonisin B₁ contaminated cornmeal at 218°C for 15 min resulted in almost complete degradation of the toxin (Castelo *et al.* 1998). The presence of ammonia during extrusion of aflatoxin B₁ led to higher amount of degradation (Hameed 1993).

Aflatoxins are photosensitive in nature; hence, various radiations such as sunlight, UV light and gamma rays have been employed for degradation studies. Sunlight was efficiently used for degrading aflatoxin B₁ in olive oil, groundnut oil, etc. (Shantha and Sreenivasa Murthy 1977; Mahjoub and Bullerman 1988). Aflatoxin B₁ was found to be more susceptible to irradiation when present in liquid medium than in solid media. The cytotoxicity and mutagenicity of aflatoxin B₁ has been shown to reduce after treatment with UV in aqueous medium (Liu *et al.* 2011). On the other hand, it was reported that irradiated fungal inocula may produce increased levels of mycotoxins especially aflatoxins (Applegate and Chipley 1974) and ochratoxin (Applegate and Chipley 1976; Paster *et al.* 1985).

5.2 Chemical treatment

Treatment with chemicals efficiently degrades aflatoxin B₁; however, formation of degradation products was observed. Acids convert aflatoxin B₁ into several products such as aflatoxins B₂, B_{2a}, D₁, etc., rather than complete degradation. Shukla *et al.* (2002) reported the conversion of aflatoxin B₁ into aflatoxin B₂ and aflatoxin G₁ into aflatoxin G₂ by lactic acid. In a recent study, lactic acid has been shown to degrade aflatoxin B₁ into aflatoxin B₂ and B_{2a} efficiently, with aflatoxin B_{2a} as the major degradation product under heat treatment (Aiko *et al.* 2015). Citric acid causes the hydration of aflatoxin B₁ at the 8,9-olefinic bond of the terminal furan ring to form aflatoxin B_{2a} (Ciegler and Peterson 1968). Treatment of aflatoxin B₁ with hydrochloric acid at elevated temperatures completely destroyed the toxin without the formation of toxic residues (Williams and Dutton 1988; Wattanapat *et al.* 1995). Other acids like salicylic, sulphamic, sulposalicylic, anthranilic, benzoic, boric, oxalic and propionic acids were efficiently used for degrading aflatoxin B₁ by more than 90% in sorghum (Hasan 1996). Alkalis cause the hydrolysis of the lactone ring in aflatoxin B₁; however, it can revert back under acidic conditions (Price and Jorgensen 1985; Camou-Arriola and Price 1989). Boiling aflatoxin B₁ contaminated corn with NaOH decreased the level of aflatoxin B₁ by 93%, with 18% reversion level after treatment with acid (Camou-Arriola and Price 1989). Nixtamalization (alkaline cooking of grains) was reported to be an efficient method in degrading fumonisin. Fumonisin-contaminated kernel corn on nixtamalization resulted in reduced toxicity of fumonisin (Voss *et al.* 2012). Sodium bisulphate and hydrogen peroxide were also used in degrading aflatoxin B₁ efficiently (Altug *et al.* 1990).

Among the many chemicals used for detoxification of mycotoxins, ammonia is the most efficient and it has been accepted for use by the corn production industry. Ammonia degrades aflatoxin B₁ into aflatoxin D₁ which has reduced toxicity and mutagenic potential (Lee and Cucullu 1978). Ozone has been used to degrade aflatoxin B₁ by more than 90% in animal feed (Prudente and King 2002). Maeba *et al.* (1988) showed that ozone-treated aflatoxins were not toxic and mutagenic. Aflatoxin B₂ and G₂, fumonisin, ochratoxin, patulin and zearalenone were also efficiently degraded by ozone.

5.3 Biological treatment

5.3.1 Microorganism: Few strains of lactic acid bacteria have been reported to remove aflatoxins B₁ and M₁ by binding non-covalently (El-Nezami *et al.* 1998). Heat-treated and acid-treated *Lactobacillus rhamosus* GG and *L. rhamosus* were able to remove zearalenone, indicating

that binding and not metabolism is the mechanism by which the toxins are removed (El-Nezami *et al.* 2002). Another bacteria *Flavobacterium aurantiacum* B-184 was found to remove aflatoxin B₁ irreversibly (Lillehoj *et al.* 1967). Line *et al.* (1994) studied the mechanism and suggested that the degradation of aflatoxin B₁ by *F. aurantiacum* is probably a mineralization phenomenon. A number of fungal species, especially *Phoma* sp., were reported to prevent the synthesis of aflatoxin B₁ and degrade the toxin as well (Shantha 1999). Degradation of mycotoxins occurs during fermentation of various foods such as milk (Megalla and Mohran 1984), dough fermentation in making bread (El-Banna and Scott 1983) or during beer brewing (Chu *et al.* 1975). Some toxigenic fungi (*Aspergillus parasiticus*, *A. flavus*) have been reported to degrade their own toxins (Doyle and Marth 1978; Hamid and Smith 1987). Degradation of aflatoxin B₁ depends on the type of substrate and the fungal strains used.

5.3.2 *Enzymes*: Several fungal enzymes have been reported to degrade aflatoxin B₁. *Armellaria tabescens* produced a multienzyme system which detoxified aflatoxin B₁ by opening the difuran ring (Liu *et al.* 1998). The enzyme peroxidase from *A. flavus* and *A. parasiticus* has been shown to degrade aflatoxins B₁ and G₁ (Singh 1998; Doyle and Marth 1979). A horseradish peroxidase enzyme from the plant *Raphanus sativa* has also been reported to degrade aflatoxin B₁ (Das and Mishra 2000).

5.3.3 *Plant extracts*: Use of botanicals as anti-fungal and anti-mycotoxin agent is considered safe to humans and environmental friendly. Various extracts from plants such as piperine from black and long peppers (Singh *et al.* 1994); lutein and xanthophylls from Aztec marigold (Meija *et al.* 1997); carotenoids from fruits and vegetables (Rauscher *et al.* 1998) were reported to suppress the toxicity and mutagenicity of aflatoxin B₁. The essential oils of several plants have been documented to possess strong anti-microbial property. The oil of *Illicium verum*, *Cymbopogon martini*, *Eucalyptus globulus*, *Cinnamomum zylenium*, etc., are reported to be anti-fungals (Bansod and Rai 2008; Huang *et al.* 2010). The powder and essential oil of *Cymbopogon*

citratum have been successfully used for inhibiting aflatoxin B₁ contamination and preserving the quality of melon seed under storage (Bankole and Joda 2004). The essential oils of *Cinnamomum jensenianum* (Tian *et al.* 2011), *Ocimum sanctum* (Kumar *et al.* 2010) and *Zataria multiflora* (Gandomi *et al.* 2009) were used efficiently against toxigenic fungi and aflatoxin B₁ and their safe use as natural preservative of food has been implicated. Among the number of plants, *Syzygium aromaticum* (clove) has been extensively studied for its anti-microbial property (Pinto *et al.* 2009). The oil of clove and its main component, eugenol, has been reported to inhibit *Aspergillus* growth and aflatoxin B₁ production by various investigators (Bullerman *et al.* 1977; Jayashree and Subramanyam 1999). Whole clove has also been shown to inhibit the growth of *A. flavus* and *P. citrinum* and their toxins in culture media and rice grains (Aiko and Mehta 2013a, b). Removal of toxigenic fungi and mycotoxins by botanicals are usually preferred over chemical treatments.

6. Management and regulation of mycotoxins

The Food and Agricultural Organization has reported that 25% of the world's food crops are contaminated with mycotoxins. This not only causes economic loss but also reduces the world's food supply. The contaminating fungi and mycotoxins are found in food crops as well as in a number of processed foods intended for human consumption. Mycotoxins pose higher risk of causing cancer than contaminants in food such as anthropogenic contaminants, pesticides, phycotoxins and food additives (Kuiper-Goodman 1998).

Several national and international organizations and agencies have set regulations and safety limits of various

Table 2. The regulation of mycotoxins in human food (µg/kg)

Mycotoxins	European Union	US FDA
Aflatoxin B ₁	2–8	20
Aflatoxin M ₁	0.05	0.5
Deoxynivalenol	200–700	1000
Fumonisin (FB ₁ , FB ₂ , FB ₃)	200–1000	2000–4000
Ochratoxin	3–10	-
Patulin	10–50	-
Zearalenone	20–200	-

Table 3. Permissible limits of Aflatoxin B₁ in food set by various countries (Moss 2002)

Country	Aflatoxin B ₁ (µg/kg)	Products
Argentina	0	Groundnuts, maize and products
Brazil	15	All foodstuffs
China	10	Rice and edible oils
Czech Republic	5	All foods
Hungary	5	All foods
India	30	All foods
Japan	10	All foods
Nigeria	20	All foods
Poland	0	All foods
South Africa	5	All foods
Zimbabwe	5	Foods

mycotoxins. The maximum levels for mycotoxins in foods and feeds have been set to ensure the safety of the consumers. The US Food and Drug Administration (FDA) and European Union (EU) have set the maximum limit of the major mycotoxins in human food as given in table 2. With the recognition of aflatoxin B₁ as a human carcinogen, several countries have set the regulation of aflatoxin B₁ in foods (table 3).

7. Conclusion

Food safety is a major concern around the world. A number of researches are focused on the prevention and removal or detoxification of mycotoxins from food and feed. Control of mycotoxins largely depends on taking proper care during pre-harvest and post-harvest conditions. Use of fertilizers, pest control and fungal-resistant crops, and maintaining low moisture content and temperature during storage conditions can prevent fungal and mycotoxin contamination. However, prevention of mycotoxin contamination is not always possible; hence, many reduction or detoxification methods have been developed as mentioned above. These methods either degrade mycotoxins completely or reduce the toxin concentration to a safe level. Ammonia is currently used for degrading aflatoxin B₁ in feedstuffs; however, it also forms a degradation product aflatoxin D₁ which is not completely non-toxic (Lee and Cucullu 1978). It is of utmost importance to develop a safe and suitable detoxification technique without compromising the nutritional value of food. There is scope for developing an efficient and safer technique for mycotoxin detoxification and the intense research in the field can be very useful.

References

- Aiko V and Mehta A 2013a Control of *Aspergillus flavus* and aflatoxin production with spices in culture medium and rice. *World Mycotoxin J.* **6** 43–50
- Aiko V and Mehta A 2013b Inhibitory effect of clove (*Syzygium aromaticum*) on the growth of *Penicillium citrinum* and citrinin production. *J. Food Saf.* **33** 440–444
- Aiko V, Edamana P and Mehta A 2015 Decomposition and detoxification of aflatoxin B₁ by lactic acid. *J. Sci. Food Agric.* doi:10.1002/jsfa.7304
- Akiyama H, Uraroongroj M, Miyahara M, Goda Y and Toyoda M 1998 Quantitation of fumonisins in corn by HPLC with *o*-phthalaldehyde postcolumn derivatisation and their identification by LC/MS. *Mycopathologia* **140** 157–161
- Alborch L, Bragulat MR, Castella G, Abarca ML and Cabanes FJ 2012 Mycobiota and mycotoxin contamination of maize flours and popcorn kernels for human consumption commercialized in Spain. *Food Microbiol.* **32** 97–103
- Altug T, Yousef AE and Martin EH 1990 Degradation of aflatoxin B₁ in dried figs by sodium bisulfite with or without heat, ultraviolet energy or hydrogen peroxide. *J. Food Prot.* **53** 581–582
- Applegate KL and Chipley JR 1974 Effect of ⁶⁰Coγ on aflatoxin B₁ and B₂ production by *Aspergillus flavus*. *Mycologia* **66** 436–445
- Applegate KL and Chipley JR 1976 Production of ochratoxin A by *Aspergillus ochraceus* NRRL-3174 before and after exposure with ⁶⁰Co irradiation. *Appl. Environ. Microbiol.* **31** 349–353
- Bankole SA and Joda AO 2004 Effect of lemon grass (*Cymbopogon citratus* Stapf) powder and essential oil on mould deterioration and aflatoxin contamination of melon seeds (*Colocynthis citrullus* L.). *Afr. J. Biotechnol.* **3** 52–59
- Bansod S and Rai M 2008 Antifungal activity of essential oils from Indian medicinal plants against human pathogenic *Aspergillus fumigatus* and *A. niger*. *World J. Med. Sci.* **3** 81–88
- Betina V 1985 Thin-layer chromatography of mycotoxins. *J. Chromatogr.* **334** 211–276
- Bhat RV, Ramakrishna Y, Beedu SR and Munshi KL 1989 Outbreak of trichothecene mycotoxicosis associated with consumption of mould-damaged wheat products in Kashmir valley, India. *Lancet* **333** 35–37
- Binder EM, Tan LM, Chin LJ, Handl J and Richard J 2007 World-wide occurrence of mycotoxins in commodities, feeds and feed ingredients. *Anim. Feed Sci. Technol.* **137** 265–282
- Blount WP 1961 Turkey 'X' disease. *J. Brit. Turkey Federation.* **9** 55–58
- Bokhari FM 2007 Spices mycobiota and mycotoxins available in Saudi Arabia and their abilities to inhibit growth of some toxigenic fungi. *Mycobiology* **35** 47–53
- Braunberg RC, Gantt O, Barton C and Friedman L 1992 In vitro effects of the nephotoxins ochratoxin A and citrinin upon biochemical function of porcine kidney. *Arch. Environ. Contam. Toxicol.* **22** 464–470
- Bugno A, Almodovar AAB, Pereira TC, Pinto TJA and Sabino M 2006 Occurrence of toxigenic fungi in herbal drugs. *Braz. J. Microbiol.* **37** 47–51
- Bullerman LB and Bianchini A 2007 Stability of mycotoxins during food processing. *Int. J. Food Microbiol.* **119** 140–146
- Bullerman LB, Lieu FY and Seier SA 1977 Inhibition of growth and aflatoxin production by cinnamon and clove oil, cinnamic aldehyde and eugenol. *J. Food Sci.* **42** 1107–1109
- Camou-Arriola JP and Price RL 1989 Destruction of aflatoxin and reduction of mutagenicity of naturally contaminated corn during production of corn snack. *J. Food Prot.* **52** 814–817
- Candlish AAG, Aidoo KE, Smith JE and Pearson SM 2000 A limited survey of aflatoxins and fumonisins in retail maize-based products in the UK using immunoassay detection. *Mycotoxin Res.* **16** 2–8
- Castelo MM, Sumner SS and Bullerman LB 1998 Stability of fumonisins in thermally processed corn products. *J. Food Prot.* **61** 1030–1033
- Cavaliere C, D'Ascenzo G, Foglia P, Pastorini E, Samperi R and Lagana A 2005 Determination of type B trichothecenes and macrocyclic lactone mycotoxins in field contaminated maize. *Food Chem.* **92** 559–568
- Cho SH, Lee CH, Jang MR, Son YW, Lee SM, Choi IS, Kim SM and Kim DB 2008 Aflatoxin contamination in spices and

- processed spice products commercialized in Korea. *Food Chem.* **107** 1283–1288
- Chourasia HK 1995 Mycobiota and mycotoxins in herbal drugs of Indian pharmaceutical industries. *Mycol. Res.* **99** 697–703
- Chu FS, Chang CC, Ashoor SH and Prentice N 1975 Stability of aflatoxin B₁ and ochratoxin A in brewing. *Appl. Microbiol.* **29** 313–316
- Ciegler A and Peterson RE 1968 Aflatoxin detoxification: hydroxydihydro-Aflatoxin B₁. *Appl. Microbiol.* **16** 665–666
- Colvin BM and Harrison LR 1992 Fumonisin-induced pulmonary edema and hydrothorax in swine. *Mycopathologia* **117** 79–82
- Council for Agricultural Science and Technology (CAST) 2003 Mycotoxins: risks in plant, animal and human systems, Task force report, No.139. (Ames: Council for Agricultural Science and Technology) pp 1–191
- Croteau SM, Prelusky DB and Trenholm HC 1994 Analysis of trichothecene mycotoxins by gas chromatography with electron capture detection. *J. Agric. Food Chem.* **42** 928–933
- Croy RG, Essigmann JM, Reinhold VN and Wogan GN 1978 Identification of the principal aflatoxin B₁-DNA adduct formed in vitro in rat liver. *Proc. Natl. Acad. Sci. USA* **75** 1745–1749
- Das C and Mishra HN 2000 *In vitro* degradation of aflatoxin B₁ by horseradish peroxidase. *Food Chem.* **68** 309–313
- Dell MPK, Haswell SJ, Roch OG, Coker RD, Medlock VFP and Tomlins K 1990 Analytical methodology for the determination of aflatoxins in peanut butter: comparison of high-performance thin-layer chromatographic, enzyme-linked immunosorbent assay and high-performance liquid chromatographic methods. *Analyst* **115** 1435–1439
- Delmulle BS, De Saeger SM, Sibanda L, Barna-Vetro I and Van Peteghem CH 2005 Development of an immunoassay-based lateral flow dipstick for the rapid detection of aflatoxin B₁ in pig feed. *J. Agric. Food Chem.* **53** 3364–3368
- Doyle MP and Marth EH 1978 Aflatoxin is degraded by mycelia from toxigenic and non-toxigenic strains of aspergilli grown on different substrates. *Mycopathologia* **63** 145–153
- Doyle MP and Marth EH 1979 Peroxidase activity in mycelia of *Aspergillus parasiticus* that degrade aflatoxin. *Appl. Microbiol. Biotechnol.* **7** 211–217
- El-Banna AA and Scott PM 1983 Fate of mycotoxins during processing of foodstuffs I- aflatoxin B₁ during making of Egyptian bread. *J. Food Prot.* **46** 301–304
- El-Nezami H, Kankaanpaa P, Salminen S and Ahokas J 1998 Ability of lactic acid bacteria to bind to a common food carcinogen, aflatoxin B-1. *Food Chem. Toxicol.* **36** 321–326
- El-Nezami H, Polychronaki N, Salminen S and Mykkanen H 2002 Binding rather than metabolism may explain the interaction of two food-grade *Lactobacillus* strains with zearalenone and its derivatives α -zearalenone. *Appl. Environ. Microbiol.* **68** 3545–3549
- Feinberg B and McLaughlin CS 1989 Biochemical mechanism of action of trichothecene mycotoxins; in *Trichothecene mycotoxicoses: pathophysiological effects*, Vol. I (ed) VR Beasley (Boca Raton: CRC Press) pp 27–35
- Fukal L 1990 A survey of cereals, cereal products, feedstuffs and porcine kidneys for ochratoxin A by radioimmunoassay. *Food Addit. Contam.* **7** 253–258
- Gandomi H, Misaghi A, Basti AA, Bokaei S, Khosravi A, Abbasifar A and Javan AJ 2009 Effect of *Zataria multiflora* Boiss. Essential oil on growth and aflatoxin formation by *Aspergillus flavus* in culture media and cheese. *Food Chem. Toxicol.* **47** 2397–2400
- Gerding J, Ali N, Schwartzbord J, Cramer B, Brown DL, Degen GH and Humpf HU 2015 A comparative study of the human urinary mycotoxin excretion patterns in Bangladesh, Germany, and Haiti using a rapid and sensitive LC-MS/MS approach. *Mycotoxin Res.* **31** 127–136
- Gray SL, Lackey BR, Tate PL, Riley MB and Camper ND 2004 Mycotoxins in root extract of American and Asian ginseng bind estrogen receptors α and β . *Exp. Biol. Med.* **229** 560–568
- Hameed HG 1993 Extrusion and chemical treatments for destruction of aflatoxin in naturally-contaminated corn, Ph.D. thesis, University of Arizona
- Hamid AB and Smith JE 1987 Degradation of aflatoxin by *Aspergillus flavus*. *J. Gen. Microbiol.* **133** 2023–2029
- Hasan HAH 1996 Destruction of aflatoxin B₁ on sorghum grain with acids, salts and ammonia derivatives. *Cryptogam. Mycol.* **17** 129–134
- Hernandez-Hierro JM, Garcia-Villanova RJ and Gonzalez-Martin I 2008 Potential of near infrared spectroscopy for the analysis of mycotoxins applied to naturally contaminated red paprika found in the Spanish market. *Anal. Chim. Acta.* **622** 189–194
- Herzallah SM 2009 Determination of aflatoxins in eggs, milk, meat and meat products using HPLC fluorescent and UV detectors. *Food Chem.* **114** 1141–1146
- Hu W, Chen H, Zhang H, He G, Li X, Zhang X, Liu Y and Li CM 2014 Sensitive detection of multiple mycotoxins by SPRi with gold nanoparticles as signal amplification tags. *J. Colloid Interface Sci.* **431** 71–75
- Huang F, Zhao J, Zhou L, Wang J, Gong Y, Chen X, Guo Z, Wang Q, et al. 2010 Antifungal activity of the essential oil of *Illicium verum* fruit and its main component *trans*-Anethole. *Molecules* **15** 7558–7569
- Huang LC, Zheng N, Zheng BQ, Wen F, Cheng JB, Han RW, Xu XM, Li SL, et al. 2014 Simultaneous determination of aflatoxin M₁, ochratoxin A, zearalenone and α -zearalenol in link by UHPLC-MS/MS. *Anal. Methods* **146** 242–249
- Hult K, Piestina R, Habazin-Novak V, Radic B and Ceovic S 1982 Ochratoxin A in human blood and Balkan endemic nephropathy. *Arch. Toxicol.* **51** 313–321
- Hussaini AM, Timothy AG, Olufunmilayo HA, Ezekiel AS and Godwin HO 2007 Fungi and some mycotoxins contaminating rice (*Oryza sativa*) in Niger state, Nigeria. *Afr. J. Biotechnol.* **6** 99–108
- International Agency for Research on Cancer (IARC) 2002 *Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. Monograph on the evaluation of carcinogenic risk to humans*, Vol. 82 (Lyon: IARC)
- Jayashree T and Subramanyam C 1999 Antiaflatoxigenic activity of eugenol is due to inhibition of lipid peroxidation. *Lett. Appl. Microbiol.* **28** 179–183
- Kitabatake N, Trivedi AB and Doi E 1991 Thermal decomposition and detoxification of Citrinin under various moisture conditions. *J. Agric. Food Chem.* **39** 2240–2244

- Kolosova AY, De Saeger S, Sibanda L, Verheijen R and Van Peteghem C 2007 Development of a colloidal gold-based lateral-flow immunoassay for the rapid simultaneous detection of zearalenone and deoxynivalenol. *Anal. Bioanal. Chem.* **389** 2103–2107
- Kos G, Lohninger H and Krska R 2003 Development of a method for the determination of *Fusarium* fungi on corn using mid-infrared spectroscopy with attenuated total reflection and chemometrics. *Anal. Chem.* **75** 1211–1217
- Krishnamachari KAVR, Bhat RV, Nagarajan V and Tilak TBG 1975 Hepatitis due to aflatoxicosis. An outbreak in western India. *Lancet* **1** 1061–1063
- Kuiper-Goodman T 1998 Food safety: mycotoxins and phycotoxins in perspective; in *Mycotoxins and Phycotoxins- development in chemistry, toxicology and food safety. Proceedings of the IX IUPAC International Symposium* (eds) M Minaglia, H Van Egmond, C Brera and J Gilbert (Fort Collins: Alaken Inc) pp 25–48
- Kuiper-Goodman T and Scott PM 1989 Risk assessment of the mycotoxin ochratoxin A. *Biomed. Environ. Sci.* **2** 179–248
- Kumar A, Shukla R, Singh P and Dubey NK 2010 Chemical composition, antifungal and antiaflatoxic activities of *Ocimum sanctum* L. essential oil and its safety assessment as plant based antimicrobial. *Food Chem. Toxicol.* **48** 539–543
- Kurtz HJ and Mirocha J 1978 Zearalenone (F2) induced estrogenic syndrome in swine; in *Mycotoxic fungi, mycotoxins, mycotocoses*, Vol. 2 (eds) TD Wyllie and LG Morehouse (New York: Marcel Dekker) pp 1256–1264
- Lee SL and Cucullu AF 1978 Conversion of aflatoxin B₁ to aflatoxin D₁ in ammoniated peanut and cottonseed meals. *J. Agric. Food Chem.* **26** 881–884
- Levi C 1980 Mycotoxins in coffee. *J. AOAC Int.* **63** 1282–1285
- Lillehoj EB, Ciegler A and Hall HH 1967 Aflatoxin B₁ uptake by *Flavobacterium aurantiacum* and resulting toxic effects. *J. Bacteriol.* **93** 464–4715
- Lin L, Zhang J, Wang P, Wang Y and Chen J 1998 Thin-layer chromatography of mycotoxins and comparison with other chromatographic methods. *J. Chromatogr. A.* **815** 3–20
- Line EJ, Brackett RE and Wilkinson RE 1994 Evidence for degradation of aflatoxin B₁ by *Flavobacterium aurantiacum*. *J. Food Prot.* **57** 788–791
- Liu DL, Yao DS, Liang R, Ma L, Cheng WA and Gu LQ 1998 Detoxification of aflatoxin B₁ by enzymes isolated from *Armellaria tabescens*. *Food Chem. Toxicol.* **36** 563–574
- Liu R, Chang M, Jin Q, Huang J, Liu Y and Wang X 2011 Degradation of aflatoxin B₁ in aqueous medium through UV irradiation. *Eur. Food Res. Technol.* **233** 1007–1012
- Maeba H, Takamoto Y, Kamimura M and Miura T 1988 Destruction and detoxification of aflatoxins with ozone. *J. Food Sci.* **53** 667–668
- Magan N and Olsen M 2004 Mycotoxins in food: detection and control (Woodhead Publishing Series in Food Science, Technology and Nutrition)
- Mahjoub AM and Bullerman LB 1988 Effects of storage time, sunlight, temperature and frying on stability of aflatoxin B₁ in olive oil. *Lebensm. Wiss. Technol.* **21** 429–435
- Mann GE, Codifer LP Jr and Doller FG 1967 Effects of heat on aflatoxin in oilseed meals. *J. Agric. Food Chem.* **15** 1090–1092
- Marasas WFO, Kellerman TS, Gelderblom WCA, Coetzer JEW, Thiel PG and Van der Lugt JJ 1988 Leukoencephalomalacia in horse induced by fumonisin B₁ isolated from *Fusarium moniliforme*. *Onderstepoort J. Vet. Res.* **55** 197–203
- McLean M and Dutton MF 1995 Cellular interactions and metabolism of aflatoxin: an update. *Pharmacol. Ther.* **65** 163–192
- Megalla SE and Mohran MA 1984 Fate of aflatoxin B₁ in fermented dairy products. *Mycopathologia* **88** 27–29
- Meija EG, Pina GL and Gomez MR 1997 Antimutagenic activity of natural xanthophylls against aflatoxin B₁ in *Salmonella typhimurium*. *Environ. Mol. Mutagen.* **30** 346–353
- Milanez TV and Valente-Soares LM 2006 Gas chromatography-mass spectrometry determination of trichothecene mycotoxins in commercial corn harvested in the state of Sao Paulo; Brazil. *J. Braz. Chem. Soc.* **17** 412–416
- Molinelli A, Grossalber K, Fuhrer M, Baumgartner S, Sulyok M and Krska R 2008 Development of qualitative and semiquantitative immunoassay-based rapid strip tests for the detection of T-2 toxin in wheat and oat. *J. Agric. Food Chem.* **56** 2589–2594
- Moss MO 2002 Risk assessment for aflatoxins in foodstuffs. *Int. Biodeterior. Biodegrad.* **50** 137–142
- Mushtaq M, Sultana B, Anwar F, Khan MZ and Ashrafuzzaman M 2012 Occurrence of aflatoxins in selected processed foods from Pakistan. *Int. J. Mol. Sci.* **13** 8324–8337
- Mutere BN and Oqana G 2005 Aflatoxin levels in maize products during the 2004 food poisoning outbreak in Eastern Province of Kenya. *East Afr. Med. J.* **82** 275–279
- Nguyen MT, Tozlovanu M, Tran TL and Pfohl-Leszkowicz A 2007 Occurrence of aflatoxin B₁, citrinin and ochratoxin A in rice in five provinces of the central region of Vietnam. *Food Chem.* **105** 42–47
- Olsen M 1989 Metabolism of zearalenone in farm animals; in *Fusarium mycotoxins, taxonomy and pathogenicity*, 1st edition (eds) J Chelkowski (Amsterdam-Oxford-New York) pp 167–177
- Osman NA, Abdelgadir AM, Moss MO and Bener A 1999 Aflatoxin contamination of rice in the United Arab Emirates. *Mycotoxin Res.* **15** 39–44
- Pal A and Dhar TK 2004 An analytical device for on-site immunoassay. Demonstration of its applicability in semiquantitative detection of aflatoxin B₁ in a batch of samples with ultrahigh sensitivity. *Anal. Chem.* **76** 98–104
- Paster N, Barkai-Golan R and Padova R 1985 Effect of gamma radiation on ochratoxin production by the fungus *Aspergillus ochraceus*. *J. Sci. Food Agric.* **36** 445–449
- Peraica M, Radic B, Lucic A and Pavlovic M 1999 Toxic effects of mycotoxins in humans. *Bull. World Health Organ.* **77** 754–766
- Pettersson H and Aberg L 2003 Near-infrared spectroscopy for determination of mycotoxins in cereals. *Food Control.* **14** 229–232
- Pfohl-Leszkowicz A, Grosse Y, Castegnaro M, Nicolov IG, Chernozemsky IN, Bartsch H, Betbeder AM, Creppy EE, et al. 1993 Ochratoxin A-related DNA adducts in urinary tract tumours of Bulgarian subjects. *IARC Sci. Publ.* **124** 141–148
- Pinto E, Vale-Silva L, Cavaleiro C and Salgueiro L 2009 Antifungal activity of the clove essential oil from *Syzygium aromaticum* on *Candida*, *Aspergillus* and dermatophyte species. *J. Med. Microbiol.* **58** 1454–1462

- Pons WA Jr and Franz AO Jr 1978 High performance liquid chromatographic determination of aflatoxins in peanut products. *J. AOAC Int.* **61** 793–800
- Price RL and Jorgensen KV 1985 Effects of processing on aflatoxin levels and on mutagenic potential of tortillas made from naturally contaminated corn. *J. Food Sci.* **50** 347–357
- Prudente AD Jr and King JM 2002 Efficacy and safety evaluation of ozonation to degrade aflatoxin in corn. *J. Food Sci.* **67** 2866–2872
- Rauscher R, Edenharder R and Platt KL 1998 *In vitro* mutagenic and *in vivo* anticlastogenic effects of carotenoids and solvent extracts from fruits and vegetables rich in carotenoids. *Mutat. Res.* **413** 129–142
- Reddy KRN, Reddy CS and Muralidharan K 2009 Detection of *Aspergillus* spp. and aflatoxin B₁ in rice in India. *Food Microbiol.* **26** 27–31
- Richard JL 2003 Mycotoxins and human disease; in *Clinical mycology* (eds EJ Anaissie, MR McGinnis and MA Pfaller (New York: Churchill Livingstone) pp 589–598
- Rizzo I, Vedoya G, Maurutto S, Haidukowski M and Varsavsky E 2004 Assessment of toxicogenic fungi on Argentinean medicinal herbs. *Microbiol. Res.* **159** 113–120
- Rodriguez-Carraso Y, Berrado H, Font G and Manes J 2012 Multi-mycotoxin analysis in wheat semolina using an acetonitrile-based extraction procedure and gas chromatography-tandem spectrometry. *J. Chromatogr. A* **1270** 28–40
- Romagnoli B, Menna V, Gruppioni N and Bergamini C 2007 Aflatoxins in spices, aromatic herbs, herb-teas and medicinal plant marketed in Italy. *Food Control.* **18** 697–701
- Roy AK and Kumari V 1991 Aflatoxin and citrinin in seeds of some medicinal plants under storage. *Pharm. Biol.* **29** 62–65
- Saha D, Acharya D, Roy D, Shrestha D and Dhar TK 2007 Simultaneous enzyme immunoassay for the screening of aflatoxin B₁ and ochratoxin A in chili samples. *Anal. Chim. Acta* **584** 343–349
- Santos L, Marin S, Sanchis V and Ramos AJ 2009 Screening of mycotoxin multicontamination in medicinal and aromatic herbs sampled in Spain. *J. Sci. Food Agric.* **89** 1802–1807
- Scott PM 1995 Mycotoxin methodology. *Food Addit. Contam.* **12** 395–403
- Serralheiro ML and Quinta ML 1985 Rapid thin layer chromatographic determination of aflatoxin M₁ in powdered milk. *J. AOAC Int.* **68** 952–954
- Sewram V, Shephard GS, Merwe LVD and Jacobs TV 2006 Mycotoxin contamination of dietary and medicinal wild plants in the Eastern Cape Province of South Africa. *J. Agric. Food Chem.* **54** 5688–5693
- Shantha T 1999 Fungal degradation of aflatoxin B₁. *Nat. Toxins.* **7** 175–178
- Shantha T and Sreenivasa Murthy V 1977 Photo-degradation of aflatoxin in groundnut oil. *Indian J. Technol.* **15** 453–454
- Shukla RS, Verma RJ and Mehta DN 2002 Kinetic and mechanistic investigations on reductions of aflatoxins by lactic acid. *Bioorg. Med. Chem. Lett.* **12** 2737–2741
- Singh VP 1998 Enzymatic degradation of aflatoxins in shake flask cultures of *A. flavus* CMI 1102566: A possible role of peroxidase in biological control mechanism. *J. Phytol. Res.* **11** 7–10
- Singh J, Reen RK and Wrehel FJ 1994 Piperine, a major ingredient of black and long peppers, protects against aflatoxin B₁-induced cytotoxicity and micronuclei formation H411EC3 rat hepatoma cells. *Cancer Lett.* **86** 195–200
- Spanjer MC, Rensen PM and Scholten JM 2008 LC-MS/MS multi-method for mycotoxins after single extraction, with validation data for peanut, pistachio, wheat, maize, cornflakes, raisins and figs. *Food Addit. Contam. Part A Chem. Anal. Control Exp. Risk Assess.* **25** 472–489
- Stack ME and Pohland AE 1975 Collaborative study of a method for chemical confirmation of the identity of aflatoxin. *J. AOAC Int.* **58** 110–113
- Sugita-Konishi Y, Nakajima M, Tabata S, Ishikuro E, Tanaka T, Norizuki H, Itoh Y, Aoyama K, *et al.* 2006 Occurrence of aflatoxins, ochratoxins A, and fumonisins in retail foods in Japan. *J. Food Prot.* **69** 1365–1370
- Sun PS and Chu FS 1977 A simple solid-phase radioimmunoassay for aflatoxin B₁. *J. Food Saf.* **1** 67–75
- Teshima R, Hirai K, Sato M, Ikebuchi H, Ichinoe M and Terao T 1990 Radioimmunoassay of nivalenol in barley. *Appl. Environ. Microbiol.* **56** 764–768
- Thirumala-Devi K, Mayo MA, Reddy G, Emmanuel KE, Larondelle Y and Reddy DVR 2001 Occurrence of ochratoxin A in black pepper, coriander, ginger and turmeric in India. *Food Addit. Contam.* **18** 830–835
- Tian J, Huang B, Luo Z, Zeng H, Ban X, He J and Wang Y 2011 The control of *Aspergillus flavus* with *Cinnamomum jensenianum* Hand.-Mazz essential oil and its potential use as a food preservative. *Food Chem.* **130** 520–527
- Torres MR, Sanchis V and Ramos AJ 1998 Occurrence of fumonisins in Spanish beers analyzed by an enzyme-linked immunosorbent assay method. *Int. J. Food Microbiol.* **39** 139–143
- Tosch D, Walkling AE and Schlesier JF 1984 Comparison of liquid chromatography and high performance thin layer chromatography for determination of aflatoxin in peanut products. *J. AOAC Int.* **67** 337–339
- Trivedi AB, Doi E and Kitabatake N 1992 Detoxification of ochratoxin A on heating under acidic and alkaline conditions. *Biosci. Biotechnol. Biochem.* **56** 741–745
- Trucksess MW, Weaver CM, Oles CJ, D'Ovidio K and Rader JI 2006 Determination of aflatoxins and ochratoxin A in ginseng and other botanical roots by immunoaffinity column cleanup and liquid chromatography with fluorescence detection. *J. AOAC Int.* **89** 624–630
- Trung TS, Tabuc C, Bailly S, Querin A, Guerre P and Bailly JD 2008 Fungal mycoflora and contamination of maize from Vietnam with aflatoxin B₁ and fumonisin B₁. *World Mycotoxin J.* **1** 87–94
- Visconti A, Pascale M and Centonze G 1999 Determination of ochratoxin A in wine by means of immunoaffinity column clean-up and high performance liquid chromatography. *J. Chromatogr. A* **864** 89–101
- Voss KA, Riley RT, Moore ND and Burns TD 2012 Alkaline cooking (nixtamalization) and the reduction in the *in vivo* toxicity of fumonisin-contaminated corn in a rat feeding bioassay. *Food Addit. Contam., Part A* **30** 1415–1421
- Vrabcheva T, Usleber E, Dietrich R and Martlbauer E 2000 Co-occurrence of ochratoxin A and citrinin in cereals from

- Bulgarian villages with a history of Balkan endemic nephropathy. *J. Agric. Food Chem.* **48** 2483–2488
- Wang Y, Dostalek J and Knoll W 2009 Long range surface plasmon-enhanced fluorescence spectroscopy for the detection of aflatoxin M₁ in milk. *Biosens. Bioelectron.* **24** 2264–2267
- Wang SK, Wang TT, Huang GL, Shi RF, Yang LG and Sun GJ 2014 Stimulation of the proliferation of human normal esophageal epithelial cells by fumonisin B₁ and its mechanism. *Exp. Ther. Med.* **7** 55–60
- Wattanapat R, Nakayama T and Beuchat LR 1995 Characteristics of acid hydrolysate from defatted peanut flour. *J. Food Sci.* **60** 443–445
- Wild CP and Gong YY 2010 Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* **31** 71–82
- Williams KR and Dutton MF 1988 Destruction of Aflatoxin during the production of hydrolysed vegetable protein. *J. Food Prot.* **51** 887–891
- Zhang DH, Li PW, Zhang Q, Yang Y, Zhang W, Guan D and Ding XX 2012 Extract-free immunochromatographic assay for on-site tests of aflatoxin M₁ in milk. *Anal. Methods* **4** 3307–3313
- Zuoxin L, Junxia G and Jiujiang Y 2006 Aflatoxin in stored maize and rice grains in Liaoning Province, China. *J. Stored Prod. Res.* **42** 468–479

MS received 08 April 2015; accepted 12 August 2015

Corresponding editor: LUIS M CORROCHANO