

# Remediation of chromium contaminants using bacteria

P. Kanmani · J. Aravind · D. Preston

Received: 6 February 2011 / Revised: 24 April 2011 / Accepted: 25 August 2011 / Published online: 26 November 2011  
© CEERS, IAU 2011

**Abstract** Chromium contaminants emanating from industrial activities pose a significant threat to human's well-being. Chromium (III) and Chromium (VI) are the forms in which they are commonly encountered, of which the trivalent form is relatively benign. Hence, biological reduction of hexavalent chromium has been widely explored by researchers, yielding fruitful outcomes, opening up exciting avenues and also throwing up new challenges. This article attempts to review this area of research. Microbes, especially bacteria capable of Chromium (VI) reduction, belonging to a heterogeneous group have been isolated from contaminated sites. They exhibit plasmid-mediated chromate resistance and the reduction is enzymatically mediated. Reduction studies have been carried out with free and immobilized enzymes as well as whole cells. Experiments have been carried out in specifically designed bioreactors operated in batch and continuous modes. Although significant progress has been made, much needs to be done for its successful in situ application as the organism may not withstand the Chromium concentration or may be impeded by the presence of other toxicants. With molecular engineering, it may be possible to derive strains with improved performance even under stressful field conditions.

**Keywords** Bioreduction · Biosorption · Chromate reductase · Chromium · Resistance · Toxicity

P. Kanmani · J. Aravind (✉)  
Department of Biotechnology, Kumaraguru College  
of Technology, Coimbatore 641049, India  
e-mail: aravindj@gmail.com

D. Preston  
School of Biosciences and Technology, VIT University,  
Vellore 632014, India

## Introduction

Heavy metals that are frequently discharged in industrial effluents are of grave concern owing to their toxic effects. Chromium compounds find widespread industrial applications in electroplating, wood preservation, leather tanning and steel manufacturing sectors and subsequently end up as environmental contaminants.

## Chemistry of chromium

Chromium is a steel-gray, lustrous, hard, brittle metal of Group VI-B of the transition series with atomic number of 24 and atomic weight of 51.996. It exists in a number of valence states ranging from  $-2$  to  $+6$ . However, Cr(III) and Cr(VI) are of significance owing to their stability in natural environments, although Cr with valences of I, II, IV and V have also been shown to exist in a number of compounds. The recommended permissible limit in drinking water is  $50 \mu\text{g/L}$  for Cr(VI) (Krishnamurthy and Wilkens 1994).

## Transport and toxicity

Cr(III) is an essential nutrient in this diet, aiding in the metabolism of glucose and lipids (Anderson 1997). However, ingesting large amounts of it may cause health problems including lung cancer (Katz and Salem 1994). Cr(VI) on the other hand is a potent, extremely toxic carcinogen and has been estimated to be 10–100 times more toxic than Cr(III) (Katz and Salem 1994). This may be due to the fact that Cr(VI) penetrates mammalian cells more readily than Cr(III). The United States Environmental

Protection Agency (USEPA) determined its reference dose to be  $1 \text{ mg kg}^{-1} \text{ day}^{-1}$  (Katz and Salem 1994).

Cr(VI) enters the cell through the membrane sulphate transport channels (Ohtake and Silver 1994; Cervantes et al. 2000; Silver et al. 2001) and reacts with intracellular reductants such as glutathione and ascorbate. This generates short-lived intermediates Cr(V) and/or Cr(IV), free radicals and the end product Cr(III). Cr(V) undergoes a one-electron redox cycle to regenerate Cr(VI) by transferring the electron to oxygen. This process results in the formation of reactive oxygen species (ROS) capable of combining with DNA–protein complexes (Fig. 1). Cr(IV) also binds to cellular materials and impedes their normal physiological functions (Pesti et al. 2000; Cervantes et al. 2000). Skin and nasal irritation, eardrum perforation and lung carcinoma are a few of the adverse effects linked with Cr(VI) (Gibb et al. 2000a, b). Apart from these, it can have teratogenic effects too (Saxena et al. 1990). Environmental effects include alteration in soil microbial communities and reduction in their growth and activity (Turpeinen et al. 2004). Hence, it becomes imperative to remove chromium contaminants, especially Cr(VI) from the environment.


Most of the intracellular Cr(VI) reductants are one electron reducers, which generate Cr(V) and result in the formation of ROS that are predominantly responsible for the adverse effects of Cr(VI) (modified from Cheung and Gu 2007).

Remediation of Cr contaminants: physico-chemical versus biological methods


Conventional chromium removal consists of chemical reduction followed by alkali precipitation or use of ion exchange and adsorption. The former technique generates large quantities of sludge while the latter are expensive and also less specific when other anions of heavy metals are present. Hence, attention has been shifted to biological methods of remediation. Biotransformation and biosorption are commonly used technologies, which utilize the potential of microorganisms or plants to transform or adsorb heavy metals. Of these, biotransformation (i.e. reduction of toxic, mobile Cr(VI) to less toxic, less mobile Cr(III)) may be considered to be the better alternative. In biosorption, the contaminant gets accumulated in the biomass which again has to be disposed off, while in biotransformation, it is converted to an innocuous form. Under aerobic conditions, microbial chromate reduction is brought about by soluble reductases. In the absence of oxygen, both soluble and membrane bound reductases including cytochromes have been implicated. A large array of compounds including carbohydrates, proteins, fats, hydrogen, NAD(P)H and endogenous electron reserves can serve as electron donors in the reduction process (Wang 2000) (Fig. 2).

Under aerobic conditions, ChrR of *Pseudomonas putida* MK1 catalyzes one as well as two electron transfers to

**Fig. 1** Schematic diagram of Cr(VI) transport into the cell and its fate



**Fig. 2** Mechanisms likely to be involved in enzymatic reduction of Cr(VI) under aerobic and anaerobic conditions



Cr(VI) with the transient formation of Cr(V), whereas YieF of *Escherichia coli* brings about a four electron transfer resulting in the direct reduction of Cr(VI) to Cr(III), with the remaining electron getting transferred to oxygen. A membrane-associated Cr<sup>6+</sup> reductase isolated from *Bacillus megaterium* TKW3 has not been characterized. Under anaerobic conditions, both soluble and membrane-associated Cr(VI) reducing enzymes, including cytochromes, associated with the electron transfer system are involved (enzymes involved in the reduction of Cr(VI) are given in boxes, SR and MR represent soluble and membrane-associated reductases, respectively) (modified from Cheung and Gu 2007).

#### Focus of this review

This review broadly covers various aspects of Cr bioremediation, with the focus on bacteria-mediated Cr(VI) reduction, with more emphasis on recent findings. The genetic basis of Cr(VI) resistance which is plasmid rather than chromosome mediated and is attributed to reduced uptake, regulated efflux mechanisms or reduction along with parameters influencing the rate and extent of reduction by the isolates viz., temperature, pH, availability of oxygen, initial Cr concentration, cell density, exogenous electron donors, presence of nitrates–sulphates, other heavy metals, have been discussed. Reduction potentials of growing cells, resting cells and cell-free supernatants have been compared; purification of chromate reductases and study of enzyme kinetics in case of free and immobilized

systems have also been reviewed. Cr(VI) reduction by a few fungi and yeast has been briefly included for comparison sake.

#### Bacterial Cr(VI) reduction

Since the discovery of the first microbe *P. dechromaticans* capable of reducing Cr(VI) in the 1970s (Romanenko and Korenkov 1977), the search for chromate reducing microorganisms has continued and much information has been gathered on microbial-mediated Cr(VI) reduction, making its practical application feasible. Cr(VI)-reducing bacteria belonging to diversified genera have been isolated from a variety of environments, indicating that such potential is widespread in nature and can be harnessed to facilitate remediation of Cr contaminated sites.

Among tannery isolates, *Microbacterium* sp. MP30 grew aerobically in a medium supplemented with 15 mM chromate but did not reduce it. Under anaerobic conditions, it reduced 100 μM sodium chromate within 30 h at the expense of acetate as electron donor (Pattanapitpaisal et al. 2001). Moderately halophilic *Nesterenkonia* sp. MF2 was tolerant to a much higher concentration of 600 mM chromate and completely reduced 200 μM Cr(VI) in 24 h under aerobic condition. Initial Cr(VI) concentration up to 400 μM did not have a significant effect on Cr(VI) reduction. The isolate was capable of chromate reduction in the presence of various concentrations of salts (Amoozegar et al. 2007). A strain of *B. circulans* exhibited comparatively far better activity and reduced as much as

190 mM chromate both under anaerobic and aerobic conditions, maximum reduction being observed at 28 h of incubation (Chaturvedi et al. 2007). *Exiguobacterium* sp. ZM-2 showed resistance to 12.37 mM potassium chromate and reduced 500  $\mu\text{M}$  Cr(VI) within 56 h. The presence of metabolic inhibitors sodium azide and sodium cyanide severely affected chromate reduction, while 2, 4-dinitrophenol, and an uncoupling agent, stimulated chromate reduction (Alam and Malik 2008).

In one study using landfill isolates, *Bacillus* sp. XW-4 was tolerant to and reduced 1.9 mM Cr(VI). Chromate reduction by the strain was significantly enhanced by the presence of glucose (Liu et al. 2006). In another study, suspended cultures of *B. cereus* GIDM20, *B. fusiformis* GIDM22 and *B. sphearicus* GIDM64 exhibited more than 85% reduction of 1 mM Cr(VI) within 30 h. The activity was found to be mainly associated with the soluble fraction of cells, expressed constitutively and unaffected by presence of different metal ions except  $\text{Hg}^{2+}$  and  $\text{Ag}^+$ . Addition of NADH enhanced Cr(VI) reduction (Desai et al. 2008). Another bacterium, *Ochrobactrum* sp. showed Cr(VI) reduction ability under alkaline conditions. This isolate was tolerant to and reduced as much as 15.4 mM Cr(VI). In this case too, the addition of glucose caused a dramatic increase in Cr(VI)-reduction, while the presence of sulphate or nitrate had no influence (He et al. 2009). In another study, filamentous bacterium *Sphaerotilus natans* CSCr-3 isolated from activated sludge systems aerobically reduced up to 1.5 mM Cr(VI) in the presence of a carbon source (Caravelli et al. 2008). This reduction potential can be considered significant as *S. natans* is previously known only for its biosorption ability.

*Bacillus* sp. KCH2 and KCH3, *Leucobacter* KCH4 and *Exiguobacterium* KCH5 were isolated from Cr contaminated soil. Of these, KCH3 and KCH4 showed higher Cr(VI) tolerance (2 mM) and reduction (1.5 mM) than KCH5 (1.5 and 0.75 mM, respectively). Cr(VI) reduction was inhibited by  $\text{Hg}^{2+}$  and enhanced by  $\text{Cu}^{2+}$  (Sarangi and Krishnan 2008). Enriched cultures of indigenous bacteria isolated from a stream sediment contaminated by a pigment manufacturing industry reduced 31% of the 250  $\mu\text{M}$  Cr(VI), which is equivalent to the exchangeable Cr(VI) concentration in the sediment. The removal efficiency was increased with external supply of electron donor in the form of glucose, lactate or acetate (Lee et al. 2008).

During fermentative growth on D-xylose, isolates ES6 and WS01 (*Cellulomonas*) from US Department of Energy (DoE) Site decreased Cr(VI) concentrations from 0.04 mM to below the detection limit (0.002 mM) in less than 3 days and retained their ability to reduce Cr(VI) even after 4 months of incubation. Washed ES6 and WS01 cells also reduced Cr(VI) under non-growth conditions for over 4 months (Viamajala et al. 2007). Stimulation of such

organisms may lead to effective long-term, in situ passive reactive barriers for Cr(VI) removal.

*Staphylococcus epidermidis* L-02 isolated from a bacterial consortium used for the remediation of a chromate-contaminated constructed wetland system reduced Cr(VI) by using pyruvate as an electron donor under anaerobic conditions. The presence of nitrate increased the specific reduction rate. Under denitrifying conditions, Cr(VI) reduction was not inhibited by nitrite. The maximum specific reduction rate was as high as 8.8–9.8  $\mu\text{M}$  Cr  $10^{10}$  cells/h (Vatsouria et al. 2005). *Intrasporangium* sp. Q5-1 isolated from a Mn/Cr contaminated site showed 17 mM/L minimum inhibitory concentration (MIC) for Cr(VI). Resting cells also reduced chromate. When immobilized with compounding beads containing 4% Poly Vinyl Alcohol (PVA), 3% sodium alginate, 1.5% active carbon and 3% diatomite, the reduction rate remained unchanged (Yang et al. 2009). These immobilized cells have the advantage of stability, reusability and less clogging in continuous systems.

In an attempt to bring about Cr(VI) remediation using psychrophiles, core samples obtained from a Cr(VI) contaminated aquifer were enriched in Vogel Bonner medium supplemented with Cr(VI). The isolate *Arthrobacter aureescens* P4 was resistant to 19.2 mM Cr(VI) but reduction was slow or not observed at and above 1.9 mM Cr(VI) (Horton et al. 2006). This species has previously not shown to be capable of low temperature (10°C) Cr(VI) reduction.

Apart from such sites polluted by anthropogenic activities, chromate resistant bacteria have also been isolated from natural environments. *B. sphearicus* from naturally occurring chromium percolated soil of Andaman Islands was tolerant to 15.4 mM Cr(VI) and reduced >80% of it during growth. Soluble fraction of the cell was responsible for aerobic reduction of chromate by this organism (Pal and Paul 2004). *Burkholderia cepacia* MCMB-821 isolated from the alkaline crater lake of Lonar was resistant to 19.2 mM Cr(VI) and reduced 98% of the 1.4 mM Cr(VI) within 36 h in the presence of 2% salt and lactose as the electron donor. Electron paramagnetic resonance (EPR) spectroscopy data suggested the formation of transient Cr(V) intermediate during reduction (Wani et al. 2007) (Table 1).

#### Other novel findings

In an endeavour to assess the toxic effect of chromate on soil microbial communities, soil microcosms were artificially contaminated with up to 1,000 mg Cr(VI)/kg soil. This affected the structure and diversity of the soil bacterial community. Bacterial strains isolated from the microcosm contaminated with highest concentration of chromate belonged to the genus *Pseudomonas* and no cyanobacterial

**Table 1** Chromate reduction potentials of various bacterial isolate

Strain	Chromate conc. the isolate is capable of reducing (mg/L)	References
<i>Microbacterium</i> sp. MP-30	100 $\mu$ M	Pattanapitpaisal et al. (2001)
<i>Thermus scotoeductus</i> SA-01	500 $\mu$ M	Opperman and Heerden (2007)
<i>Exiguobacterium</i> sp. KCH-5	0.75 mM	Sarangi and Krishnan (2008)
<i>Bacillus</i> sp. KCH-3, <i>Leucobacter</i> KCH-4	1.5 mM	Sarangi and Krishnan (2008)
<i>Sphaerotilus natans</i>	1.5 mM	Caravelli et al. (2008)
<i>Arthrobacter aureescens</i> P4	1.9 mM	Horton et al. (2006)
<i>Ochrobactrum</i> sp. CSCr-3	15.3 M	He et al. (2009)

growth was detected in this, indicating their sensitivity. Some enzymes were inhibited, others stimulated and the ATP content in microcosms was strongly affected by chromate (Viti et al. 2006). Thus, the soil microbial community responds to chromate pollution through changes in community structure, in metabolic activity, and in selection for Cr(VI)-resistance.

Cr(VI) is usually known to be reduced to Cr(III). However, as a rare occurrence, *Shewanella oneidensis*, reduced Cr(VI) to Cr(II). Individual cell microanalysis demonstrated that the Cr(II) concentrated near the cytoplasmic membrane, suggesting intracellular localization of the terminal reduction pathway (Daulton et al. 2007).

Acetate was added to aid in the bioremediation of two closed soil-water systems representative of the subsurface environment close to chromium ore processing residue disposal sites; one had a pH of 7.7, the other 9.3. Cr(VI) reduction occurred in both systems as part of a cascade of microbially mediated terminal electron-accepting processes, occurring between nitrate and iron reduction. Cr(VI) and subsequently iron reduction took longer to start and were slower in the more alkaline system. At the point when Cr(VI) reduction was essentially complete, the microbial populations in both systems showed an increase in species closely related to  $\beta$ -proteobacteria that are capable of nitrate reduction (Stewart et al. 2007).

Cr(III) produced as a result of Cr(VI) reduction is mostly considered to be insoluble Cr(OH)<sub>3</sub>. However, Cr(VI) reduction in the presence of cellular organic metabolites formed both soluble and insoluble organo-Cr(III) end-products (Puzon et al. 2005). These organo-Cr(III) complexes may account for the mobile form of Cr(III) detected in the environment. Such soluble complexes have been demonstrated with four bacterial cultures (*S. oneidensis* MR1, *Cellulomonas* sp. ES6, *Rhodococcus* sp. and *Desulfovibrio vulgaris* var *Hildenborough*). Structural characterization of the complexes indicated that they were composed of multiple chromium ions bound together by the cellular metabolites (Puzon et al. 2008).

A better understanding of the organo-Cr(III) pathway will be useful in designing remediation strategies for Cr-impacted soils.

The shape of *Leucobacter* Ch1 cells before and after chromate reduction was observed using scanning and transmission electron microscopy. The composition of reduction products was determined using Energy Dispersive X-ray Analysis and EPR Spectroscopy and was found to contain 28.2% Cr in the form of Cr(OH)<sub>3</sub> precipitate (Zhu et al. 2008).

Cr(VI) migrating downwards from a chromium ore processing residue was of concern because groundwater emerging from the site was alkaline, visibly yellow and had an elevated Cr(VI) concentration. Sandy clay from immediately beneath the waste contained up to 3,000 mg/kg of Cr, and around 60% of 0.5 N HCl extractable iron in the form of reduced Fe(II). Very recent microcosm experiments showed that the soil contained a viable bacterial population capable of iron-reduction. This sandy clay layer, despite a pH value of 10.5, acted as a natural reactive zone beneath the waste as it accumulated chromium (Stewart et al. 2010). The mechanism of Cr(VI) reduction is believed, most likely to be an abiotic reaction with the Fe(II) present in the soil, and the Fe(II) being replenished by microbial iron-reduction.

#### Purification and characterization of bacterial chromate reductases

Chromate reductases have been purified and characterized by several researchers. Cloning the genes that encode chromate reductases can result in further advancements in bioremediation. Characterization of chromate reduction by a flavin reductase from *E. coli* revealed faster reduction when compared to chemical reduction by NADH and glutathione suggesting that unlike eukaryotes, enzymatic reduction is the dominant mechanism in bacterial cells (Puzon et al. 2002). Chromate reductase encoding genes vary according to the microorganism and also occurs in



plasmids, few literatures are available describing the gene sequence (Barak et al. 2006; Gonzalez et al. 2005).

A soluble Cr(VI) reductase purified from the cytoplasm of *E. coli* ATCC 33456 was a dimer with a molecular mass of 84 kDa. The pI was 4.66, and optimal enzyme activity was obtained at pH 6.5 and 37°C. The purified enzyme used both NADPH and NADH as electron donors for Cr(VI) reduction, while NADPH was the better, conferring 61% higher activity than NADH. The  $K_m$  values for NADPH and NADH were 47.5 and 17.2  $\mu\text{mol}$ , and the  $V_{\text{max}}$  values 322.2 and 130.7  $\mu\text{mol Cr(VI) min/mg protein}$ , respectively. The activity was strongly inhibited by *N*-ethylmaleimide,  $\text{Ag}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Hg}^{2+}$  and  $\text{Zn}^{2+}$  (Bae et al. 2005).

Chromate reductase from *P. putida* was purified by  $[\text{NH}_4]_2\text{SO}_4$  precipitation, anion exchange (DEAE sepharose CL-6B) and gel filtration (Sephadex-G-25) chromatography. The enzyme activity was dependent on NADH or NADPH, the temperature and pH optima were 75°C and 8.5, respectively, and the  $K_m$  was 260  $\mu\text{M}$  with a  $V_{\text{max}}$  of 8.8  $\mu\text{mol/min/mg}$  of protein. It had a molecular weight of 20 kDa (Renuga et al. 2008).

*Thermus scotoductus* SA-01, a South African gold mine isolate reduced Cr(VI) when grown aerobically in an organic medium containing 0.5 mM Cr(VI) (Opperman and Heerden 2007). Cell suspensions also reduced Cr(VI) under non-growth conditions. The reduction was found to be catalysed by a cytoplasmic, constitutively expressed enzyme. This enzyme was purified to homogeneity and was found to be a homodimeric protein, with a monomer molecular mass of 36 kDa, containing a non-covalently bound Flavin Mononucleotide cofactor. Enzyme activity was dependent on NADH or NADPH, with a preference for NADPH. The catalytic efficiency of chromate reduction was 50-fold greater than that of the quinone reductases and 180-fold more efficient than that of the nitroreductases. It was encoded by an open reading frame (ORF) of 1,050 bp, and was under the regulation of an *E. coli*  $\sigma^{70}$ -like promoter (Opperman and Heerden 2008). *T. scotoductus* also has a second strictly anaerobic membrane bound enzyme capable of chromate reduction. This enzyme showed a preference towards NADH. It was also a homodimer, but with a monomeric molecular weight of 48 kDa and a non-covalently bound FAD coenzyme. The catalytic efficiency of this enzyme was comparable to that of quinone reductases but more efficient than the nitroreductases. N-terminal sequencing and subsequent screening of a genomic library of *T. scotoductus* revealed an ORF of 1,386 bp, homologous (84%) to the dihydrolipoamide dehydrogenase gene of *T. thermophilus* HB8 (Opperman et al. 2008). These results extend the knowledge of chromate reductases mediating Cr(VI) reduction via noncovalently bound or free redox-active flavin groups and the activity of

dihydrolipoamide dehydrogenases towards physiologically unrelated substrates.

The kinetics of enzymatic reduction has also been studied using immobilized enzyme systems. In a recent work, chromate reductases associated with cell-free extracts of *A. rhombi*-RE were immobilized with different matrices and calcium alginate beads proved to be most effective (Elangovan et al. 2010). However, the performance was not encouraging under continuous mode of operation (Table 2).

#### Bioreactor studies for bacterial Cr(VI) remediation

Membrane bioreactors have been used in order to minimize the toxic effects of Cr(VI) on microorganisms or to facilitate the isolation of chromium that is reduced. Immobilized cells have also been used in such membrane reactors. One example is the use of *Pseudomonas* immobilized on agar–agar films for chromate reduction (Konovalova et al. 2003).

The relationship between sulphate starvation and chromate reduction was studied in a  $\text{H}_2$ -fed fixed-film bioreactor under continuous feed conditions. The bioreactor was inoculated with a bacterial consortium containing *Desulfomicrobium norvegicum*. With 500 mg/L of sulphate in the feed solution, total chromate-reduction was observed in the effluent whereas sulphate-reduction was strongly decreased. In the absence of sulphate, chromate-reduction activity was still observed but was lower than in the presence of sulphate.

In another study, columns packed with quartz sand alone and hydrous ferric oxide (HFO)-coated quartz sand were inoculated with *Cellulomonas* sp. ES6, fed nutrients to stimulate growth and then nutrient-free Cr(VI) solutions were injected into the column. In HFO-containing columns, Cr(VI) reduction was significantly prolonged and effluent Cr(VI) concentrations remained below detectable levels for periods of up to 66 residence times after nutrient injection was discontinued as against 5.7 residence times for columns packed with quartz sand alone. Fe was continuously detected in the effluent indicating that the insoluble Fe(III) bearing solids were being reduced to form soluble Fe(II) resulting in prolonged abiotic Cr(VI) reduction (Viamajala et al. 2008). Thus, growth of *Cellulomonas* resulted in formation of permeable reactive barriers that could reduce Cr(VI) and Fe(III) for extended periods even in the absence of external electron donors. This may be a viable technology for in situ remediation of Cr(VI)-contaminated groundwater.

Chromate reduction by *Arthrobacter* CR47 isolated from a landfarming soil sample was studied in batch operated packed bed biofilm reactors and in recirculating packed bed biofilm reactors. The reduction was evaluated

**Table 2** Characterization of chromate reductases from a few bacteria

Bacteria from which the chromate reductase is purified	Mol. Wt. (kDa)	Temp. optima (°C)	pH optima	$K_M$	$V_{max}$ (Cr(VI) reduced/min/mg protein)	References
<i>Bacillus</i> sp. QC1-2	44	37	7	0.35 mM	50 nM	Garcia et al. (1997)
<i>E. coli</i> ATCC 33456	84	37	6.5	47.5 $\mu$ M	322 $\mu$ M	Bae et al. (2005)
<i>Pseudomonas putida</i>	20	75	8.5	260 $\mu$ M	8.8 $\mu$ M	Renuga et al. (2008)
<i>Thermus scotoductus</i> SA-01 (soluble reductases)	72	65	6.3	8.4 $\mu$ M	16 $\mu$ M	Opperman and Heerden (2008)
<i>Thermus scotoductus</i> SA-01 (membrane bound reductase)	96	65	6.5	55.5 $\mu$ M	23 $\mu$ M	Opperman et al. (2008)
<i>Exiguobacterium</i> sp. ZM-2	–	–	–	106.1 $\mu$ M	1.24 $\mu$ M	Alam and Malik (2008)
<i>Bacillus</i> sp. KCH-3,	–	35	5.5	45–55 $\mu$ M	–	Saranghi and Krishnan (2008)
<i>Leucobacter</i> KCH-4	–	30	5.5	45–55 $\mu$ M	–	Saranghi and Krishnan (2008)
<i>Exiguobacterium</i> sp. KCH-5	–	35	6	200 $\mu$ M	–	Saranghi and Krishnan (2008)
<i>Bacillus</i> sp. GIDM-22	–	–	–	200 $\mu$ M	5.5 $\mu$ M	Desai et al. (2008)
<i>Arthrobacter rhombi</i> -RE	–	30	5.5	48 $\mu$ M	4.09 nM	Elangovan et al. (2010)

with laboratory solutions and industrial effluents containing 30 mg/L Cr(VI). Under batch mode, the reduction reaction by the biofilm fit an exponential-decay model with a first order kinetic parameter of 0.071 mg (L/h) Cr(VI). In the recirculating reactor fed with laboratory solutions, the removal rate was 0.79 mg/L/h. In the reactor fed with the industrial model solutions, the maximum Cr(VI) removal rate attained was 0.49 mg/L/h (Cordoba et al. 2008).

In yet another attempt, chromate reduction was carried out in a pilot-scale trickling filter packed with plastic media under continuous mode of operation with recirculation. Mixed bacterial cultures originated from industrial sludge were used for this study. The effect of the organic carbon concentration was examined for constant Cr(VI) influent concentration at about 5.5 mg/L and volumetric flow rates ranging from 60 to 900 mL/min. The highest reduction rate achieved was 1,117 g Cr(VI)/m<sup>2</sup>/day for a volumetric flow rate of 900 mL/min (Dermou and Vayenas 2008). Even though the system mandated frequent backwashing of the filter, the high reduction rates combined with low operating cost make this an option worth considering for the treatment of industrial chromate effluents.

Recently, activated sludge process was used for continuous Cr(VI) removal and the highest specific Cr(VI) removal rate was attained with cheese whey or lactose as electron donors. Batch assays with different nitrogen to carbon source ratios demonstrated that Cr(VI) reduction was associated with the cell multiplication phase; as a

result, maximum removal occurred when there was no substrate limitation (Orozco et al. 2010).

#### Bacterial Cr(VI) resistance

Cr(VI) resistant bacteria have been isolated from a number of environments and the genetic basis of resistance has been studied. Such studies have abundantly been carried out with tannery isolates. In one such work, among several isolates, all of which harboured plasmid DNA, *Corynebacterium hoagie* showed maximal resistance to 22 mM Cr(VI) (Viti et al. 2003). Thermotolerant strains resistant to Cr too have been isolated from tannery sources. These plasmid bearing isolates showed resistance to other heavy metals and antibiotics as well, which make them particularly useful. Three of the isolates belonged to uropathogenic *E. coli* serotype 04 (Verma et al. 2004). In another study, *B. brevis* showed good plasmid-mediated resistance and reduction potential as established by transfer and curing studies (Verma et al. 2009).

*Enterobacter cloacae* CYS-25, isolated from a chromate plant showed a strong tendency for chromate resistance due to reduced uptake and not reduction. Atomic force microscopic investigation showed that the average length of the bacterium in the stationary phase is about  $2.3 \pm 0.6 \mu$ M, while under the stimulation of 400 mg/L  $\text{CrO}_4^{2-}$ , the length increased to  $3.2 \pm 0.7 \mu$ M. Moreover, the smooth surface of the bacterium changed into one with discontinuous features. These compact convex patches are organic components stimulated by  $\text{CrO}_4^{2-}$  (Yang et al. 2007).



Hence, an over expression of extracellular biological components for preventing the permeability of  $\text{CrO}_4^{2-}$  into the cell helps the bacteria to survive chromate stress.

DNA helicases are involved in chromate resistance by *P. aeruginosa* PAO1. Chromate-hypersensitive mutants were generated using transposon insertion mutagenesis. Comparison of the nucleotide sequences of the regions interrupted within the PAO1 genome showed that mutant strains were affected in ORF PA0967 and PA5345, which correspond to the *ruvB* and *recG* genes, respectively. These genes encode helicases RuvB and RecG involved in DNA replication, recombination and repair. The chromate resistance phenotype was restored by cosmids bearing wild type *ruvB* or *recG* genes (Miranda et al. 2009). *P. aeruginosa* RuvB and RecG helicases are thus involved in repairing DNA damage caused by chromate.

The genome of *Arthrobacter* sp. FB24 is known to contain a chromate resistance determinant (CRD), consisting of a cluster of eight genes located on a 10.6 kb fragment of a 96 kb plasmid. The CRD includes *chrA*, which encodes a putative chromate efflux protein, and three genes with amino acid similarities to the amino and carboxyl termini of *ChrB*, a putative regulatory protein. There are also three novel genes that have not been previously associated with chromate resistance; they encode an oxidoreductase (most similar to malate:quinone oxidoreductase), a functionally unknown protein with a WD40 repeat domain and a lipoprotein. In one study, chromate-sensitive mutant (strain D11) of this bacterium was generated by curing FB24 of its 96-kb plasmid. The mutants accumulated three times more chromium than wild type cells. Introduction of the CRD into strain D11 conferred chromate resistance comparable to wild-type levels, whereas deletion of specific regions of the CRD led to decreased resistance. Using real-time reverse transcriptase PCR, the expression of each gene within the CRD was shown to be specifically induced in response to chromate. Higher levels of *chrA* expression were achieved when the *chrB* orthologs and the WD40 repeat domain genes were present, suggesting their possible regulatory roles (Henne et al. 2009). These findings indicate that chromate resistance in *Arthrobacter* FB24 is due to chromate efflux through the *ChrA* transport protein (Table 3).

#### Cr(VI) reduction by fungi, yeast and actinomycetes

Although a wide variety of bacterial cultures have been reported for their capability to reduce Cr(VI), little information exists on filamentous fungi (Pal. 1997) and yeasts (Muter et al. 2001; Ramirez-Ramirez et al. 2004). One such study was the Cr(VI) reduction by *Trichoderma viride* in stirred tank and concentric tube airlift bioreactors. The

**Table 3** Chromate resistance of bacterial isolates

Strain	Chromate conc. to which the isolate is resistant	References
<i>Bacillus brevis</i>	180 $\mu\text{M}$	Verma et al. (2009)
<i>Exiguobacterium</i> sp. KCH-5	1.5 mM	Sarangi and Krishnan (2008)
<i>Bacillus sphearicus</i>	15.3 mM	Pal and Paul (2004)
<i>Ochrobactrum</i> sp. CSCr-3	15.3 mM	He et al. (2009)
<i>Arthrobacter aurescens</i> P4	19.2 mM	Horton et al. (2006)
<i>Bacillus circulans</i> MNI	19.2 mM	Chaturvedi et al. (2007)
<i>Burkholderia cepacia</i> sp. MCMB-821	19.2 mM	Wani et al. (2007)
<i>Nesterenkonia</i> sp. MF-2	600 mM	Amoozegar et al. (2007)

airlift reactor performed better because the agitation in stirred tank bioreactor inflicted physical damage on the mycelia, causing their defragmentation (Morales-Barrera and Cristiani-Urbina 2005). *Lecytophora* sp. NGV-1, *Candida* sp. NGV-9 and *Aureobasidium pullulans* VR-8 isolated from tannery effluents and from a nickel-copper mine in Argentina showed tolerance to multiple heavy metals including Cr(VI). Growth in YNB-glucose medium supplemented with 0.5 and 1 mM Cr(VI) revealed that resistance was mainly due to reduction rather than bioaccumulation (Villegas et al. 2008).

Actinomycetes have also been screened for chromate reduction. *Streptomyces griseus* NCIM 2020 reduced Cr(VI) at the expense of a variety of electron donors. Sulphate, nitrate, chloride and carbonate had no effect on chromate reduction while cations such as Cd, Ni, Co and Cu were inhibitory. Cell-free extracts also reduced chromate and addition of NAD(P)H resulted in two- to three-fold increase in activity. Sonication was the best method to release the chromate reductase expressed constitutively. The enzyme showed optimum activity at 28°C and pH 7 (Poopal and Laxman 2009a). Different matrices were tested for whole cell immobilization of this chromate-reducing organism and PVA-alginate was found to be the most effective and was used in a bioreactor. Cr(VI) reduction efficiency decreased as Cr(VI) was increased from 2 to 12 mg/L but increased with an increase in biomass concentration. However, increasing the flow rate from 2 to 8 ml/h did not significantly affect Cr(VI) reduction (Poopal and Laxman 2009b). Such immobilized *S. griseus* cells could be applied for large-scale bioremediation of chromate-containing wastewaters.



## Chromium biosorption

Several biosorption studies have also been carried out for Cr. Although voluminous research has been carried out on this front, this paper takes only a fleeting glance at it, confining itself largely to biological reduction. Microorganisms such as bacteria (Srinath et al. 2002), fungi (Bai and Abraham 2003) and algae (Aravindhan et al. 2004) have been used in biosorption studies. The bacterium *P. aeruginosa* (Kang et al. 2006) fungus *Mucor hiemalis* (Tewari et al. 2005), yeast *Pichia guilliermondi* (Kaszycki et al. 2004) and microalgae *Scenedesmus incrassatulus* (Pena-Castro et al. 2004) are a few examples of organisms whose metal-binding capacities have been demonstrated.

As biosorption of metals using living biomass is time-consuming and it is expensive to find appropriate biomass, research interest turned to the use of dead microbial biomass and other biomaterials which are of low cost and are abundant. The removal of heavy metals by biomass by-products from agricultural, industrial or pharmaceutical industry has proved to be highly efficient. These biomaterials are of little commercial value and serve as substitutes for more expensive sources. They possess large quantity of adsorption sites (Law et al. 2003; Tarley and Arruda 2004) and functional groups for ion-exchange such as hydroxyl, carboxyl and phosphate groups as revealed by Fourier transform infrared spectrum (Law et al. 2003; Tarley and Arruda 2004; Chen et al. 2005). Cone biomass of *Pinus sylvestris* showed biosorption ability for Cr(VI) (Ucun et al. 2002). The agricultural biomass by-product *Lentinus edodes* also exhibited Cr(VI) biosorption potential. Nitrogen oxides and carboxyl groups were increased after adsorption and cation exchange was the underlying mechanism (Chen et al. 2006).

In another interesting revelation, chromate-tolerant bacteria isolated from the rhizosphere of aquatic plant *Eichhornia crassipes* enhanced Cr uptake by the plant. Plants inoculated with RA1 (*P. diminuta*), RA2 (*Brevundimonas diminuta*), RA3 (*Nitrobacteria irancium*), RA5 (*Ochrobactrum anthropi*), RA7 and RA8 (*B. cereus*) had 7-, 11-, 24-, 29-, 35- and 21-fold, respectively, higher Cr concentrations in roots compared to the control (Abou-Shanab et al. 2007). These bacterial isolates can play a supportive role in phytoremediation of Cr contaminants.

## Conclusion

Bacterial reduction of hexavalent Cr to its trivalent form is an option worth exploring for the remediation of diverse chromium contaminated sites. Its advantages over physical-chemical treatments are innumerable and research in this direction has come out with promising findings. Yet,

large scale field application of this technology warrants an improved understanding of the mechanisms involved in resistance and reduction of Cr(VI). Application of genetic and protein engineering tools to broaden enzyme activity should also be further researched. Enzyme purification, characterization and cloning of genes encoding chromate reductases are prerequisites for this, areas in which work has progressed well. Furthermore, successful implementation of a bioremediation program necessitates a thorough assessment of the contaminated site and choice of an appropriate technology as well as microbial strain that is best suited for the prevailing conditions.

## References

- Abou-Shanab RAI, Angle JS, Van Berkum P (2007) Chromate-tolerant bacteria for enhanced metal uptake by *Eichhornia crassipes* (MART). Int J Phytorem 9(2):91–105
- Alam MZ, Malik A (2008) Chromate resistance, transport and bioreduction by *Exiguobacterium* sp. ZM-2 isolated from agricultural soil irrigated with tannery effluent. J Basic Microbiol 48(5):416–420
- Amoozegar MA, Ghasemi A, Razavi MR, Naddaf S (2007) Evaluation of Hexavalent chromium reduction by chromate resistant moderate halophile *Nesterenkonia* sp. strain MF2. Process Biochem 42(10):1475–1479
- Anderson RA (1997) Chromium as an essential nutrient for humans. Regul Toxicol Pharmacol 26:S35–S41
- Aravindhan R, Madhan B, Rao JR (2004) Bioaccumulation of chromium from tannery wastewater: an approach for chrome recovery and reuse. Environ Sci Technol 38:300–306
- Bae WC, Lee HK, Choe YC, Jahng DJ, Lee SH, Kim SJ, Lee JH, Jeong BC (2005) Purification and characterization of NADPH-dependent Cr(VI) reductase from *Escherichia coli* ATCC 33456. J Microbiol 43(1):21–27
- Bai RS, Abraham TE (2003) Studies on Chromium(VI) adsorption-desorption using immobilized fungal biomass. Bioresour Technol 87:17–26
- Barak Y, Ackerley DF, Dodge CJ, Banwari L, Alex C, Francis AJ, Matin A (2006) Analysis of novel soluble chromate and uranyl reductase and generation of an improved enzyme by directed evolution. Appl Environ Microbiol 72(11):7074–7082
- Caravelli AH, Giannuzzi L, Zaritzky NE (2008) Reduction of hexavalent chromium by *Sphaerotilus natans* a filamentous microorganism present in activated sludges. J Hazard Mater 156(1–3):214–222
- Cervantes C, Campos-Garcia J, Devars S, Gutierrez-Corona F, Loza-Tavera H, Torres-Guzman JC, Moreno-Sanchez R (2000) Interactions of chromium. Am J Ind Med 38:127–131
- Chaturvedi MK, Bhatt S, Shukla NP (2007) Hexavalent chromium removal under anaerobic and aerobic conditions by *Bacillus circulans* strain MNI. Pollut Res 26(1):5–11
- Chen GQ, Zeng GM, Tu X (2005) A novel biosorbent: characterization of the spent mushroom compost and its application for removal of heavy metals. J Environ Sci 17:756–760
- Chen GQ, Zeng GM, Tu X, Niu CG, Huang GH, Jiang W (2006) Application of a by-product of *Lentinus edodes* to the bioremediation of chromate contaminated water. J Hazard Mater 135(1–3):249–255



- Cheung KH, Gu J (2007) Mechanism of hexavalent chromium detoxification by microorganisms and bioremediation application potential: a review. *Int Biodeter Biodegr* 59:8–15
- Cordoba A, Vargas P, Dussan J (2008) Chromate reduction by *Arthrobacter* CR47 in biofilm packed bed reactors. *J Hazard Mater* 151(1):274–279
- Daulton TL, Little BJ, Meehan J, Blom DA, Allard LF (2007) Microbial reduction of chromium from the hexavalent to the divalent state. *Geochem Cosmochim Acta* 71(3):557–565
- Dermou E, Vayenas DV (2008) Biological Cr(VI) reduction in a trickling filter under continuous operation with recirculation. *J Chem Technol Biotechnol* 83(6):871–877
- Desai C, Jain K, Madamwar D (2008) Evaluation of in vitro Cr(VI) reduction potential in cytosolic extracts of three indigenous *Bacillus* sp. isolated from Cr(VI) polluted industrial landfill. *Bioresour Technol* 99(14):6059–6069
- Elangovan R, Philip L, Chandraraj K (2010) Hexavalent chromium reduction by free and immobilized cell-free extract of *Arthrobacter rhombi*-RE. *Appl Biochem Biotechnol* 160(1):81–97
- Garcia J, Cadena G, Gonzalez R, Cervantes C (1997) Purification and characterization of a chromate reductase from *Bacillus*. *Rev Latinoam Microb* 39(1–2):73–81
- Gibb HJ, Lees PS, Pinsky FP, Rooney BC (2000a) Clinical findings of irritation among chromium chemical production workers. *Am J Ind Med* 38:127–131
- Gibb HJ, Lees PS, Pinsky FP, Rooney BC (2000b) Lung cancer among workers in chromium chemical production. *Am J Ind Med* 38:115–126
- Gonzalez CF, Ackerley DF, Lynch SV, Matin A (2005) ChrR, a soluble Quinone reductases of *Pseudomonas putida* that defends against H<sub>2</sub>O<sub>2</sub>. *J Biol Chem* 280(24):22590–22595
- He Z, Gao F, Sha T, Hu Y, He C (2009) Isolation and characterization of a Cr(VI) reduction *Ochrobactrum* sp. strain CSCr-3 from chromium landfill. *J Hazard Mater* 163(2–3):869–873
- Henne KL, Nakatsu CH, Thompson DK, Konopka AE (2009) High-level chromate resistance in *Arthrobacter* sp. strain FB24 requires previously uncharacterized accessory genes. *BMC Microbiol* 9:199
- Horton RN, Apel WA, Thompson VS, Sheridan PP (2006) Low temperature reduction of hexavalent chromium by a microbial consortium and a novel strain of *Arthrobacter aurescens*. *BMC Microbiol* 6:5
- Kang SY, Lee JU, Kim KW (2006) Biosorption of Cr(III) and Cr(VI) onto the cell surface of *Pseudomonas aeruginosa*. *Biochem Eng J* 36(1):54–58
- Kaszycki P, Fedorovych D, Ksheminska H, Babyak L, Wojcik D, Koloczek (2004) Chromium accumulation by living yeast at various environmental conditions. *Microbiol Res* 159:11–17
- Katz SA, Salem H (1994) The biological and environmental chemistry of chromium. The toxicology of chromium and its compound. VCH Publishers, Inc, New York
- Konovalova VV, Dmytrenko GM, Nigmatullin RR, Bryk MT, Gvozdyak PI (2003) Chromium (VI) reduction in a membrane bioreactor with immobilized *Pseudomonas* cells. *Enzyme Microb Technol* 33:899–907
- Krishnamurthy S, Wilkens MW (1994) Environmental chemistry of Cr. *Northeast Geol* 16(1):14–17
- Law WM, Lau WN, Lo KL (2003) Removal of biocide pentachlorophenol in water system by the spent mushroom compost of *Pleurotus pulmonarius*. *Chemosphere* 52:1531–1537
- Lee SE, Lee JU, Chon HT, Lee JS (2008) Reduction of Cr(VI) by indigenous bacteria in Cr-contaminated sediment under aerobic condition. *J Geochem Explor* 96(2–3):144–147
- Liu YG, Xu WH, Zeng GM, Li X, Gao H (2006) Cr (VI) reduction by *Bacillus* sp. isolated from chromium landfill. *Process Biochem* 41:1981–1986
- Miranda AT, Gonzalez MV, Gonzalez G, Vargas E, Campos-Garcia J, Cervantes C (2009) Involvement of DNA helicases in chromate resistance by *Pseudomonas aeruginosa* PA01. *Mut Res Fundam Mol Mech Mutagen* 578(1–2):202–209
- Morales-Barrera L, Cristiani-Urbina E (2005) Removal of hexavalent chromium by *Trichoderma viride* in an airlift bioreactor. *Enzyme Microb Technol* 40:107–113
- Muter O, Patmalnieks A, Rapoport A (2001) Interactions of the yeast *Candida utilis* and Cr (VI): metal reduction and its distribution in the cell and medium. *Process Biochem* 36:963–970
- Ohtake H, Silver S (1994) Bacterial detoxification of toxic chromate. In: Chaudry GR (ed) *Biological degradation and bioremediation of toxic chemicals*. Chapman and Hall, London
- Opperman DJ, Heerden V (2007) Aerobic Cr(VI) reduction by *Thermus scotoductus* strain SA-01. *J Appl Microbiol* 103(5):1907–1913
- Opperman DJ, Heerden V (2008) A membrane associated protein with Cr(VI)-reducing activity from *Thermus scotoductus*. *FEMS Microbiol Lett* 280(2):210–218
- Opperman DJ, Piater LA, Heerden V (2008) A novel chromate reductase from *Thermus scotoductus* SA-01 related to old yellow enzyme. *J Bacteriol* 190(8):3076–3082
- Orozco AMF, Contreras EM, Zaritsky NE (2010) Cr(VI) reduction capacity of activated sludge as affected by nitrogen and carbon sources, microbial acclimation and cell multiplication. *J Hazard Mater* 176(1–3):657–665
- Pal N (1997) Reduction of hexavalent chromium to trivalent chromium by *Phanerochaete chrysosporium*. In situ and on-site bioremediation, vol 1. Batelle Press, OH, pp 511–517
- Pal A, Paul AK (2004) Aerobic chromate reduction by chromium resistant bacteria isolated from serpentine soil. *Microbiol Res* 159:347–354
- Pattanapitipaisal P, Brown NL, Macaskie LE (2001) Chromate reduction and 16S rRNA identification of bacteria isolated from a Cr (VI) contaminated site. *Appl Microbiol Biotechnol* 57(1–2):257–261
- Pena-Castro JM, Martinez-Jeronimo F, Esparza-Garcia F, Canires-Villanueva RO (2004) Heavy metals removal by the microalga *Scenedesmus incrassatulus* in continuous cultures. *Bioresour Technol* 94:219–222
- Pesti M, Gazdag Z, Belagyi J (2000) In-vivo interaction of trivalent chromium with yeast plasma membrane as revealed by EPR spectroscopy. *FEMS Microbiol Lett* 182:375–380
- Poopal AC, Laxman RS (2009a) Chromate reduction by PVA alginate immobilized *Streptomyces griseus* in a bioreactor. *Biotechnol Lett* 31(1):71–76
- Poopal AC, Laxman RS (2009b) Studies on biological reduction of chromate by *Streptomyces griseus*. *J Hazard Mater* 169(1–3):539–549
- Puzon GJ, Peterson JN, Roberts AJ, Kramer DM, Xun L (2002) A bacterial flavin reductase system reduces chromate to a soluble chromium(III)-NAD<sup>+</sup> complex. *Biochem Biophys Res Commun* 294:76–81
- Puzon GJ, Robert AG, Kramer DM, Xun L (2005) Formation of soluble organo-Cr(III) complexes after chromate reduction in the presence of cellular organics. *Environ Sci Technol* 39(8):2811–2817
- Puzon GJ, Xun L, Tokala R, Zhang Z, Clark S, Peyton B, Yonge D (2008) Production and fate of organo-Cr(III) complexes from microbial reduction of chromate: evidence and implications. *AIHS-AISH Publ* 324:420–427
- Ramirez-Ramirez R, Calvo-Mendez C, Avila-Rodriguez M, Lappe P, Ulloa M, Vazquez-Juarez R (2004) Cr(VI) reduction in chromium resistant strain of *Candida maltosa* isolated from the leather industry. *Antonie van Leeuwenhoek* 85(1):63–68
- Renuga G, Lakshminarayan S, Babu A, Arumugam KR (2008) Purification of chromate reductase and their bioremediation potential. *Pollut Res* 27(1):129–136

- Romanenko VI, Korenkov VN (1977) A pure culture of bacterial cells assimilating chromates and bichromates as hydrogen acceptors when grown under anaerobic conditions. *Mikrobiologiya* 46:414–417
- Sarangi A, Krishnan C (2008) Comparison of in-vitro Cr(VI) reduction by CFEs of chromate resistant bacteria isolated from chromate contaminated soil. *Bioresour Technol* 99(10):4130–4137
- Saxena DK, Murthy RC, Jain VK, Chandra SV (1990) Fetoplacental-maternal uptake of hexavalent chromium administered orally in rats and mice. *Bull Environ Contam Toxicol* 45:430–435
- Silver S, Schottel J, Weiss A (2001) Bacterial resistance to toxic metals determined by extrachromosomal R factors. *Int Biodegrad Biodegrad* 48(1–4):263–281
- Srinath T, Verma T, Ramteke PW (2002) Cr(VI) biosorption and bioaccumulation by chromate resistant bacteria. *Chemosphere* 48:427–435
- Stewart DI, Burke IT, Mortimer RJG (2007) Stimulation of microbially mediated chromate reduction in alkaline soil-water systems. *Geomicrobiol J* 24(7–8):655–669
- Stewart DI, Burke IT, Hughes-Berry DV, Whittleston RA (2010) Microbially mediated chromate reduction in soil contaminated by highly alkaline leachate from chromium containing waste. *Ecol Eng* 36(2):211–221
- Tarley CRT, Arruda MAZ (2004) Biosorption of heavy metals using rice milling by-products: characterization and application for removal of metals from aqueous effluents. *Chemosphere* 54(7):987–995
- Tewari N, Vasudevan P, Guha BK (2005) Study on biosorption of Cr(VI) by *Mucor hiemalis*. *Biochem Eng J* 23:185–192
- Turpeinen R, Kairesalo T, Haggblom MM (2004) Microbial community structure and activity in arsenic, chromium and copper contaminated soils. *FEMS Microbiol Ecol* 47(1):39–50
- Ucun H, Bayhan YK, Kaya Y (2002) Biosorption of chromium(VI) from aqueous solution by cone biomass of *Pinus sylvestris*. *Bioresour Technol* 85:155–158
- Vatsouria A, Vainshtein M, Kuschik P, Wiessner A, Kosolapov D, Kaestner M (2005) Anaerobic co-reduction of chromate and nitrate by bacterial cultures of *Staphylococcus epidermidis* L-02. *J Ind Microbiol Biotechnol* 32(9):409–414
- Verma T, Garg SK, Ramteke PW (2009) Genetic correlation between chromium resistance and reduction in *Bacillus brevis* isolated from tannery effluent. *J Appl Microbiol* 107(5):1425–1432
- Verma T, Ramteke PW, Garg SK (2004) Occurrence of chromium resistant thermotolerant coliforms in tannery effluent. *Indian J Exp Biol* 42(11):1112–1116
- Viamajala S, Peyton BM, Gerlach R, Sivaswamy V, Apel WA, Peterson JN (2008) Permeable reactive biobarriers for in situ Cr(VI) reduction: bench scale tests using *Cellulomonas* sp. strain ES6. *Biotechnol Bioeng* 101(6):1150–1162
- Viamajala S, Smith WA, Sani RK, Apel WA, Peterson JN, Neal AL, Roberto FF, Peyton BM (2007) Isolation and characterization of Cr(VI) reducing *Cellulomonas* sp. from subsurface soils: implications for long-term chromate reduction. *Bioresour Technol* 98(3):612–622
- Villegas LB, Fernandez PM, Amoroso MJ, De Figueroa LIC (2008) Chromate removal by yeasts isolated from sediment of a tanning factory and a mine site in Argentina. *Biometals* 21(5):591–600
- Viti C, Mini A, Ranalli G, Lustrato G, Giovannetti L (2006) Response of microbial communities to different doses of chromate in soil microcosms. *Appl Soil Ecol* 34(2–3):125–139
- Viti C, Pace A, Giovannetti L (2003) Characterization of Cr(VI) resistant bacteria isolated from chromium-contaminated soil by tannery activity. *Curr Microbiol* 46(1):1–5
- Wang YT (2000) Microbial reduction of chromate. In: Loveley DR (ed) *Environmental microbe-metal interactions*. American Society for Microbiology Press, Washington DC
- Wani R, Kodam KM, Gawai KR, Dhakephalkar PK (2007) Chromate reduction by *Burkholderia cepacia* MCMB-821, isolated from the pristine habitat of alkaline crater lake. *Appl Microbiol Biotechnol* 75(3):627–632
- Yang C, Cheng Y, Ma X, Zhu Y, Wang C (2007) Surface-mediated chromate-resistant mechanism of *Enterobacter cloacae* bacteria investigated by atomic force microscopy. *Langmuir* 23(8):4480–4485
- Yang J, He M, Wang G (2009) Removal of toxic chromate using free and immobilized Cr(VI) reducing bacterial cells of *Intrasporangium* sp Q5-1. *World J Microbiol Biotechnol* 25(9):1579–1587
- Zhu WJ, Long HZ, Yang ZH, Chai LY (2008) Bio-reduction of Cr(VI) by *Leucobacter* and element analysis of reduction products. *J Cent South Univ Sci Technol* 39(3):443–447

