

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-(4-Chlorophenyl)-5-phenyl-1,2,4-triazolo[3,4-a]isoquinoline

F. Nawaz Khan,^a P. Manivel,^a K. Prabakaran,^a Venkatesha R. Hathwar^b and Mehmet Akkurt^{c*}

^aOrganic and Medicinal Chemistry Research Laboratory, Organic Chemistry Division, School of Advanced Sciences, VIT University, Vellore 632 014, Tamil Nadu, India, ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India, and ^cDepartment of Physics, Faculty of Arts and Sciences, Erciyes University, 38039 Kayseri, Turkey
Correspondence e-mail: akkurt@erciyes.edu.tr


Received 3 April 2010; accepted 13 April 2010

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.097; data-to-parameter ratio = 13.7.

In the title molecule, $\text{C}_{22}\text{H}_{14}\text{ClN}_3$, the triazoloisoquinoline ring system is approximately planar, with an r.m.s. deviation of 0.033 (2) Å and a maximum departure from the mean plane of 0.062 (1) Å for the triazole ring C atom, bonded to the benzene ring. The benzene and phenyl rings are twisted by 57.02 (6) and 62.16 (6)°, respectively, to the mean plane of the triazoloisoquinoline ring system. The molecule is stabilized by a weak intramolecular π - π interaction [centroid-centroid distance = 3.7089 (10) Å] between the benzene and phenyl rings. In the crystal structure, weak intermolecular C—H...N hydrogen bonds and C—H... π interactions link the molecules.

Related literature

For the synthesis and antihelmintic activity of triazolo compounds similar to the title compound, see: Nadkarni *et al.* (2001); Hui *et al.* (1999). For related structures, see: Khan *et al.* (2010); Zou *et al.* (2004).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{14}\text{ClN}_3$
 $M_r = 355.81$

Monoclinic, $P2_1/c$
 $a = 7.9841$ (3) Å

$b = 9.0679$ (4) Å
 $c = 23.9881$ (11) Å
 $\beta = 93.078$ (4)°
 $V = 1734.20$ (13) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.23$ mm⁻¹
 $T = 290$ K
 $0.40 \times 0.32 \times 0.25$ mm

Data collection

Oxford Xcalibur Eos (Nova) CCD detector diffractometer
Absorption correction: multi-scan (*CrysAlis PRO RED*; Oxford Diffraction, 2009)
 $T_{\min} = 0.902$, $T_{\max} = 0.945$

19413 measured reflections
3216 independent reflections
2090 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.097$
 $S = 0.97$
3216 reflections

235 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.11$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg_1 , Cg_2 and Cg_3 are the centroids of the N1–N3/C1/C16, N1/C1/C2/C7–C9 and C2–C7 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{N2}^{\text{i}}$	0.93	2.59	3.514 (2)	170
$\text{C8}-\text{H8}\cdots\text{N3}^{\text{i}}$	0.93	2.62	3.496 (2)	156
$\text{C18}-\text{H18}\cdots\text{Cg1}^{\text{ii}}$	0.93	2.70	3.4524 (17)	138
$\text{C21}-\text{H21}\cdots\text{Cg3}^{\text{iii}}$	0.93	2.89	3.7139 (19)	149
$\text{C22}-\text{H22}\cdots\text{Cg2}^{\text{iii}}$	0.93	2.90	3.5442 (18)	128

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis PRO CCD*; data reduction: *CrysAlis PRO RED* (Oxford Diffraction, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

We thank the FIST program for the data collection at SSCU, IISc, Bangalore. We also thank Professor T. N. Guru Row, IISc, Bangalore, for his help with the data collection. FNK thanks the DST for Fast Track Proposal funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2341).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Hui, X. P., Zhang, L. M. & Zhang, Z. Y. (1999). *Indian J. Chem. Sect. B*, **38**, 1066–1069.
Khan, F. N., Manivel, P., Prabakaran, K., Hathwar, V. R. & Ng, S. W. (2010). *Acta Cryst.* **E66**, o488.
Nadkarni, B. A., Kamat, V. R. & Khadse, B. G. (2001). *Arzneim. Forsch.* **51**, 569–573.
Oxford Diffraction (2009). *CrysAlis PRO CCD* and *CrysAlis PRO RED*. Oxford Diffraction Ltd, Yarnton, England.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Zou, K.-H., Cai, X.-Q., Chen, J.-X., Zhang, L.-X., Zhang, A.-J. & Hu, M.-L. (2004). *Acta Cryst.* **E60**, o1736–o1738.

supplementary materials

Acta Cryst. (2010). E66, o1094 [doi:10.1107/S1600536810013668]

3-(4-Chlorophenyl)-5-phenyl-1,2,4-triazolo[3,4-*a*]isoquinoline

F. N. Khan, P. Manivel, K. Prabakaran, V. R. Hathwar and M. Akkurt

Comment

As part of our search for new isoquinoline analogues, we focused on the synthesis of the titled compound, which crystal structure is reported.

In the title molecule (I), Fig. 1, the triazoloisoquinoline ring system (N1–N3/C1–C9/C16) is approximately planar, with an r.m.s. deviation of 0.033 (2) Å and a maximum departure from the mean plane of -0.062 (1) Å for the triazole ring C16 atom, bonded to the benzene ring (C17–C22). The benzene (C17–C22) and phenyl (C10–C15) rings are twisted by 57.02 (6) and 62.16 (6)° with respect to the mean plane of the triazoloisoquinoline ring system. The dihedral angle between the benzene (C17–C22) and phenyl (C10–C15) rings is 22.21 (8)°.

The molecule is stabilized by a weak intramolecular π - π interaction [$Cg4 \cdots Cg5(x, y, z) = 3.7089(10)$ Å; $Cg4$ and $Cg5$ are the centroids of the rings C10–C15 and C17–C22, respectively]. In the crystal structure, weak intermolecular C—H \cdots N hydrogen bonds and C—H \cdots π interactions (Table 1, Fig. 2) link the molecules to each other.

Experimental

2-(3-Phenylisoquinolin-1-yl)hydrazine (1 mmol) was condensed with, 4-chlorobenzaldehyde (1.1 mmol) under refluxing conditions in isopropanol (10 ml) solvent to give the corresponding hydrazone in high yield. After removal of the solvent the compound was then oxidatively cyclized in nitrobenzene (10 ml) at 473 K. The product was recrystallized from dichloromethane to give block-shaped crystals.

Refinement

H atoms were placed at calculated positions with C–H = 0.93 Å and were included in the refinement in the riding model approximation, with $U_{iso}(H) = 1.2U_{eq}(C)$.

Figures




Fig. 1. The title molecule with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level.




Fig. 2. A general view of the packing diagram and the hydrogen bonding of (I). H atoms not involved in the motif shown have been omitted for clarity.

3-(4-Chlorophenyl)-5-phenyl-1,2,4-triazolo[3,4-a]isoquinoline

Crystal data

$C_{22}H_{14}ClN_3$	$F(000) = 736$
$M_r = 355.81$	$D_x = 1.363 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 953 reflections
$a = 7.9841 (3) \text{ \AA}$	$\theta = 1.7\text{--}20.4^\circ$
$b = 9.0679 (4) \text{ \AA}$	$\mu = 0.23 \text{ mm}^{-1}$
$c = 23.9881 (11) \text{ \AA}$	$T = 290 \text{ K}$
$\beta = 93.078 (4)^\circ$	Block, colourless
$V = 1734.20 (13) \text{ \AA}^3$	$0.40 \times 0.32 \times 0.25 \text{ mm}$
$Z = 4$	

Data collection

Oxford Xcalibur Eos (Nova) CCD detector diffractometer	3216 independent reflections
Radiation source: Enhance (Mo) X-ray Source graphite	2090 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.042$
Absorption correction: multi-scan (CrysAlis Pro RED; Oxford Diffraction, 2009)	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 3.0^\circ$
$T_{\text{min}} = 0.902$, $T_{\text{max}} = 0.945$	$h = -9 \rightarrow 9$
19413 measured reflections	$k = -10 \rightarrow 10$
	$l = -29 \rightarrow 29$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 0.97$	$w = 1/[\sigma^2(F_o^2) + (0.0512P)^2]$
3216 reflections	where $P = (F_o^2 + 2F_c^2)/3$
235 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.11 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds

in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.31231 (8)	0.36657 (7)	0.47872 (2)	0.0881 (2)
N1	0.59862 (15)	0.38872 (12)	0.21077 (5)	0.0397 (3)
C9	0.76959 (18)	0.38731 (15)	0.23083 (7)	0.0420 (4)
C7	0.84736 (19)	0.37494 (16)	0.13340 (7)	0.0438 (4)
C8	0.8862 (2)	0.37965 (16)	0.19258 (7)	0.0465 (4)
H8	0.9984	0.3773	0.2051	0.056*
C2	0.67799 (19)	0.38170 (16)	0.11339 (7)	0.0428 (4)
C1	0.55381 (19)	0.39096 (16)	0.15404 (7)	0.0421 (4)
C10	0.81034 (19)	0.39579 (17)	0.29162 (7)	0.0434 (4)
C17	0.41623 (19)	0.38678 (16)	0.29626 (7)	0.0438 (4)
C16	0.44668 (19)	0.39497 (16)	0.23649 (7)	0.0435 (4)
N2	0.38951 (16)	0.40120 (14)	0.14584 (6)	0.0515 (4)
N3	0.32375 (16)	0.40415 (15)	0.19789 (6)	0.0514 (4)
C6	0.9712 (2)	0.36397 (18)	0.09438 (8)	0.0555 (5)
H6	1.0837	0.3616	0.1066	0.067*
C20	0.3518 (2)	0.3739 (2)	0.40853 (7)	0.0543 (5)
C18	0.4740 (2)	0.26909 (18)	0.32926 (7)	0.0484 (4)
H18	0.5349	0.1940	0.3133	0.058*
C3	0.6365 (2)	0.37670 (19)	0.05624 (8)	0.0577 (5)
H3	0.5248	0.3823	0.0432	0.069*
C19	0.4425 (2)	0.26216 (19)	0.38509 (7)	0.0536 (5)
H19	0.4819	0.1832	0.4068	0.064*
C21	0.2899 (2)	0.4902 (2)	0.37646 (8)	0.0608 (5)
H21	0.2271	0.5640	0.3923	0.073*
C5	0.9281 (2)	0.3568 (2)	0.03855 (8)	0.0668 (5)
H5	1.0115	0.3473	0.0132	0.080*
C11	0.7686 (2)	0.51875 (18)	0.32226 (7)	0.0519 (4)
H11	0.7176	0.5991	0.3042	0.062*
C22	0.3224 (2)	0.49554 (19)	0.32059 (8)	0.0541 (5)
H22	0.2806	0.5735	0.2989	0.065*
C15	0.8920 (2)	0.27925 (19)	0.31932 (7)	0.0552 (5)
H15	0.9245	0.1971	0.2993	0.066*
C12	0.8017 (2)	0.5234 (2)	0.37932 (8)	0.0627 (5)
H12	0.7719	0.6062	0.3994	0.075*
C4	0.7606 (3)	0.3635 (2)	0.01925 (8)	0.0705 (6)
H4	0.7329	0.3591	-0.0189	0.085*
C14	0.9250 (2)	0.2852 (2)	0.37642 (8)	0.0679 (6)
H14	0.9791	0.2067	0.3947	0.081*

supplementary materials

C13	0.8784 (3)	0.4062 (2)	0.40646 (8)	0.0692 (6)
H13	0.8988	0.4086	0.4450	0.083*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0944 (4)	0.1095 (5)	0.0616 (4)	0.0059 (3)	0.0163 (3)	0.0034 (3)
N1	0.0311 (7)	0.0376 (8)	0.0498 (8)	0.0006 (5)	-0.0026 (6)	-0.0017 (6)
C9	0.0333 (8)	0.0365 (9)	0.0552 (11)	-0.0003 (7)	-0.0060 (8)	-0.0025 (7)
C7	0.0389 (9)	0.0396 (10)	0.0527 (11)	-0.0010 (7)	-0.0007 (8)	-0.0034 (8)
C8	0.0324 (8)	0.0480 (10)	0.0579 (11)	0.0006 (7)	-0.0081 (8)	-0.0035 (8)
C2	0.0394 (9)	0.0379 (10)	0.0506 (11)	0.0012 (7)	-0.0031 (8)	-0.0051 (8)
C1	0.0369 (9)	0.0377 (10)	0.0507 (11)	0.0006 (7)	-0.0077 (8)	-0.0028 (7)
C10	0.0359 (8)	0.0429 (10)	0.0505 (10)	-0.0047 (7)	-0.0054 (7)	-0.0001 (8)
C17	0.0349 (8)	0.0376 (10)	0.0590 (11)	-0.0045 (7)	0.0047 (8)	0.0024 (8)
C16	0.0346 (8)	0.0368 (10)	0.0589 (11)	-0.0023 (7)	0.0011 (8)	0.0003 (8)
N2	0.0355 (8)	0.0589 (10)	0.0593 (10)	-0.0008 (6)	-0.0046 (7)	-0.0029 (7)
N3	0.0361 (7)	0.0562 (9)	0.0614 (10)	-0.0033 (6)	-0.0009 (7)	0.0018 (7)
C6	0.0410 (9)	0.0599 (12)	0.0654 (13)	0.0036 (8)	0.0017 (9)	-0.0061 (9)
C20	0.0513 (10)	0.0557 (12)	0.0565 (12)	-0.0062 (9)	0.0074 (9)	0.0003 (9)
C18	0.0453 (10)	0.0369 (10)	0.0634 (12)	0.0004 (7)	0.0058 (8)	-0.0005 (9)
C3	0.0483 (10)	0.0688 (13)	0.0549 (12)	0.0052 (9)	-0.0065 (9)	-0.0086 (9)
C19	0.0523 (11)	0.0455 (11)	0.0629 (12)	-0.0016 (8)	0.0012 (9)	0.0083 (9)
C21	0.0576 (11)	0.0516 (12)	0.0750 (14)	0.0062 (9)	0.0187 (10)	0.0001 (10)
C5	0.0571 (12)	0.0837 (14)	0.0604 (13)	0.0074 (10)	0.0111 (10)	-0.0074 (11)
C11	0.0538 (11)	0.0430 (10)	0.0582 (12)	-0.0032 (8)	-0.0030 (9)	-0.0006 (9)
C22	0.0489 (10)	0.0447 (11)	0.0695 (13)	0.0059 (8)	0.0103 (9)	0.0087 (9)
C15	0.0503 (11)	0.0506 (11)	0.0636 (12)	0.0043 (8)	-0.0075 (9)	0.0006 (9)
C12	0.0687 (12)	0.0608 (13)	0.0586 (13)	-0.0125 (10)	0.0018 (10)	-0.0137 (10)
C4	0.0704 (14)	0.0933 (16)	0.0473 (11)	0.0114 (11)	-0.0019 (10)	-0.0106 (10)
C14	0.0628 (12)	0.0744 (14)	0.0644 (14)	0.0050 (11)	-0.0154 (10)	0.0145 (11)
C13	0.0693 (13)	0.0853 (17)	0.0517 (12)	-0.0141 (12)	-0.0087 (10)	0.0011 (12)

Geometric parameters (\AA , $^\circ$)

C11—C20	1.7307 (18)	C20—C21	1.381 (2)
N1—C1	1.3886 (19)	C20—C19	1.383 (2)
N1—C16	1.3913 (19)	C18—C19	1.377 (2)
N1—C9	1.4228 (18)	C18—H18	0.9300
C9—C8	1.343 (2)	C3—C4	1.371 (3)
C9—C10	1.479 (2)	C3—H3	0.9300
C7—C6	1.401 (2)	C19—H19	0.9300
C7—C2	1.412 (2)	C21—C22	1.380 (2)
C7—C8	1.437 (2)	C21—H21	0.9300
C8—H8	0.9300	C5—C4	1.393 (3)
C2—C3	1.394 (2)	C5—H5	0.9300
C2—C1	1.430 (2)	C11—C12	1.381 (2)
C1—N2	1.3193 (19)	C11—H11	0.9300
C10—C11	1.386 (2)	C22—H22	0.9300

C10—C15	1.392 (2)	C15—C14	1.382 (2)
C17—C22	1.386 (2)	C15—H15	0.9300
C17—C18	1.393 (2)	C12—C13	1.374 (3)
C17—C16	1.469 (2)	C12—H12	0.9300
C16—N3	1.315 (2)	C4—H4	0.9300
N2—N3	1.3804 (19)	C14—C13	1.375 (3)
C6—C5	1.366 (2)	C14—H14	0.9300
C6—H6	0.9300	C13—H13	0.9300
C1—N1—C16	104.42 (12)	C19—C18—H18	119.5
C1—N1—C9	121.57 (13)	C17—C18—H18	119.5
C16—N1—C9	133.95 (14)	C4—C3—C2	119.83 (17)
C8—C9—N1	117.18 (14)	C4—C3—H3	120.1
C8—C9—C10	123.53 (14)	C2—C3—H3	120.1
N1—C9—C10	119.28 (14)	C18—C19—C20	119.27 (16)
C6—C7—C2	118.25 (15)	C18—C19—H19	120.4
C6—C7—C8	122.66 (15)	C20—C19—H19	120.4
C2—C7—C8	119.09 (15)	C22—C21—C20	119.20 (17)
C9—C8—C7	123.76 (14)	C22—C21—H21	120.4
C9—C8—H8	118.1	C20—C21—H21	120.4
C7—C8—H8	118.1	C6—C5—C4	120.67 (18)
C3—C2—C7	120.39 (16)	C6—C5—H5	119.7
C3—C2—C1	122.38 (14)	C4—C5—H5	119.7
C7—C2—C1	117.21 (14)	C12—C11—C10	120.77 (16)
N2—C1—N1	110.42 (15)	C12—C11—H11	119.6
N2—C1—C2	128.52 (15)	C10—C11—H11	119.6
N1—C1—C2	121.06 (13)	C21—C22—C17	121.20 (16)
C11—C10—C15	118.54 (15)	C21—C22—H22	119.4
C11—C10—C9	121.21 (14)	C17—C22—H22	119.4
C15—C10—C9	120.25 (14)	C14—C15—C10	120.20 (17)
C22—C17—C18	118.43 (16)	C14—C15—H15	119.9
C22—C17—C16	119.77 (15)	C10—C15—H15	119.9
C18—C17—C16	121.76 (15)	C13—C12—C11	120.20 (18)
N3—C16—N1	109.02 (14)	C13—C12—H12	119.9
N3—C16—C17	122.28 (14)	C11—C12—H12	119.9
N1—C16—C17	128.65 (14)	C3—C4—C5	120.23 (18)
C1—N2—N3	106.85 (13)	C3—C4—H4	119.9
C16—N3—N2	109.26 (13)	C5—C4—H4	119.9
C5—C6—C7	120.60 (16)	C13—C14—C15	120.54 (17)
C5—C6—H6	119.7	C13—C14—H14	119.7
C7—C6—H6	119.7	C15—C14—H14	119.7
C21—C20—C19	120.84 (17)	C12—C13—C14	119.70 (18)
C21—C20—C11	119.56 (15)	C12—C13—H13	120.2
C19—C20—C11	119.60 (14)	C14—C13—H13	120.2
C19—C18—C17	121.02 (16)		
C1—N1—C9—C8	-3.74 (19)	C18—C17—C16—N1	-55.3 (2)
C16—N1—C9—C8	179.68 (14)	N1—C1—N2—N3	-0.67 (16)
C1—N1—C9—C10	175.62 (12)	C2—C1—N2—N3	178.84 (14)
C16—N1—C9—C10	-1.0 (2)	N1—C16—N3—N2	1.22 (16)

supplementary materials

N1—C9—C8—C7	0.9 (2)	C17—C16—N3—N2	-176.41 (12)
C10—C9—C8—C7	-178.46 (13)	C1—N2—N3—C16	-0.35 (16)
C6—C7—C8—C9	-178.68 (15)	C2—C7—C6—C5	-1.4 (2)
C2—C7—C8—C9	1.4 (2)	C8—C7—C6—C5	178.64 (15)
C6—C7—C2—C3	0.3 (2)	C22—C17—C18—C19	-1.5 (2)
C8—C7—C2—C3	-179.71 (13)	C16—C17—C18—C19	-179.15 (14)
C6—C7—C2—C1	179.22 (14)	C7—C2—C3—C4	0.7 (2)
C8—C7—C2—C1	-0.8 (2)	C1—C2—C3—C4	-178.11 (16)
C16—N1—C1—N2	1.37 (15)	C17—C18—C19—C20	0.1 (2)
C9—N1—C1—N2	-176.09 (12)	C21—C20—C19—C18	1.3 (3)
C16—N1—C1—C2	-178.19 (13)	C11—C20—C19—C18	-179.73 (12)
C9—N1—C1—C2	4.4 (2)	C19—C20—C21—C22	-1.3 (3)
C3—C2—C1—N2	-2.6 (2)	C11—C20—C21—C22	179.77 (13)
C7—C2—C1—N2	178.59 (14)	C7—C6—C5—C4	1.5 (3)
C3—C2—C1—N1	176.91 (13)	C15—C10—C11—C12	-2.3 (2)
C7—C2—C1—N1	-1.9 (2)	C9—C10—C11—C12	177.63 (15)
C8—C9—C10—C11	116.02 (18)	C20—C21—C22—C17	-0.2 (3)
N1—C9—C10—C11	-63.30 (19)	C18—C17—C22—C21	1.6 (2)
C8—C9—C10—C15	-64.0 (2)	C16—C17—C22—C21	179.24 (15)
N1—C9—C10—C15	116.67 (16)	C11—C10—C15—C14	2.2 (2)
C1—N1—C16—N3	-1.56 (15)	C9—C10—C15—C14	-177.82 (15)
C9—N1—C16—N3	175.42 (14)	C10—C11—C12—C13	0.7 (3)
C1—N1—C16—C17	175.88 (14)	C2—C3—C4—C5	-0.7 (3)
C9—N1—C16—C17	-7.1 (2)	C6—C5—C4—C3	-0.4 (3)
C22—C17—C16—N3	-55.7 (2)	C10—C15—C14—C13	-0.3 (3)
C18—C17—C16—N3	121.87 (17)	C11—C12—C13—C14	1.2 (3)
C22—C17—C16—N1	127.15 (17)	C15—C14—C13—C12	-1.4 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1, Cg2 and Cg3 are the centroids of the N1–N3/C1/C16, N1/C1/C2/C7–C9 and C2–C7 rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6 \cdots N2 ⁱ	0.93	2.59	3.514 (2)	170
C8—H8 \cdots N3 ⁱ	0.93	2.62	3.496 (2)	156
C18—H18 \cdots Cg1 ⁱⁱ	0.93	2.70	3.4524 (17)	138
C21—H21 \cdots Cg3 ⁱⁱⁱ	0.93	2.89	3.7139 (19)	149
C22—H22 \cdots Cg2 ⁱⁱⁱ	0.93	2.90	3.5442 (18)	128

Symmetry codes: (i) $x+1, y, z$; (ii) $-x+1, y-1/2, -z+1/2$; (iii) $-x+1, y+1/2, -z+1/2$.

Fig. 1





Fig. 2



Copyright of Acta Crystallographica: Section E (International Union of Crystallography - IUCr) is the property of International Union of Crystallography - IUCr and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.