

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4,5,6,7,8,9-Hexahydro-2*H*-cycloocta- [c]pyrazol-1-ium-3-olate

 Hoong-Kun Fun,^{a*}‡ Chin Sing Yeap,^{a§} R. Venkat
Ragavan,^b V. Vijayakumar^b and S. Sarveswari^b
^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bOrganic Chemistry Division, School of Advanced Sciences, VIT University, Vellore 632 014, India

Correspondence e-mail: hkfun@usm.my


Received 19 October 2010; accepted 27 October 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.037; wR factor = 0.095; data-to-parameter ratio = 14.4.

The title compound, $\text{C}_9\text{H}_{14}\text{N}_2\text{O}$, exists in the zwitterionic form in the crystal. The cyclooctane ring adopts a twisted boat-chair conformation. In the crystal, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into sheets lying parallel to the bc plane. The structure is also stabilized by $\pi-\pi$ interactions, with a centroid-to-centroid distance of 3.5684 (8) Å.

Related literature

For pyrazole derivatives and their microbial activities, see: Ragavan *et al.* (2009, 2010). For a related structure, see: Xiong *et al.* (2007). For the stability of the temperature controller used for data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

 $\text{C}_9\text{H}_{14}\text{N}_2\text{O}$
 $M_r = 166.22$

 Monoclinic, $P2_1/c$
 $a = 12.8078$ (2) Å
 $b = 6.7758$ (1) Å
 $c = 10.7096$ (2) Å
 $\beta = 111.620$ (1)°
 $V = 864.03$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100$ K
 $0.54 \times 0.24 \times 0.11$ mm

Data collection

 Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2009)
 $T_{\min} = 0.956$, $T_{\max} = 0.991$

 6990 measured reflections
 1680 independent reflections
 1474 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.095$
 $S = 1.05$
 1680 reflections
 117 parameters

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.25$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N1}\cdots\text{O1}^{\text{i}}$	0.938 (19)	1.757 (19)	2.6900 (14)	173.0 (19)
$\text{N2}-\text{H1N2}\cdots\text{O1}^{\text{ii}}$	0.925 (19)	1.789 (19)	2.7056 (14)	170.1 (18)

 Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and CSY thank Universiti Sains Malaysia (USM) for a Research University Grant (No. 1001/PFIZIK/811160). VV is grateful to DST-India for funding through the Young Scientist Scheme (Fast Track Proposal).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2356).

References

- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Ragavan, R. V., Vijayakumar, V. & Kumari, N. S. (2009). *Eur. J. Med. Chem.* **44**, 3852–3857.
- Ragavan, R. V., Vijayakumar, V. & Kumari, N. S. (2010). *Eur. J. Med. Chem.* **45**, 1173–1180.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Xiong, Y., Gao, W.-Y., Deng, K.-Z., Chen, H.-X. & Wang, S.-J. (2007). *Acta Cryst.* **E63**, o333–o334.

‡ Thomson Reuters ResearcherID: A-3561-2009.

§ Thomson Reuters ResearcherID: A-5523-2009.

supplementary materials

Acta Cryst. (2010). E66, o3019 [doi:10.1107/S1600536810043904]

4,5,6,7,8,9-Hexahydro-2*H*-cycloocta[*c*]pyrazol-1-ium-3-olate

H.-K. Fun, C. S. Yeap, R. V. Ragavan, V. Vijayakumar and S. Sarveswari

Comment

Antibacterial and antifungal activities of the azoles are most widely studied and some of them are in clinical practice as anti-microbial agents. However, the azole-resistant strains led to the development of new antimicrobial compounds. In particular pyrazole derivatives are extensively studied and used as antimicrobial agents. Pyrazole is an important class of heterocyclic compounds and many pyrazole derivatives are reported to have a broad spectrum of biological properties such as anti-inflammatory, antifungal, herbicidal, anti-tumour, cytotoxic, molecular modelling and antiviral activities. Pyrazole derivatives also act as antiangiogenic agents, A3 adenosine receptor antagonists, neuropeptide YY5 receptor antagonists, kinase inhibitor for treatment of type 2 diabetes, hyperlipidemia, obesity, and thrombopietinmimetics. Recently urea derivatives of pyrazoles have been reported as potent inhibitors of p38 kinase. Since the high electronegativity of halogens (particularly chlorine and fluorine) in the aromatic part of the drug molecules play an important role in enhancing their biological activity, we are interested to have 4-fluoro or 4-chloro substitution in the aryls of 1,5-diaryl pyrazoles. As part of our on-going research aiming on the synthesis of new antimicrobial compounds, we have reported the synthesis of novel pyrazole derivatives and their microbial activities (Ragavan *et al.*, 2009, 2010).

The title compound exists in an zwitterionic form (Fig. 1). The cyclooctane ring adopts a twisted boat-chair conformation which similar to Xiong *et al.* (2007). In the crystal structure, intermolecular N1—H1N1···O1 and N2—H1N2···O1 hydrogen bonds link the molecules into planes parallel to the *bc* plane (Fig. 2). The structure is stabilized by the π - π interactions [$Cg1 \cdots Cg1^{iii} = 3.5684(8) \text{ \AA}$; $Cg1$ is centroid of N1–N2–C1–C8–C9 ring; (iii) $1 - x, 1 - y, 1 - z$].

Experimental

The compound has been synthesized using the method available in the literature Ragavan *et al.*, (2010) and recrystallized using the ethanol–chloroform 1:1 mixture. Yield: 74%, m.p. 221.6–228.8 °C.

Refinement

The N-bound H atoms were located from difference Fourier map and refined freely. The rest of H atoms were positioned geometrically [$C-H = 0.97 \text{ \AA}$] and refined using a riding model [$U_{iso}(H) = 1.2U_{eq}$].

Figures




Fig. 1. The molecular structure of the title compound with atom labels and 50% probability ellipsoids for non-H atoms.




Fig. 2. The crystal packing of title compound, viewed down *b* axis, showing the molecules are linked into planes parallel to the *bc* plane. Intermolecular hydrogen bonds are shown as dashed lines.

4,5,6,7,8,9-Hexahydro-2*H*-cycloocta[*c*]pyrazol-1-ium-3-olate

Crystal data

$C_9H_{14}N_2O$

$M_r = 166.22$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.8078$ (2) Å

$b = 6.7758$ (1) Å

$c = 10.7096$ (2) Å

$\beta = 111.620$ (1)°

$V = 864.03$ (2) Å³

$Z = 4$

$F(000) = 360$

$D_x = 1.278$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3802 reflections

$\theta = 3.5$ – 30.1 °

$\mu = 0.09$ mm⁻¹

$T = 100$ K

Plate, colourless

$0.54 \times 0.24 \times 0.11$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

Absorption correction: multi-scan (*SADABS*; Bruker, 2009)

$T_{\min} = 0.956$, $T_{\max} = 0.991$

6990 measured reflections

1680 independent reflections

1474 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 26.0$ °, $\theta_{\min} = 1.7$ °

$h = -15 \rightarrow 15$

$k = -8 \rightarrow 8$

$l = -13 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.037$

$wR(F^2) = 0.095$

$S = 1.05$

1680 reflections

117 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.047P)^2 + 0.3516P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.25$ e Å⁻³

0 restraints

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.41579 (8)	0.06094 (13)	0.31883 (8)	0.0190 (2)
N1	0.44456 (9)	0.21181 (16)	0.52370 (10)	0.0160 (3)
N2	0.41423 (9)	0.38447 (16)	0.56789 (11)	0.0154 (3)
C1	0.34257 (10)	0.48051 (18)	0.46046 (12)	0.0149 (3)
C2	0.28502 (11)	0.66563 (19)	0.47482 (13)	0.0177 (3)
H2A	0.2711	0.7476	0.3961	0.021*
H2B	0.3335	0.7384	0.5525	0.021*
C3	0.17257 (11)	0.6209 (2)	0.49123 (12)	0.0189 (3)
H3A	0.1883	0.5734	0.5818	0.023*
H3B	0.1306	0.7430	0.4807	0.023*
C4	0.09909 (11)	0.4691 (2)	0.39200 (12)	0.0185 (3)
H4A	0.0295	0.4555	0.4075	0.022*
H4B	0.1371	0.3426	0.4113	0.022*
C5	0.07002 (11)	0.5164 (2)	0.24164 (12)	0.0191 (3)
H5A	-0.0105	0.5025	0.1954	0.023*
H5B	0.0888	0.6534	0.2339	0.023*
C6	0.12881 (11)	0.38840 (19)	0.16907 (12)	0.0189 (3)
H6A	0.0905	0.4065	0.0731	0.023*
H6B	0.1202	0.2510	0.1889	0.023*
C7	0.25452 (11)	0.42986 (19)	0.20469 (12)	0.0177 (3)
H7A	0.2795	0.3581	0.1424	0.021*
H7B	0.2640	0.5695	0.1918	0.021*
C8	0.32877 (10)	0.37496 (19)	0.34518 (12)	0.0153 (3)
C9	0.39626 (10)	0.20329 (18)	0.38724 (12)	0.0150 (3)
H1N1	0.4969 (14)	0.125 (3)	0.5823 (17)	0.030 (4)*
H1N2	0.4229 (14)	0.397 (3)	0.6572 (19)	0.037 (5)*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0272 (5)	0.0181 (5)	0.0123 (4)	0.0062 (4)	0.0082 (4)	0.0006 (3)
N1	0.0201 (6)	0.0158 (5)	0.0122 (5)	0.0031 (4)	0.0061 (4)	0.0007 (4)
N2	0.0181 (6)	0.0167 (5)	0.0122 (5)	0.0012 (4)	0.0065 (4)	-0.0012 (4)
C1	0.0149 (6)	0.0163 (6)	0.0150 (6)	-0.0015 (5)	0.0072 (5)	0.0013 (5)
C2	0.0214 (7)	0.0155 (6)	0.0166 (6)	-0.0001 (5)	0.0076 (5)	-0.0017 (5)
C3	0.0204 (7)	0.0207 (7)	0.0165 (6)	0.0025 (5)	0.0078 (5)	-0.0021 (5)
C4	0.0179 (7)	0.0212 (7)	0.0182 (6)	0.0003 (5)	0.0090 (5)	-0.0014 (5)
C5	0.0179 (7)	0.0217 (7)	0.0163 (7)	0.0025 (5)	0.0045 (5)	-0.0006 (5)
C6	0.0223 (7)	0.0203 (7)	0.0130 (6)	0.0034 (5)	0.0051 (5)	0.0004 (5)
C7	0.0227 (7)	0.0192 (6)	0.0126 (6)	0.0053 (5)	0.0082 (5)	0.0035 (5)
C8	0.0165 (6)	0.0168 (6)	0.0144 (6)	0.0001 (5)	0.0078 (5)	0.0012 (5)
C9	0.0168 (6)	0.0172 (6)	0.0118 (6)	-0.0001 (5)	0.0061 (5)	0.0011 (5)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.2902 (15)	C4—C5	1.5467 (17)
N1—C9	1.3622 (16)	C4—H4A	0.9700
N1—N2	1.3708 (15)	C4—H4B	0.9700
N1—H1N1	0.939 (18)	C5—C6	1.5343 (17)
N2—C1	1.3459 (16)	C5—H5A	0.9700
N2—H1N2	0.925 (19)	C5—H5B	0.9700
C1—C8	1.3807 (17)	C6—C7	1.5377 (18)
C1—C2	1.4912 (17)	C6—H6A	0.9700
C2—C3	1.5438 (17)	C6—H6B	0.9700
C2—H2A	0.9700	C7—C8	1.5007 (17)
C2—H2B	0.9700	C7—H7A	0.9700
C3—C4	1.5283 (18)	C7—H7B	0.9700
C3—H3A	0.9700	C8—C9	1.4199 (17)
C3—H3B	0.9700		
C9—N1—N2	109.39 (10)	H4A—C4—H4B	107.4
C9—N1—H1N1	128.4 (10)	C6—C5—C4	115.83 (11)
N2—N1—H1N1	121.9 (10)	C6—C5—H5A	108.3
C1—N2—N1	107.96 (10)	C4—C5—H5A	108.3
C1—N2—H1N2	128.7 (11)	C6—C5—H5B	108.3
N1—N2—H1N2	119.5 (11)	C4—C5—H5B	108.3
N2—C1—C8	109.65 (11)	H5A—C5—H5B	107.4
N2—C1—C2	121.73 (11)	C5—C6—C7	115.82 (11)
C8—C1—C2	128.51 (11)	C5—C6—H6A	108.3
C1—C2—C3	111.34 (10)	C7—C6—H6A	108.3
C1—C2—H2A	109.4	C5—C6—H6B	108.3
C3—C2—H2A	109.4	C7—C6—H6B	108.3
C1—C2—H2B	109.4	H6A—C6—H6B	107.4
C3—C2—H2B	109.4	C8—C7—C6	114.98 (10)
H2A—C2—H2B	108.0	C8—C7—H7A	108.5

C4—C3—C2	114.47 (10)	C6—C7—H7A	108.5
C4—C3—H3A	108.6	C8—C7—H7B	108.5
C2—C3—H3A	108.6	C6—C7—H7B	108.5
C4—C3—H3B	108.6	H7A—C7—H7B	107.5
C2—C3—H3B	108.6	C1—C8—C9	106.16 (11)
H3A—C3—H3B	107.6	C1—C8—C7	126.38 (11)
C3—C4—C5	115.79 (11)	C9—C8—C7	127.44 (11)
C3—C4—H4A	108.3	O1—C9—N1	122.31 (11)
C5—C4—H4A	108.3	O1—C9—C8	130.92 (11)
C3—C4—H4B	108.3	N1—C9—C8	106.76 (11)
C5—C4—H4B	108.3		
C9—N1—N2—C1	2.98 (13)	C2—C1—C8—C9	-175.54 (12)
N1—N2—C1—C8	-2.13 (13)	N2—C1—C8—C7	178.99 (11)
N1—N2—C1—C2	174.25 (11)	C2—C1—C8—C7	2.9 (2)
N2—C1—C2—C3	-89.09 (14)	C6—C7—C8—C1	-77.60 (16)
C8—C1—C2—C3	86.55 (15)	C6—C7—C8—C9	100.55 (14)
C1—C2—C3—C4	-46.15 (14)	N2—N1—C9—O1	176.18 (11)
C2—C3—C4—C5	-55.64 (15)	N2—N1—C9—C8	-2.61 (13)
C3—C4—C5—C6	108.08 (13)	C1—C8—C9—O1	-177.37 (13)
C4—C5—C6—C7	-72.87 (15)	C7—C8—C9—O1	4.2 (2)
C5—C6—C7—C8	68.15 (15)	C1—C8—C9—N1	1.29 (13)
N2—C1—C8—C9	0.52 (14)	C7—C8—C9—N1	-177.16 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N1...O1 ⁱ	0.938 (19)	1.757 (19)	2.6900 (14)	173.0 (19)
N2—H1N2...O1 ⁱⁱ	0.925 (19)	1.789 (19)	2.7056 (14)	170.1 (18)

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $x, -y+1/2, z+1/2$.

Fig. 1





Fig. 2



Copyright of Acta Crystallographica: Section E (International Union of Crystallography - IUCr) is the property of International Union of Crystallography - IUCr and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.