

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,3-Dimethyl-2,6-diphenylpiperidin-4-one

 P. Nithya,^a Venkatesha R. Hathwar,^b T. Maiyalagan,^a Canan Kazak^c and F. Nawaz Khan^{a*}
^aChemistry Division, School of Science and Humanities, VIT University, Vellore 632 014, Tamil Nadu, India, ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India, and ^cOndokuz Mayıs University, Arts and Sciences Faculty, Department of Physics, 55139 Samsun, Turkey

Correspondence e-mail: nawaz_f@yahoo.co.in


Received 22 January 2009; accepted 28 January 2009

 Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.040; wR factor = 0.116; data-to-parameter ratio = 16.4.

In the title molecule, $\text{C}_{19}\text{H}_{21}\text{NO}$, the 4-piperidone ring adopts a chair conformation in which the two benzene rings and the methyl group attached to C atoms all have equatorial orientations. In the crystal structure, centrosymmetric dimers are formed through weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds [the dihedral angle between the aromatic rings is $58.51(5)^\circ$].

Related literature

For general background, see: Badorrey *et al.* (1999); Grishina *et al.* (1994); Nalanishi *et al.* (1974); Perumal *et al.* (2001); Ponnuswamy *et al.* (2002). For a related crystal structure, see: Gayathri *et al.* (2008). For the synthesis, see: Noller & Baliah (1948). For puckering and asymmetry parameters, see: Cremer & Pople (1975); Nardelli (1983, 1995).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{21}\text{NO}$
 $M_r = 279.37$

 Triclinic, $P\bar{1}$
 $a = 5.9201(2)$ Å

 $b = 10.9749(3)$ Å

 $c = 12.8247(3)$ Å

 $\alpha = 80.2961(12)^\circ$
 $\beta = 86.673(2)^\circ$
 $\gamma = 76.4499(11)^\circ$
 $V = 798.30(4)$ Å³
 $Z = 2$

 Mo $K\alpha$ radiation

 $\mu = 0.07$ mm⁻¹
 $T = 290(2)$ K

 $0.28 \times 0.21 \times 0.18$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.943$, $T_{\max} = 0.987$

12316 measured reflections

3143 independent reflections

 2446 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.116$
 $S = 1.02$

3143 reflections

192 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.14$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C1}-\text{H1}\cdots\text{O1}^i$	0.98	2.56	3.3535 (16)	139

 Symmetry code: (i) $-x, -y + 1, -z + 2$.

Data collection: SMART (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXL97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2003); software used to prepare material for publication: PLATON.

We thank the Department of Science and Technology, India, for use of the CCD facility set up under the IRHPA-DST program at the IISc. We thank Professor T. N. Guru Row, IISc, Bangalore, for useful crystallographic discussions. FNK thanks the DST for Fast Track Proposal funding.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2761).

References

- Badorrey, R., Cativiela, C., Diaz-de-Villegas, M. D. & Galvez, J. A. (1999). *Tetrahedron*, **55**, 7601–7612.
- Bruker (2004). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gayathri, P., Thiruvalluvar, A., Manimekalai, A., Sivakumar, S. & Butcher, R. J. (2008). *Acta Cryst.* **E64**, o1973.
- Grishina, G. V., Gaidarova, E. L. & Zefirov, N. S. (1994). *Chem. Heterocycl. Compd.* **30**, 401–426.
- Nalanishi, M., Shiraki, M., Kobayakawa, T. & Kobayashi, R. (1974). Jpn Patent 74-03987.
- Nardelli, M. (1983). *Acta Cryst.* **C39**, 1141–1142.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Noller, C. & Baliah, V. (1948). *J. Am. Chem. Soc.* **70**, 3853–3855.
- Perumal, R. V., Agiraj, M. & Shanmugapandiyam, P. (2001). *Indian Drugs*, **38**, 156–159.
- Ponnuswamy, S., Venkatraj, M., Jeyaraman, R., Suresh Kumar, M., Kumaran, D. & Ponnuswamy, M. N. (2002). *Indian J. Chem. Sect. B*, **41**, 614–627.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2009). E65, o439 [doi:10.1107/S1600536809003419]

1,3-Dimethyl-2,6-diphenylpiperidin-4-one

P. Nithya, Venkatesha R. Hathwar, T. Maiyalagan, Canan Kazak and F. Nawaz Khan

S1. Comment

The synthesis of 4-piperidones is of current interest due to their potential medical applications (Grishina *et al.*, 1994, Ponnuswamy *et al.*, 2002). 4-Piperidones have been found to exhibit blood cholesterol-lowering activities (Nalanishi *et al.*, 1974). Various piperidones and piperidine derivatives are present in numerous alkaloids (Badorrey *et al.*, 1999). Piperidones are also reported to possess analgesic, anti-inflammatory, central nervous system (CNS), local anaesthetic, anticancer and antimicrobial activity (Perumal *et al.*, 2001).


In the title molecule, C₁₉H₂₀NO (Fig. 1), the piperidine ring adopts a chair conformation (Cremer & Pople, 1975; Nardelli, 1995). The phenyl rings at positions 2 and 6 and the methyl group attached at position 3 all have equatorial orientations. In the related crystal structure of r-2,c-6-Bis(4-chlorophenyl)-t-3-isopropyl-1-nitrosopiperidin-4-one, the piperidine ring also adopts a chair conformation (Gayathri *et al.*, 2008) but the three substituents on the C atoms of the ring are in axial orientations. In the crystal structure, centrosymmetric dimers are formed through weak intermolecular C—H...O hydrogen bonds (Fig. 2).

S2. Experimental


The synthesis was based on a procedure in the literature (Noller & Baliah, 1948). Benzaldehyde (0.20 mol), 3-methyl-2-butanone (0.10 mol) and ammonium acetate (0.10 mol) were dissolved in 80 ml of distilled ethanol and heated over a boiling water bath, with shaking until a yellow colour developed and ultimately changed to orange. The solution was left undisturbed for 14 h. The precipitated solid was filtered and purified by recrystallization from ethanol. The piperidone intermediate was then dissolved in acetone and was alkylated with methyl iodide in the presence of potassium carbonate.

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H bond lengths of 0.93, 0.97 and 0.96 Å for aromatic, methylene and methyl H atoms, respectively and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C})$ for methyl bound H atoms.

**Figure 1**

The molecular structure of (I) shown with 50% probability displacement ellipsoids.

**Figure 2**

Part of the crystal structure of (I) with dashed lines indicating intermolecular C—H...O hydrogen bonds.

1,3-Dimethyl-2,6-diphenylpiperidin-4-one

Crystal data

C₁₉H₂₁NO $M_r = 279.37$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 5.9201$ (2) Å $b = 10.9749$ (3) Å $c = 12.8247$ (3) Å $\alpha = 80.2961$ (12)° $\beta = 86.673$ (2)° $\gamma = 76.4499$ (11)° $V = 798.30$ (4) Å³ $Z = 2$ $F(000) = 300$ $D_x = 1.162$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 873 reflections

 $\theta = 1.9$ – 20.8 ° $\mu = 0.07$ mm⁻¹ $T = 290$ K

Block, colorless

 $0.28 \times 0.21 \times 0.18$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.943$, $T_{\max} = 0.987$

12316 measured reflections

3143 independent reflections

2446 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.019$ $\theta_{\text{max}} = 26.0$ °, $\theta_{\text{min}} = 1.6$ ° $h = -6 \rightarrow 7$ $k = -13 \rightarrow 13$ $l = -15 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.116$ $S = 1.02$

3143 reflections

192 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0539P)^2 + 0.1409P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.14$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.17010 (18)	0.40892 (9)	0.73876 (8)	0.0419 (3)
O1	0.25711 (17)	0.47960 (11)	1.03053 (7)	0.0615 (3)
C1	0.1134 (2)	0.54013 (12)	0.76264 (10)	0.0419 (3)

H1	-0.0457	0.5582	0.7916	0.050*
C2	0.2788 (3)	0.55376 (14)	0.84570 (11)	0.0532 (4)
H2A	0.4336	0.5475	0.8146	0.064*
H2B	0.2276	0.6370	0.8665	0.064*
C3	0.2882 (2)	0.45443 (14)	0.94154 (10)	0.0485 (3)
C4	0.3367 (3)	0.32119 (14)	0.91755 (11)	0.0532 (4)
H4	0.4923	0.3030	0.8850	0.064*
C5	0.1607 (2)	0.31444 (12)	0.83478 (10)	0.0432 (3)
H5	0.0040	0.3339	0.8661	0.052*
C6	0.2066 (2)	0.18211 (12)	0.80603 (10)	0.0467 (3)
C7	0.4060 (3)	0.13480 (14)	0.75020 (12)	0.0571 (4)
H7	0.5153	0.1837	0.7319	0.069*
C8	0.4446 (3)	0.01556 (15)	0.72124 (14)	0.0711 (5)
H8	0.5799	-0.0148	0.6840	0.085*
C9	0.2867 (4)	-0.05793 (15)	0.74677 (15)	0.0748 (5)
H9	0.3126	-0.1374	0.7261	0.090*
C10	0.0903 (4)	-0.01377 (16)	0.80298 (17)	0.0791 (5)
H10	-0.0170	-0.0639	0.8214	0.095*
C11	0.0495 (3)	0.10595 (15)	0.83308 (14)	0.0656 (4)
H11	-0.0846	0.1348	0.8717	0.079*
C12	0.1264 (2)	0.63589 (12)	0.66384 (10)	0.0428 (3)
C13	-0.0613 (3)	0.73494 (13)	0.63339 (12)	0.0536 (4)
H13	-0.1987	0.7419	0.6733	0.064*
C14	-0.0475 (3)	0.82404 (14)	0.54407 (13)	0.0658 (4)
H14	-0.1752	0.8902	0.5245	0.079*
C15	0.1535 (3)	0.81493 (15)	0.48471 (13)	0.0675 (5)
H15	0.1629	0.8751	0.4252	0.081*
C16	0.3413 (3)	0.71672 (17)	0.51319 (13)	0.0687 (5)
H16	0.4776	0.7101	0.4725	0.082*
C17	0.3284 (3)	0.62776 (15)	0.60201 (12)	0.0579 (4)
H17	0.4565	0.5616	0.6207	0.070*
C18	0.0031 (3)	0.39902 (14)	0.66175 (12)	0.0623 (4)
H18A	-0.1504	0.4132	0.6928	0.093*
H18B	0.0439	0.3158	0.6422	0.093*
H18C	0.0069	0.4616	0.5999	0.093*
C19	0.3365 (4)	0.22361 (19)	1.01678 (14)	0.0942 (7)
H19A	0.4369	0.2368	1.0679	0.141*
H19B	0.3910	0.1398	0.9993	0.141*
H19C	0.1814	0.2326	1.0458	0.141*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0505 (6)	0.0408 (5)	0.0341 (6)	-0.0089 (5)	-0.0065 (4)	-0.0056 (4)
O1	0.0534 (6)	0.0987 (8)	0.0390 (6)	-0.0236 (5)	0.0017 (4)	-0.0217 (5)
C1	0.0439 (7)	0.0439 (7)	0.0377 (7)	-0.0076 (5)	-0.0009 (5)	-0.0097 (5)
C2	0.0664 (9)	0.0558 (8)	0.0426 (8)	-0.0190 (7)	-0.0068 (6)	-0.0132 (6)
C3	0.0408 (7)	0.0708 (9)	0.0369 (7)	-0.0152 (6)	-0.0045 (5)	-0.0120 (6)

C4	0.0567 (8)	0.0602 (8)	0.0391 (7)	-0.0083 (7)	-0.0098 (6)	-0.0021 (6)
C5	0.0426 (7)	0.0471 (7)	0.0378 (7)	-0.0091 (5)	0.0007 (5)	-0.0032 (5)
C6	0.0483 (8)	0.0455 (7)	0.0437 (7)	-0.0102 (6)	-0.0050 (6)	0.0006 (6)
C7	0.0632 (9)	0.0483 (8)	0.0588 (9)	-0.0138 (7)	0.0082 (7)	-0.0069 (7)
C8	0.0846 (12)	0.0519 (9)	0.0729 (11)	-0.0073 (8)	0.0069 (9)	-0.0134 (8)
C9	0.0964 (14)	0.0446 (8)	0.0832 (12)	-0.0136 (9)	-0.0167 (10)	-0.0081 (8)
C10	0.0831 (13)	0.0558 (9)	0.1021 (14)	-0.0323 (9)	-0.0123 (11)	0.0036 (10)
C11	0.0572 (9)	0.0586 (9)	0.0795 (11)	-0.0193 (7)	0.0017 (8)	0.0004 (8)
C12	0.0511 (7)	0.0410 (6)	0.0388 (7)	-0.0118 (5)	-0.0038 (5)	-0.0105 (5)
C13	0.0614 (9)	0.0448 (7)	0.0522 (8)	-0.0046 (6)	-0.0024 (6)	-0.0112 (6)
C14	0.0908 (12)	0.0407 (7)	0.0607 (10)	-0.0037 (7)	-0.0145 (9)	-0.0052 (7)
C15	0.1055 (14)	0.0528 (9)	0.0486 (9)	-0.0318 (9)	-0.0065 (9)	0.0003 (7)
C16	0.0750 (11)	0.0822 (11)	0.0516 (9)	-0.0312 (9)	0.0058 (8)	-0.0018 (8)
C17	0.0536 (9)	0.0651 (9)	0.0509 (9)	-0.0106 (7)	0.0006 (6)	-0.0018 (7)
C18	0.0852 (11)	0.0505 (8)	0.0540 (9)	-0.0167 (7)	-0.0292 (8)	-0.0049 (7)
C19	0.146 (2)	0.0797 (12)	0.0527 (11)	-0.0258 (12)	-0.0310 (11)	0.0125 (9)

Geometric parameters (Å, °)

N1—C18	1.4711 (16)	C9—C10	1.365 (3)
N1—C5	1.4777 (15)	C9—H9	0.9300
N1—C1	1.4800 (15)	C10—C11	1.395 (2)
O1—C3	1.2128 (15)	C10—H10	0.9300
C1—C12	1.5145 (18)	C11—H11	0.9300
C1—C2	1.5363 (18)	C12—C13	1.3820 (19)
C1—H1	0.9800	C12—C17	1.390 (2)
C2—C3	1.493 (2)	C13—C14	1.387 (2)
C2—H2A	0.9700	C13—H13	0.9300
C2—H2B	0.9700	C14—C15	1.367 (2)
C3—C4	1.503 (2)	C14—H14	0.9300
C4—C19	1.519 (2)	C15—C16	1.372 (2)
C4—C5	1.5520 (18)	C15—H15	0.9300
C4—H4	0.9800	C16—C17	1.381 (2)
C5—C6	1.5175 (18)	C16—H16	0.9300
C5—H5	0.9800	C17—H17	0.9300
C6—C11	1.383 (2)	C18—H18A	0.9600
C6—C7	1.384 (2)	C18—H18B	0.9600
C7—C8	1.384 (2)	C18—H18C	0.9600
C7—H7	0.9300	C19—H19A	0.9600
C8—C9	1.364 (3)	C19—H19B	0.9600
C8—H8	0.9300	C19—H19C	0.9600
C18—N1—C5	109.46 (10)	C8—C9—C10	119.37 (16)
C18—N1—C1	108.63 (10)	C8—C9—H9	120.3
C5—N1—C1	111.85 (9)	C10—C9—H9	120.3
N1—C1—C12	111.33 (10)	C9—C10—C11	120.50 (16)
N1—C1—C2	110.45 (10)	C9—C10—H10	119.7
C12—C1—C2	109.68 (10)	C11—C10—H10	119.7

N1—C1—H1	108.4	C6—C11—C10	120.59 (16)
C12—C1—H1	108.4	C6—C11—H11	119.7
C2—C1—H1	108.4	C10—C11—H11	119.7
C3—C2—C1	112.04 (11)	C13—C12—C17	118.07 (13)
C3—C2—H2A	109.2	C13—C12—C1	120.95 (12)
C1—C2—H2A	109.2	C17—C12—C1	120.97 (12)
C3—C2—H2B	109.2	C12—C13—C14	120.86 (15)
C1—C2—H2B	109.2	C12—C13—H13	119.6
H2A—C2—H2B	107.9	C14—C13—H13	119.6
O1—C3—C2	122.76 (13)	C15—C14—C13	120.18 (15)
O1—C3—C4	123.25 (13)	C15—C14—H14	119.9
C2—C3—C4	113.99 (11)	C13—C14—H14	119.9
C3—C4—C19	112.28 (13)	C14—C15—C16	119.84 (15)
C3—C4—C5	108.87 (11)	C14—C15—H15	120.1
C19—C4—C5	112.70 (13)	C16—C15—H15	120.1
C3—C4—H4	107.6	C15—C16—C17	120.21 (16)
C19—C4—H4	107.6	C15—C16—H16	119.9
C5—C4—H4	107.6	C17—C16—H16	119.9
N1—C5—C6	110.17 (10)	C16—C17—C12	120.83 (14)
N1—C5—C4	110.93 (10)	C16—C17—H17	119.6
C6—C5—C4	110.79 (10)	C12—C17—H17	119.6
N1—C5—H5	108.3	N1—C18—H18A	109.5
C6—C5—H5	108.3	N1—C18—H18B	109.5
C4—C5—H5	108.3	H18A—C18—H18B	109.5
C11—C6—C7	117.95 (13)	N1—C18—H18C	109.5
C11—C6—C5	121.26 (13)	H18A—C18—H18C	109.5
C7—C6—C5	120.78 (12)	H18B—C18—H18C	109.5
C6—C7—C8	120.81 (15)	C4—C19—H19A	109.5
C6—C7—H7	119.6	C4—C19—H19B	109.5
C8—C7—H7	119.6	H19A—C19—H19B	109.5
C9—C8—C7	120.77 (17)	C4—C19—H19C	109.5
C9—C8—H8	119.6	H19A—C19—H19C	109.5
C7—C8—H8	119.6	H19B—C19—H19C	109.5
C18—N1—C1—C12	-59.74 (14)	N1—C5—C6—C7	55.01 (16)
C5—N1—C1—C12	179.33 (10)	C4—C5—C6—C7	-68.12 (16)
C18—N1—C1—C2	178.16 (11)	C11—C6—C7—C8	0.9 (2)
C5—N1—C1—C2	57.22 (13)	C5—C6—C7—C8	-178.08 (14)
N1—C1—C2—C3	-52.14 (15)	C6—C7—C8—C9	0.2 (3)
C12—C1—C2—C3	-175.21 (11)	C7—C8—C9—C10	-1.1 (3)
C1—C2—C3—O1	-127.65 (13)	C8—C9—C10—C11	0.8 (3)
C1—C2—C3—C4	51.65 (16)	C7—C6—C11—C10	-1.1 (2)
O1—C3—C4—C19	1.3 (2)	C5—C6—C11—C10	177.82 (14)
C2—C3—C4—C19	-178.02 (14)	C9—C10—C11—C6	0.3 (3)
O1—C3—C4—C5	126.78 (13)	N1—C1—C12—C13	124.49 (13)
C2—C3—C4—C5	-52.51 (15)	C2—C1—C12—C13	-112.96 (14)
C18—N1—C5—C6	56.39 (14)	N1—C1—C12—C17	-56.44 (15)
C1—N1—C5—C6	176.84 (10)	C2—C1—C12—C17	66.11 (15)

C18—N1—C5—C4	179.44 (11)	C17—C12—C13—C14	-0.4 (2)
C1—N1—C5—C4	-60.11 (13)	C1—C12—C13—C14	178.66 (12)
C3—C4—C5—N1	55.96 (14)	C12—C13—C14—C15	0.0 (2)
C19—C4—C5—N1	-178.79 (13)	C13—C14—C15—C16	0.5 (2)
C3—C4—C5—C6	178.65 (11)	C14—C15—C16—C17	-0.5 (2)
C19—C4—C5—C6	-56.09 (17)	C15—C16—C17—C12	0.0 (2)
N1—C5—C6—C11	-123.92 (14)	C13—C12—C17—C16	0.5 (2)
C4—C5—C6—C11	112.95 (15)	C1—C12—C17—C16	-178.65 (13)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C1—H1...O1 ⁱ	0.98	2.56	3.3535 (16)	139

Symmetry code: (i) $-x, -y+1, -z+2$.