

## 2-Chloro-6-methylquinoline-3-carbaldehyde

F. Nawaz Khan,<sup>a</sup> R. Subashini,<sup>a</sup> S. Mohana Roopan,<sup>a</sup>  
Venkatesha R. Hathwar<sup>b</sup> and Seik Weng Ng<sup>c\*</sup>

<sup>a</sup>Chemistry Division, School of Science and Humanities, VIT University, Vellore 632 014, Tamil Nadu, India, <sup>b</sup>Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 6 October 2009; accepted 6 October 2009

Key indicators: single-crystal X-ray study;  $T = 290\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.034;  $wR$  factor = 0.089; data-to-parameter ratio = 16.0.

The quinolinyl fused-ring of the title compound,  $\text{C}_{11}\text{H}_8\text{ClNO}$ , is almost planar (r.m.s. deviation =  $0.013\text{ \AA}$ ); the formyl group is slightly bent out of the plane of the fused ring system [ $\text{C}-\text{C}-\text{O}$  torsion angle =  $13.5(4)^\circ$ ].

### Related literature

For a review of the synthesis of quinolines by the Vilsmeier–Haack reaction, see: Meth-Cohn (1993).



### Experimental

#### Crystal data

$\text{C}_{11}\text{H}_8\text{ClNO}$

$M_r = 205.63$

Monoclinic, $Pc$	$Z = 2$
$a = 5.944(1)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 3.9210(19)\text{ \AA}$	$\mu = 0.37\text{ mm}^{-1}$
$c = 20.390(2)\text{ \AA}$	$T = 290\text{ K}$
$\beta = 101.377(15)^\circ$	$0.25 \times 0.15 \times 0.15\text{ mm}$
$V = 465.9(2)\text{ \AA}^3$	

#### Data collection

Oxford Diffraction Excalibur diffractometer	5980 measured reflections
Absorption correction: multi-scan ( <i>CrysAlis Pro</i> ; Oxford Diffraction, 2009)	2052 independent reflections
	1831 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.028$
	$T_{\min} = 0.913$ , $T_{\max} = 0.947$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H-atom parameters constrained
$wR(F^2) = 0.089$	$\Delta\rho_{\max} = 0.19\text{ e \AA}^{-3}$
$S = 1.00$	$\Delta\rho_{\min} = -0.17\text{ e \AA}^{-3}$
2052 reflections	Absolute structure: Flack (1983),
128 parameters	990 Friedel pairs
2 restraints	Flack parameter: 0.02 (6)

Data collection: *CrysAlis Pro* (Oxford Diffraction, 2009); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis Pro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Department of Science and Technology, India, for use of the diffraction facility at IISc under the IRHPA-DST program. FNK thanks the DST for Fast Track Proposal funding. We thank VIT University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2550).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Meth-Cohn, O. (1993). *Heterocycles*, **35**, 539–557.
- Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

# supporting information

*Acta Cryst.* (2009). E65, o2686 [doi:10.1107/S1600536809040653]

## 2-Chloro-6-methylquinoline-3-carbaldehyde


F. Nawaz Khan, R. Subashini, S. Mohana Roopan, Venkatesha R. Hathwar and Seik Weng Ng

### S1. Experimental

A Vilsmeier-Haack adduct prepared from phosphorus oxytrichloride (6.5 ml, 70 mmol) and *N,N*-dimethylformamide (2.3 ml, 30 mmol) at 273 K was added to *N*-(4-tolyl)acetamide (1.49 g, 10 mmol), and heated at 353 K for 15 h. The mixture was then poured onto ice, and the white product was collected and dried. The compound was purified by recrystallization from a petroleum ether/ethyl acetate mixture.

### S2. Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93–0.96 Å) and were included in the refinement in the riding model approximation with  $U_{\text{iso}}(\text{H})$  set to 1.2–1.5  $U_{\text{eq}}(\text{C})$ .



**Figure 1**

Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{11}\text{H}_8\text{ClNO}$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

## 2-Chloro-6-methylquinoline-3-carbaldehyde

### Crystal data

$\text{C}_{11}\text{H}_8\text{ClNO}$   
 $M_r = 205.63$   
Monoclinic,  $Pc$   
Hall symbol: P -2yc

$a = 5.944 (1)$  Å  
 $b = 3.9210 (19)$  Å  
 $c = 20.390 (2)$  Å  
 $\beta = 101.377 (15)^\circ$

$V = 465.9 (2) \text{ \AA}^3$   
 $Z = 2$   
 $F(000) = 212$   
 $D_x = 1.466 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
Cell parameters from 1352 reflections

$\theta = 2.0\text{--}20.7^\circ$   
 $\mu = 0.37 \text{ mm}^{-1}$   
 $T = 290 \text{ K}$   
Block, colorless  
 $0.25 \times 0.15 \times 0.15 \text{ mm}$

#### Data collection

Oxford Diffraction Excalibur diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
 $\omega$  scans  
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2009)  
 $T_{\min} = 0.913$ ,  $T_{\max} = 0.947$

5980 measured reflections  
2052 independent reflections  
1831 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.5^\circ$   
 $h = -7 \rightarrow 7$   
 $k = -5 \rightarrow 5$   
 $l = -26 \rightarrow 25$

#### Refinement

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.089$   
 $S = 1.00$   
2052 reflections  
128 parameters  
2 restraints  
Primary atom site location: structure-invariant direct methods  
Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0584P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.19 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$   
Absolute structure: Flack (1983), 990 Friedel pairs  
Absolute structure parameter: 0.02 (6)

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	1.00002 (8)	1.11653 (14)	0.50000 (3)	0.04862 (17)
O1	0.4626 (4)	0.5444 (6)	0.55814 (8)	0.0676 (6)
N1	0.7865 (3)	0.9885 (5)	0.37983 (9)	0.0382 (4)
C1	0.7703 (3)	0.9492 (5)	0.44201 (10)	0.0351 (5)
C2	0.5905 (3)	0.7858 (5)	0.46557 (10)	0.0340 (4)
C3	0.4137 (4)	0.6644 (5)	0.41837 (10)	0.0341 (4)
H3	0.2906	0.5561	0.4315	0.041*
C4	0.4163 (3)	0.7021 (5)	0.34984 (10)	0.0320 (4)
C5	0.2378 (4)	0.5831 (5)	0.29839 (10)	0.0366 (4)
H5	0.1096	0.4792	0.3094	0.044*
C6	0.2519 (4)	0.6196 (5)	0.23214 (10)	0.0383 (5)
C7	0.4490 (4)	0.7786 (6)	0.21674 (10)	0.0454 (5)
H7	0.4598	0.8019	0.1721	0.054*
C8	0.6212 (4)	0.8971 (6)	0.26419 (11)	0.0441 (5)
H8	0.7474	1.0015	0.2521	0.053*
C9	0.6099 (4)	0.8627 (5)	0.33289 (10)	0.0335 (4)
C10	0.5895 (4)	0.7365 (7)	0.53783 (11)	0.0470 (5)
H10	0.6930	0.8615	0.5688	0.056*

C11	0.0664 (4)	0.4932 (7)	0.17678 (10)	0.0501 (6)
H11A	-0.0398	0.3562	0.1952	0.075*
H11B	0.1332	0.3584	0.1463	0.075*
H11C	-0.0131	0.6841	0.1534	0.075*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0366 (3)	0.0578 (3)	0.0488 (3)	-0.0066 (3)	0.00193 (19)	-0.0056 (3)
O1	0.0607 (12)	0.1037 (15)	0.0375 (9)	-0.0252 (12)	0.0074 (8)	0.0170 (10)
N1	0.0348 (9)	0.0388 (8)	0.0428 (10)	-0.0031 (7)	0.0118 (7)	-0.0002 (7)
C1	0.0309 (11)	0.0349 (11)	0.0389 (11)	0.0019 (8)	0.0056 (8)	-0.0021 (8)
C2	0.0325 (11)	0.0377 (10)	0.0325 (9)	0.0060 (9)	0.0083 (8)	0.0028 (8)
C3	0.0310 (10)	0.0375 (10)	0.0354 (11)	0.0007 (8)	0.0106 (8)	0.0031 (8)
C4	0.0327 (10)	0.0302 (10)	0.0337 (10)	0.0022 (8)	0.0077 (8)	0.0012 (8)
C5	0.0346 (11)	0.0376 (11)	0.0375 (10)	-0.0002 (8)	0.0067 (8)	0.0003 (8)
C6	0.0412 (12)	0.0384 (11)	0.0348 (10)	0.0030 (9)	0.0062 (9)	-0.0016 (8)
C7	0.0575 (15)	0.0519 (12)	0.0294 (10)	0.0032 (11)	0.0148 (10)	0.0014 (9)
C8	0.0480 (13)	0.0471 (12)	0.0421 (11)	-0.0048 (10)	0.0207 (10)	0.0018 (10)
C9	0.0355 (10)	0.0326 (10)	0.0337 (10)	0.0021 (8)	0.0099 (8)	0.0003 (8)
C10	0.0412 (13)	0.0631 (14)	0.0350 (10)	-0.0026 (11)	0.0035 (9)	0.0021 (11)
C11	0.0573 (15)	0.0555 (13)	0.0346 (11)	-0.0024 (12)	0.0019 (10)	-0.0041 (10)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Cl1—C1	1.748 (2)	C5—H5	0.9300
O1—C10	1.196 (3)	C6—C7	1.416 (3)
N1—C1	1.300 (2)	C6—C11	1.498 (3)
N1—C9	1.365 (3)	C7—C8	1.345 (3)
C1—C2	1.409 (3)	C7—H7	0.9300
C2—C3	1.363 (3)	C8—C9	1.422 (3)
C2—C10	1.487 (3)	C8—H8	0.9300
C3—C4	1.408 (3)	C10—H10	0.9300
C3—H3	0.9300	C11—H11A	0.9600
C4—C9	1.414 (3)	C11—H11B	0.9600
C4—C5	1.416 (3)	C11—H11C	0.9600
C5—C6	1.377 (3)		
C1—N1—C9	116.50 (17)	C8—C7—C6	122.5 (2)
N1—C1—C2	126.42 (19)	C8—C7—H7	118.7
N1—C1—Cl1	114.64 (15)	C6—C7—H7	118.7
C2—C1—Cl1	118.93 (14)	C7—C8—C9	119.9 (2)
C3—C2—C1	116.68 (17)	C7—C8—H8	120.0
C3—C2—C10	120.06 (19)	C9—C8—H8	120.0
C1—C2—C10	123.25 (19)	N1—C9—C4	122.69 (17)
C2—C3—C4	120.41 (18)	N1—C9—C8	118.51 (19)
C2—C3—H3	119.8	C4—C9—C8	118.81 (19)
C4—C3—H3	119.8	O1—C10—C2	123.4 (2)

C3—C4—C9	117.27 (17)	O1—C10—H10	118.3
C3—C4—C5	123.20 (18)	C2—C10—H10	118.3
C9—C4—C5	119.53 (17)	C6—C11—H11A	109.5
C6—C5—C4	120.73 (19)	C6—C11—H11B	109.5
C6—C5—H5	119.6	H11A—C11—H11B	109.5
C4—C5—H5	119.6	C6—C11—H11C	109.5
C5—C6—C7	118.4 (2)	H11A—C11—H11C	109.5
C5—C6—C11	121.8 (2)	H11B—C11—H11C	109.5
C7—C6—C11	119.78 (19)		
C9—N1—C1—C2	1.0 (3)	C5—C6—C7—C8	-0.6 (3)
C9—N1—C1—Cl1	-179.69 (15)	C11—C6—C7—C8	-180.0 (2)
N1—C1—C2—C3	-1.7 (3)	C6—C7—C8—C9	0.4 (3)
Cl1—C1—C2—C3	179.00 (15)	C1—N1—C9—C4	0.9 (3)
N1—C1—C2—C10	177.0 (2)	C1—N1—C9—C8	-179.45 (19)
Cl1—C1—C2—C10	-2.3 (3)	C3—C4—C9—N1	-2.0 (3)
C1—C2—C3—C4	0.5 (3)	C5—C4—C9—N1	178.91 (17)
C10—C2—C3—C4	-178.30 (18)	C3—C4—C9—C8	178.38 (18)
C2—C3—C4—C9	1.2 (3)	C5—C4—C9—C8	-0.7 (3)
C2—C3—C4—C5	-179.73 (19)	C7—C8—C9—N1	-179.4 (2)
C3—C4—C5—C6	-178.44 (18)	C7—C8—C9—C4	0.2 (3)
C9—C4—C5—C6	0.6 (3)	C3—C2—C10—O1	13.5 (4)
C4—C5—C6—C7	0.0 (3)	C1—C2—C10—O1	-165.1 (3)
C4—C5—C6—C11	179.4 (2)		