

- Loh, W.-S., Fun, H.-K., Sarveswari, S., Vijayakumar, V. & Reddy, B. P. (2010). *Acta Cryst.* **E66**, o91–o92.
- Markees, D. G., Dewey, V. C. & Kidder, G. W. (1970). *J. Med. Chem.* **13**, 324–326.
- Michael, J. P. (1997). *Nat. Prod. Rep.* **14**, 605–608.
- Shahani, T., Fun, H.-K., Sarveswari, S., Vijayakumar, V. & Ragavan, R. V. (2010). *Acta Cryst.* **E66**, o374.
- Morimoto, Y., Matsuda, F. & Shirahama, H. (1991). *Synlett*, **3**, 202–203.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2010). E66, o1192-o1193 [doi:10.1107/S1600536810012900]

3-Acetyl-6-chloro-2-methyl-4-phenylquinolinium perchlorate

T. Shahani, H.-K. Fun, S. Sarveswari, V. Vijayakumar and B. P. Reddy

Comment

Quinolines and their derivatives are very important compounds because of their wide occurrence in natural products (Morimoto *et al.*, 1991; Michael, 1997), and biologically active compounds (Markees *et al.*, 1970 ; Campbell *et al.*, 1988). A large variety of quinolines have interesting physiological activities and found attractive applications as pharmaceuticals, agrochemicals and as synthetic building blocks, due to their great importance, the synthesis of new derivatives of quinoline remains an active research area (Katritzky & Arend, 1998; Jiang & Si, 2002).

In the title compound (Fig. 1), the asymmetric unit consists one perchlorate anion and one 3-acetyl-6-chloro-2-methyl-4-phenylquinoline-1-ium cation. The quinolinium ring system (C1/N1/C2–C9) is approximately planar, with a maximum deviation of 0.027 (1) Å at atom C1. The dihedral angle formed between quinolinium ring system and benzene ring (C10–C15) is 78.46 (3)°. Bond lengths (Allen *et al.*, 1987) and angles are normal and comparable to those related structures (Shahani *et al.*, 2010; Fun *et al.*, 2009; Loh *et al.*, 2010).

In the crystal packing (Fig. 2), intermolecular N1—H1N1...O3, C3—H3A...O2, C15—H15A...O5 and C18—H18B...O5 hydrogen bonds (Table 1) link the molecules into three-dimensional network. This crystal structure is further consolidated by C—H... π interactions involving C10–C15 benzene ring (centroid Cg1).

Experimental

A mixture of 3-acetyl-6-chloro-2-methyl-4-phenylquinoline and a catalytic amount of nickel chloride in acid medium was refluxed for about an hour and resultant compound was recrystallized from 3:1 ethanol water to yield colourless blocks of (I).

Refinement

All H atoms were located in a difference map and was refined freely. [N—H = 0.829 (19) Å, C—H = 0.76 (2)–1.025 (17) Å].

Figures




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids.




Fig. 2. The crystal packing of (I), viewed along *a* axis. H atoms not involved in intermolecular interactions (dashed lines) are omitted for clarity.

3-Acetyl-6-chloro-2-methyl-4-phenylquinolinium perchlorate

Crystal data

$C_{18}H_{15}ClNO^+ \cdot ClO_4^-$	$Z = 2$
$M_r = 396.21$	$F(000) = 408$
Triclinic, $P\bar{1}$	$D_x = 1.533 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.3862 (1) \text{ \AA}$	Cell parameters from 9929 reflections
$b = 8.8519 (2) \text{ \AA}$	$\theta = 2.7\text{--}35.1^\circ$
$c = 13.3378 (3) \text{ \AA}$	$\mu = 0.41 \text{ mm}^{-1}$
$\alpha = 92.477 (1)^\circ$	$T = 100 \text{ K}$
$\beta = 91.903 (1)^\circ$	Block, colourless
$\gamma = 99.550 (1)^\circ$	$0.58 \times 0.54 \times 0.27 \text{ mm}$
$V = 858.44 (3) \text{ \AA}^3$	

Data collection

Bruker SMART APEXII CCD diffractometer	7482 independent reflections
Radiation source: fine-focus sealed tube graphite	6933 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.019$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 35.0^\circ$, $\theta_{\text{min}} = 1.5^\circ$
$T_{\text{min}} = 0.797$, $T_{\text{max}} = 0.898$	$h = -11 \rightarrow 11$
27967 measured reflections	$k = -13 \rightarrow 14$
	$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.108$	All H-atom parameters refined
$S = 1.09$	$w = 1/[\sigma^2(F_o^2) + (0.0636P)^2 + 0.2949P]$
7482 reflections	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} < 0.001$

295 parameters

$$\Delta\rho_{\max} = 0.69 \text{ e } \text{\AA}^{-3}$$

0 restraints

$$\Delta\rho_{\min} = -1.00 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.57962 (3)	-0.15711 (3)	0.127788 (19)	0.01984 (6)
O1	-0.34868 (10)	0.25202 (9)	0.18488 (6)	0.02125 (14)
N1	-0.06256 (10)	-0.06546 (8)	0.35547 (5)	0.01256 (12)
C1	-0.15762 (11)	0.04733 (9)	0.34086 (6)	0.01226 (13)
C2	0.08539 (11)	-0.08886 (9)	0.30060 (6)	0.01178 (13)
C3	0.17738 (12)	-0.21217 (10)	0.32169 (6)	0.01442 (14)
C4	0.32724 (13)	-0.23218 (10)	0.26722 (7)	0.01553 (14)
C5	0.38693 (12)	-0.12954 (10)	0.19231 (7)	0.01427 (14)
C6	0.29771 (11)	-0.01015 (9)	0.16974 (6)	0.01300 (13)
C7	0.14195 (11)	0.01160 (9)	0.22419 (6)	0.01113 (12)
C8	0.03934 (11)	0.13153 (9)	0.20510 (6)	0.01083 (12)
C9	-0.10849 (11)	0.14720 (9)	0.26283 (6)	0.01146 (12)
C10	0.09383 (11)	0.23714 (9)	0.12361 (6)	0.01128 (12)
C11	-0.00355 (12)	0.21724 (10)	0.03113 (6)	0.01471 (14)
C12	0.05201 (13)	0.31380 (11)	-0.04601 (7)	0.01646 (15)
C13	0.20163 (13)	0.43138 (10)	-0.03026 (7)	0.01619 (15)
C14	0.29697 (13)	0.45204 (10)	0.06228 (7)	0.01649 (15)
C15	0.24533 (12)	0.35437 (10)	0.13932 (6)	0.01454 (14)
C16	-0.21861 (12)	0.27461 (10)	0.24424 (6)	0.01340 (13)
C17	-0.15358 (17)	0.42459 (12)	0.30030 (8)	0.02276 (18)
C18	-0.31035 (13)	0.06418 (11)	0.40871 (7)	0.01699 (15)
H3A	0.137 (2)	-0.2799 (18)	0.3733 (12)	0.017 (3)*
H4A	0.394 (3)	-0.310 (2)	0.2795 (13)	0.029 (4)*
H6A	0.343 (2)	0.0569 (18)	0.1200 (11)	0.016 (3)*
H11A	-0.117 (2)	0.1334 (19)	0.0189 (12)	0.022 (4)*
H12A	-0.013 (2)	0.2955 (19)	-0.1119 (13)	0.023 (4)*
H13A	0.239 (2)	0.5034 (19)	-0.0772 (12)	0.021 (4)*
H14A	0.394 (2)	0.5305 (19)	0.0737 (12)	0.021 (4)*

supplementary materials

H15A	0.306 (2)	0.3656 (18)	0.2039 (12)	0.020 (4)*
H17A	-0.033 (3)	0.460 (2)	0.2780 (14)	0.031 (4)*
H17B	-0.242 (3)	0.488 (2)	0.2843 (15)	0.037 (5)*
H17C	-0.153 (3)	0.418 (3)	0.3570 (18)	0.045 (6)*
H18A	-0.281 (3)	0.154 (3)	0.4486 (16)	0.042 (5)*
H18B	-0.424 (3)	0.078 (2)	0.3754 (14)	0.031 (4)*
H18C	-0.334 (3)	-0.026 (2)	0.4431 (15)	0.037 (5)*
H1N1	-0.089 (3)	-0.121 (2)	0.4034 (14)	0.030 (4)*
Cl2	0.28102 (3)	0.33444 (2)	0.461521 (15)	0.01541 (5)
O2	0.27428 (10)	0.49972 (8)	0.45103 (6)	0.02070 (14)
O3	0.10191 (10)	0.25407 (9)	0.48844 (6)	0.01977 (14)
O4	0.40805 (10)	0.32400 (9)	0.55505 (5)	0.01960 (13)
O5	0.35420 (12)	0.26779 (11)	0.37509 (6)	0.02648 (16)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.01521 (10)	0.01761 (10)	0.02749 (11)	0.00551 (7)	0.00402 (7)	-0.00306 (8)
O1	0.0153 (3)	0.0210 (3)	0.0279 (4)	0.0044 (2)	-0.0042 (2)	0.0044 (3)
N1	0.0142 (3)	0.0112 (3)	0.0124 (3)	0.0017 (2)	0.0016 (2)	0.0031 (2)
C1	0.0126 (3)	0.0116 (3)	0.0126 (3)	0.0015 (2)	0.0012 (2)	0.0020 (2)
C2	0.0132 (3)	0.0103 (3)	0.0118 (3)	0.0020 (2)	-0.0002 (2)	0.0013 (2)
C3	0.0180 (3)	0.0117 (3)	0.0140 (3)	0.0042 (3)	-0.0018 (3)	0.0020 (2)
C4	0.0176 (4)	0.0132 (3)	0.0166 (3)	0.0058 (3)	-0.0025 (3)	0.0003 (3)
C5	0.0129 (3)	0.0133 (3)	0.0170 (3)	0.0039 (2)	-0.0002 (3)	-0.0016 (3)
C6	0.0128 (3)	0.0118 (3)	0.0146 (3)	0.0025 (2)	0.0011 (2)	0.0005 (2)
C7	0.0116 (3)	0.0100 (3)	0.0119 (3)	0.0021 (2)	0.0001 (2)	0.0011 (2)
C8	0.0113 (3)	0.0095 (3)	0.0116 (3)	0.0012 (2)	0.0002 (2)	0.0018 (2)
C9	0.0116 (3)	0.0108 (3)	0.0122 (3)	0.0020 (2)	0.0011 (2)	0.0024 (2)
C10	0.0122 (3)	0.0104 (3)	0.0116 (3)	0.0023 (2)	0.0016 (2)	0.0025 (2)
C11	0.0155 (3)	0.0150 (3)	0.0133 (3)	0.0014 (3)	-0.0009 (2)	0.0024 (3)
C12	0.0191 (4)	0.0187 (4)	0.0126 (3)	0.0050 (3)	0.0013 (3)	0.0038 (3)
C13	0.0195 (4)	0.0149 (3)	0.0159 (3)	0.0057 (3)	0.0063 (3)	0.0054 (3)
C14	0.0170 (4)	0.0140 (3)	0.0180 (3)	-0.0002 (3)	0.0045 (3)	0.0033 (3)
C15	0.0143 (3)	0.0141 (3)	0.0144 (3)	-0.0002 (3)	0.0010 (2)	0.0022 (2)
C16	0.0132 (3)	0.0136 (3)	0.0146 (3)	0.0042 (2)	0.0033 (2)	0.0042 (2)
C17	0.0308 (5)	0.0161 (4)	0.0229 (4)	0.0101 (3)	-0.0037 (4)	-0.0030 (3)
C18	0.0164 (4)	0.0185 (4)	0.0171 (3)	0.0039 (3)	0.0060 (3)	0.0045 (3)
Cl2	0.01584 (9)	0.01583 (9)	0.01397 (9)	0.00038 (6)	-0.00013 (6)	0.00351 (6)
O2	0.0187 (3)	0.0142 (3)	0.0297 (4)	0.0028 (2)	-0.0018 (3)	0.0094 (2)
O3	0.0148 (3)	0.0220 (3)	0.0209 (3)	-0.0037 (2)	-0.0018 (2)	0.0106 (2)
O4	0.0200 (3)	0.0196 (3)	0.0181 (3)	0.0001 (2)	-0.0077 (2)	0.0070 (2)
O5	0.0264 (4)	0.0343 (4)	0.0192 (3)	0.0083 (3)	0.0016 (3)	-0.0065 (3)

Geometric parameters (\AA , $^\circ$)

Cl1—C5	1.7332 (9)	C10—C15	1.3976 (12)
O1—C16	1.2090 (11)	C11—C12	1.3946 (12)
N1—C1	1.3296 (11)	C11—H11A	1.025 (17)

N1—C2	1.3740 (11)	C12—C13	1.3903 (14)
N1—H1N1	0.829 (19)	C12—H12A	0.981 (17)
C1—C9	1.4123 (11)	C13—C14	1.3903 (13)
C1—C18	1.4919 (12)	C13—H13A	0.929 (16)
C2—C7	1.4097 (11)	C14—C15	1.3936 (12)
C2—C3	1.4115 (12)	C14—H14A	0.917 (17)
C3—C4	1.3754 (13)	C15—H15A	0.952 (16)
C3—H3A	0.954 (16)	C16—C17	1.4937 (14)
C4—C5	1.4117 (13)	C17—H17A	0.956 (19)
C4—H4A	0.926 (19)	C17—H17B	0.95 (2)
C5—C6	1.3738 (12)	C17—H17C	0.76 (2)
C6—C7	1.4159 (11)	C18—H18A	0.93 (2)
C6—H6A	0.942 (15)	C18—H18B	0.963 (19)
C7—C8	1.4295 (11)	C18—H18C	0.93 (2)
C8—C9	1.3788 (11)	Cl2—O5	1.4344 (8)
C8—C10	1.4864 (11)	Cl2—O3	1.4583 (7)
C9—C16	1.5200 (12)	Cl2—O2	1.4846 (7)
C10—C11	1.3965 (11)	Cl2—O4	1.5512 (7)
C1—N1—C2	123.82 (7)	C10—C11—H11A	121.2 (9)
C1—N1—H1N1	118.1 (13)	C13—C12—C11	120.12 (8)
C2—N1—H1N1	117.9 (13)	C13—C12—H12A	120.6 (10)
N1—C1—C9	118.77 (7)	C11—C12—H12A	119.3 (10)
N1—C1—C18	118.45 (7)	C12—C13—C14	119.96 (8)
C9—C1—C18	122.77 (8)	C12—C13—H13A	123.8 (10)
N1—C2—C7	118.94 (7)	C14—C13—H13A	116.1 (10)
N1—C2—C3	119.61 (7)	C13—C14—C15	120.52 (8)
C7—C2—C3	121.45 (8)	C13—C14—H14A	120.5 (10)
C4—C3—C2	118.80 (8)	C15—C14—H14A	119.0 (10)
C4—C3—H3A	120.4 (10)	C14—C15—C10	119.40 (8)
C2—C3—H3A	120.8 (10)	C14—C15—H15A	123.0 (10)
C3—C4—C5	119.88 (8)	C10—C15—H15A	117.5 (10)
C3—C4—H4A	121.9 (11)	O1—C16—C17	123.74 (8)
C5—C4—H4A	118.2 (11)	O1—C16—C9	119.76 (8)
C6—C5—C4	122.16 (8)	C17—C16—C9	116.48 (8)
C6—C5—C11	119.75 (7)	C16—C17—H17A	106.1 (11)
C4—C5—C11	118.09 (7)	C16—C17—H17B	105.6 (12)
C5—C6—C7	118.88 (8)	H17A—C17—H17B	114.6 (16)
C5—C6—H6A	119.5 (10)	C16—C17—H17C	112.6 (17)
C7—C6—H6A	121.6 (10)	H17A—C17—H17C	111 (2)
C2—C7—C6	118.79 (7)	H17B—C17—H17C	107 (2)
C2—C7—C8	118.43 (7)	C1—C18—H18A	110.3 (13)
C6—C7—C8	122.77 (7)	C1—C18—H18B	115.2 (11)
C9—C8—C7	119.31 (7)	H18A—C18—H18B	101.8 (17)
C9—C8—C10	121.17 (7)	C1—C18—H18C	106.8 (13)
C7—C8—C10	119.52 (7)	H18A—C18—H18C	115.6 (17)
C8—C9—C1	120.67 (7)	H18B—C18—H18C	107.4 (16)
C8—C9—C16	120.14 (7)	O5—Cl2—O3	114.23 (5)
C1—C9—C16	119.18 (7)	O5—Cl2—O2	111.92 (5)
C11—C10—C15	120.21 (7)	O3—Cl2—O2	110.06 (5)

supplementary materials

C11—C10—C8	120.13 (7)	O5—C12—O4	109.31 (5)
C15—C10—C8	119.65 (7)	O3—C12—O4	104.17 (4)
C12—C11—C10	119.78 (8)	O2—C12—O4	106.60 (4)
C12—C11—H11A	119.0 (9)		
C2—N1—C1—C9	1.85 (12)	C7—C8—C9—C16	179.58 (7)
C2—N1—C1—C18	-177.33 (8)	C10—C8—C9—C16	-0.34 (12)
C1—N1—C2—C7	0.14 (12)	N1—C1—C9—C8	-2.26 (12)
C1—N1—C2—C3	-179.90 (8)	C18—C1—C9—C8	176.88 (8)
N1—C2—C3—C4	-178.83 (8)	N1—C1—C9—C16	178.84 (7)
C7—C2—C3—C4	1.12 (13)	C18—C1—C9—C16	-2.01 (12)
C2—C3—C4—C5	0.44 (13)	C9—C8—C10—C11	-78.54 (10)
C3—C4—C5—C6	-1.37 (13)	C7—C8—C10—C11	101.54 (10)
C3—C4—C5—C11	178.54 (7)	C9—C8—C10—C15	102.80 (10)
C4—C5—C6—C7	0.70 (13)	C7—C8—C10—C15	-77.12 (10)
C11—C5—C6—C7	-179.20 (6)	C15—C10—C11—C12	0.67 (13)
N1—C2—C7—C6	178.19 (7)	C8—C10—C11—C12	-177.97 (8)
C3—C2—C7—C6	-1.77 (12)	C10—C11—C12—C13	-1.27 (14)
N1—C2—C7—C8	-1.72 (11)	C11—C12—C13—C14	0.51 (14)
C3—C2—C7—C8	178.32 (7)	C12—C13—C14—C15	0.86 (14)
C5—C6—C7—C2	0.84 (12)	C13—C14—C15—C10	-1.44 (14)
C5—C6—C7—C8	-179.25 (8)	C11—C10—C15—C14	0.67 (13)
C2—C7—C8—C9	1.28 (11)	C8—C10—C15—C14	179.32 (8)
C6—C7—C8—C9	-178.63 (7)	C8—C9—C16—O1	89.51 (11)
C2—C7—C8—C10	-178.80 (7)	C1—C9—C16—O1	-91.59 (10)
C6—C7—C8—C10	1.30 (12)	C8—C9—C16—C17	-88.92 (10)
C7—C8—C9—C1	0.70 (12)	C1—C9—C16—C17	89.98 (10)
C10—C8—C9—C1	-179.22 (7)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N1...O3 ⁱ	0.832 (18)	1.896 (18)	2.7177 (10)	169 (2)
C3—H3A...O2 ⁱⁱ	0.955 (16)	2.583 (16)	3.3010 (11)	132.2 (12)
C15—H15A...O5	0.951 (16)	2.512 (16)	3.3716 (12)	150.4 (13)
C18—H18B...O5 ⁱⁱⁱ	0.97 (2)	2.53 (2)	3.3266 (13)	139.5 (14)
C12—H12A...Cg1 ^{iv}	0.981 (17)	2.694 (17)	3.5810 (10)	150.6 (13)

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $x, y-1, z$; (iii) $x-1, y, z$; (iv) $-x, -y, -z$.

Fig. 1





Fig. 2



Copyright of Acta Crystallographica: Section E (International Union of Crystallography - IUCr) is the property of International Union of Crystallography - IUCr and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.