



http://www.e-journals.net



ISSN: 0973-4945; CODEN ECJHAO

E-Journal of Chemistry

2010, 7(1), 281-286

A Facile Microwave Assisted One-pot Strategy for the Synthesis of Bis-hexahydroquinazolin-5(6H)-ones

MILAN CH. DUTTA[§], KAUSHIK CHANDA[§],
R. L. NONGKHLAW and JAI N. VISHWAKARMA^{§*}

[§]Organic Research Lab., Department of Chemistry,
St. Anthony's College, Shillong-793 001, India.

Department of Chemistry,
North-Eastern Hill University, Shillong-793 022, India.
jnvishwakarma@rediffmail.com

Received 15 June 2009; Accepted 18 July 2009


Abstract: A facile microwave assisted one-pot synthetic protocol has been devised for the synthesis of 3,3'-(alkanediyl) bis (1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-ones (**2a-c**), 3,3'-(1,4-phenylene) bis (1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (**2d**), 3,3'-(alkanediyl) bis (7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-ones (**2e-g**) and 3,3'-(1,4-phenylene) bis(7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (**2h**) by the cyclocondensation of cyclic enaminones **1a, b** with diamines and formaldehyde.

Keywords: Enaminones, Quinazolinones, Diamines, Cyclocondensation.

Introduction

Fused pyrimidines are found in a broad variety of natural products¹, used in medicines², possess antimalarial activities³ and other important biological properties⁴. Recently, 1,2,3,4,5,6,7,8-octahydroquinazolin-2,5-dione derivatives have been reported to exhibit potent calcium antagonist activities^{5,6} and have also attracted considerable attention owing to their potential antibacterial activity against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*^{7,8}. However, synthesis of 5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines⁹ is least attended to and to the best of our knowledge bis(5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines) are unknown in the literature and hence their biological properties remain unexplored. Prompted by the promising biological properties of 1,2,3,4,5,6,7,8-octahydroquinazolines, we have recently reported¹⁰ a facile synthetic methodology for 1-aralkyl/aryl-3-alkyl/aralkyl/aryl-5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines and 1-aralkyl/aryl-3-alkyl/aralkyl/aryl-7,7-dimethyl-5-oxo-1,2,3,4,5,6,7,8-octahydroquinazolines. In continuation with our efforts on the synthesis of tetrahydropyrimidines¹¹⁻¹⁴, we now report herein a facile

one pot synthetic strategy for bis-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-ones in which the two quinazoline rings are linked through flexible aliphatic chains or through rigid aromatic ring (Scheme 1).



Scheme 1

Experimental

Melting points were recorded by open capillary method and are uncorrected. The IR spectra were recorded on a Perkin-Elmer 983 spectrometer. ¹H NMR (90 MHz) spectra were recorded on Varian EM-390 spectrometer. High resolution ¹H NMR and ¹³C NMR (300 MHz) spectra were recorded on Bruker ACF-300 spectrometer. The chemical shifts (δ ppm) and the coupling constants (Hz) are reported in the standard fashion with reference to TMS as internal reference. FAB-mass spectra (MS) were measured on JEOL 3SX 102/DA-6000 Mass spectrometer using Argon as the FAB gas and *m*-nitrobenzylalcohol as the matrix. Elemental analyses were performed on a Vario-EL III instrument. Microwave irradiation was carried out in a domestic MW oven (Samsung CE2733G) operating at 2450 MHz. Enaminones **1a** and **1b** were synthesized by our reported procedure¹⁵.

General procedure

A mixture of diamine (0.5 mmol) and formaldehyde (2 mmol, 40% solution) in 1.5 mL methanol was shaken at room temperature for 5 minutes. To this a solution of enaminones **1** (1 mmol) was added in 5-6 mL methanol and the resulting mixture was irradiated in a domestic microwave oven for specified period of time (Table 2). After the completion of the reaction (monitored by TLC), methanol was removed under reduced pressure to give a gum, which on trituration with hexane and subsequent recrystallization in appropriate solvent gave compounds **2a**, **2e**, **2g** and **2h**. In case of compounds **2b**, **2c**, **2d** and **2f** the gum was chromatographed using neutral alumina and ethylacetate (eluant). Under thermal conditions these reactions took much longer time for completion giving the desired products in comparatively lower yields.

3,3'-(Ethane-1,2-diyl)bis(1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (**2a**))

It was obtained as a pale yellow solid in 85% yield; mp 182-184 °C (EtOAc); IR (KBr): 1493, 1566 cm⁻¹; ¹H NMR (CDCl₃): δ 1.70 (m, 4H, 2 CH₂), 2.12-2.28 (t, 4H, 2 CH₂), 2.31-2.60 (t, 4H, 2 CH₂), 2.83 (s, 4H, 2 CH₂), 3.73 (s, 4H, 2 CH₂), 4.43 (s, 4H, 2 CH₂), 7.07-7.87 (m, 10H); MS: m/z 483 (MH⁺). Anal. Calc. for C₃₀H₃₄N₄O₂ (482.27): C, 74.66; H, 7.10; N, 11.61. Found: C, 74.41; H, 7.07; N, 11.66%.

3,3'-(Propane-1,3-diyl)bis(1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (**2b**))

It was obtained as yellow gum in 76% yield; IR (CCl₄): 1493, 1560 cm⁻¹; ¹H NMR (CDCl₃): δ 1.84-1.85 (m, 4H, 2 CH₂), 1.86-1.88 (m, 2H), 2.18-2.20 (t, 4H, 2 CH₂), 2.32-2.36 (t, 4H, 2 CH₂), 2.67-2.72 (t, 4H, 2 CH₂), 3.64 (s, 4H, 2 CH₂), 4.30 (s, 4H, 2 CH₂), 7.11-7.13 (m, 4H), 7.27-7.43 (m, 6H); ¹³C NMR (CDCl₃): δ 22.10, 26.07, 27.72, 36.55, 47.96, 50.95, 71.98, 104.83, 127.36, 127.50, 129.84, 143.02, 158.36, 194.94; MS: m/z, 497 (MH⁺). Anal. Calc. for C₃₁H₃₆N₄O₂ (496.28): C, 74.97; H, 7.31; N, 11.28. Found: C, 74.66; H, 7.33; N, 11.24%.

3,3'-(Butane-1,4-diyl)bis(1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2c)

It was obtained as yellow gum in 77% yield; IR (CCl₄): 1493, 1560 cm⁻¹; ¹H NMR (CDCl₃): δ 1.58 (m, 4H, 2 CH₂), 1.85-1.88 (m, 4H, 2 CH₂), 2.21 (t, 4H, 2 CH₂), 2.33 (t, 4H, 2 CH₂), 2.62 (t, 4H, 2 CH₂), 3.64 (s, 4H, 2 CH₂), 4.29 (s, 4H, 2 CH₂), 7.12-7.14 (m, 4H), 7.30-7.43 (m, 6H); ¹³C NMR (CDCl₃): δ 22.08, 25.59, 27.71, 30.51, 47.59, 52.59, 72.19, 104.76, 124.61, 127.37, 129.31, 143.05, 158.48, 195.03; MS: m/z, 511 (MH⁺). Anal. Calc. for C₃₂H₃₈N₄O₂ (510.29): C, 75.26; H, 7.50; N, 10.97. Found: C, 75.50; H, 7.48; N, 10.92%.

3,3'-(1,4-Phenylene)bis(1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2d)

It was obtained as yellow solid in 79% yield; mp 191-193 °C; IR (KBr): 1427, 1507, 1566 cm⁻¹; ¹H NMR (CDCl₃): δ 1.70-2.06 (m, 4H, 2 CH₂), 2.10-2.30 (t, 4H, 2 CH₂), 2.33-2.60 (t, 4H, 2 CH₂), 4.56 (s, 4H, 2 CH₂), 4.99 (s, 4H, 2 CH₂), 6.96-7.93 (m, 14H); MS: m/z, 531 (MH⁺). Anal. Calc. for C₃₄H₃₄N₄O₂ (530.27): C, 76.95; H, 6.46; N, 10.56. Found: C, 76.70; H, 6.45; N, 10.51%.

3,3'-(Ethane-1,2-diyl)bis(7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2e)

It was obtained as yellow solid in 87% yield; mp 193-194 °C (CHCl₃/Hexane); IR (KBr): 1493, 1566, 1619 cm⁻¹; ¹H NMR (CDCl₃): δ 1.03 (s, 12H, 4 CH₃), 2.10 (s, 4H, 2 CH₂), 2.23 (s, 4H, 2 CH₂), 2.93 (s, 4H, 2 CH₂), 3.80 (s, 4H, 2 CH₂), 4.50 (s, 4H, 2 CH₂), 7.03-7.76 (m, 10H); MS: m/z, 539 (MH⁺). Anal. Calc. for C₃₄H₄₂N₄O₂ (538.33): C, 75.80; H, 7.86; N, 10.40. Found: C, 76.05; H, 7.88; N, 10.44%.

3,3'-(Propane-1,3-diyl)bis(7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2f)

It was obtained as yellow gum in 75% yield; IR (CCl₄): 1427, 1493, 1566, cm⁻¹; ¹H NMR (CDCl₃): δ 0.90 (s, 12H, 4 CH₃), 1.59-1.61 (m, 2H), 1.99 (s, 4H, 2 CH₂), 2.13 (s, 4H, 2 CH₂), 2.60-2.65 (m, 4H, 2 CH₂), 3.58 (s, 4H, 2 CH₂), 4.24 (s, 4H, 2 CH₂), 7.01-7.08 (m, 4H), 7.25-7.41 (m, 6H); ¹³C NMR (CDCl₃): δ 24.92, 27.28, 27.36, 31.64, 39.99, 46.40, 48.65, 48.96, 49.46, 70.92, 102.15, 126.38, 127.89, 129.39, 141.78, 155.62, 193.42. Anal. Calc. for C₃₅H₄₄N₄O₂ (552.35): C, 76.05; H, 8.02; N, 10.15. Found: C, 76.31; H, 7.98; N, 10.11%.

3,3'-(Butane-1,4-diyl)bis(7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2g)


It was obtained as a pale yellow solid in 90% yield; mp 168-169 °C (EtOAc); IR (KBr): 1440, 1566, 1613 cm⁻¹; ¹H NMR (CDCl₃): δ 1.03 (s, 12H, 4 CH₃), 1.46-1.83 (m, 4H, 2 CH₂), 2.13 (s, 4H, 2 CH₂), 2.26 (s, 4H, 2 CH₂), 2.56-2.90 (m, 4H, 2 CH₂), 3.76 (s, 4H, 2 CH₂), 4.45 (s, 4H, 2 CH₂), 7.10-7.86 (m, 10H); MS: m/z, 567 (MH⁺). Anal. Calc. for C₃₆H₄₆N₄O₂ (566.36): C, 76.29; H, 8.18; N, 9.89. Found: C, 76.02; H, 8.21; N, 9.83%.

3,3'-(1,4-Phenylene)bis(7,7-dimethyl-1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (2h)

It was obtained as yellow solid in 89% yield; mp 126-128 °C (MeOH/EtOAc); IR (KBr): 1507, 1566, 1610 cm⁻¹; ¹H NMR (CDCl₃): δ 0.93 (s, 12H, 4 CH₃), 2.00 (s, 4H, 2 CH₂), 2.23 (s, 4H, 2 CH₂), 4.27 (s, 4H, 2 CH₂), 4.89 (s, 4H, 2 CH₂), 6.91-6.96 (m, 8H), 7.27-7.36 (m, 6H); ¹³C NMR (CDCl₃): δ 28.08, 28.52, 32.82, 41.24, 45.96, 50.21, 52.59, 71.21, 104.63, 119.13, 127.53, 127.62, 129.88, 142.84, 157.52, 194.22; MS: m/z, 587 (MH⁺). Anal. Calc. for C₃₈H₄₂N₄O₂ (586.33): C, 77.78; H, 7.21; N, 9.55. Found: C, 77.52; H, 7.18; N, 9.61%.

Results and Discussion

Thus, when a mixture of 3-anilino-cyclohex-2-en-1-one (**1a**), ethylenediamine and formaldehyde (2:1:4) in methanol was subjected to MWI, work-up of the reaction mixture followed by chromatographic purification yielded a solid in 85% yields, which was characterized as 3,3'-(ethane-1,2-diyl)bis(1-phenyl-1,2,3,4,7,8-hexahydroquinazolin-5(6H)-one (**2a**). The reaction was found to be general with other diamines and with corresponding **1a-b** to give the respective **2b-h** in 77-90% overall yields (Table 1). These reactions were also carried out under thermal conditions to give the expected products in comparatively lower yields and in addition they took much longer time for completion (Table 2). The structures of the Bis-quinazolines were assigned on the basis of spectral and analytical data. Thus, the infrared spectra of **2a-h** showed strong peaks in the range of 1427 to 1619 cm^{-1} due to highly delocalised double bonds and carbonyl group stretching frequencies of enaminone functionalities. In the ^1H NMR spectra of **2a** & **2e** the NCH_2 protons of ethylene chain appeared as singlets at 2.83 ppm and 2.93 ppm respectively whereas the NCH_2 protons of propylene chain in **2b** & **2f** gave triplets in the range of 2.60-2.72 ppm. The protons at C_2 of propylene chain gave multiplets in the range of 1.59-1.88 ppm. Likewise in **2c** & **2g** the protons at C_1 & C_2 of butylene chains gave multiplets in the range of 2.56-2.90 and 1.46-1.88 ppm respectively. The protons at C_2 and C_4 of quinazoline ring resonated in the range of 4.24-4.99 and 3.64-4.56 ppm respectively. In **2a-d** the C_8 protons of the parent ring appeared as triplets between 2.10 & 2.30 ppm but in **2e-h** they appeared as singlets between 1.99 & 2.10 ppm. The C_6 protons of the parent ring in **2a-d** appeared as triplets between 2.31 & 2.60 ppm but in **2e-h** they appeared as singlets between 2.13 & 2.26 ppm. The C_7 protons of the ring in **2a-d** resonated giving multiplets in the range of 1.58-2.06 ppm. The two-methyl protons at C_7 in **2e-h** gave singlets between 0.90 & 1.03 ppm. The aromatic protons resonated in their usual range of 6.91-7.93 ppm. A plausible mechanism for the formation of **2** from the cyclic enaminones **1** is worked out (Scheme 2).



Scheme 2.

Table 1. Synthesis of bis-quinazolinones (**2a-h**)











Entry	Enaminones (1)	Conditions	Bis-quinazolinones (2)
1	 1a	H ₂ N-(CH ₂) ₂ -NH ₂ , CH ₂ O, MeOH/180 W 3 minutes	 2a
2	1a	H ₂ N-(CH ₂) ₃ -NH ₂ , CH ₂ O, MeOH/180 W 2 minutes	 2b
3	1a	H ₂ N-(CH ₂) ₄ -NH ₂ , CH ₂ O, MeOH/180 W 2.5 minutes	 2c
4	1a	H ₂ N-C ₆ H ₄ -NH ₂ , CH ₂ O, MeOH/180 W, 3 minutes	 2d
5	 1b	H ₂ N-(CH ₂) ₂ -NH ₂ , CH ₂ O, MeOH/180 W, 2 minutes	 2e
6	1b	H ₂ N-(CH ₂) ₃ -NH ₂ , CH ₂ O, MeOH/180 W, 4 minutes	 2f
7	1b	H ₂ N-(CH ₂) ₄ -NH ₂ , CH ₂ O, MeOH/180 W, 1.5 minutes	 2g
8	1b	H ₂ N-C ₆ H ₄ -NH ₂ , CH ₂ O, MeOH/180 W, 2 minutes	 2h

Table 2. Comparison of microwave irradiation and thermal condition results.




Compound	Microwave Irradiation (180 W)		Thermal (Reflux)	
	Time, min	Yield, %	Time, h	Yield, %
2a	3	85	27	58
2b	2	76	22	70
2c	2.5	77	24	65
2d	3	79	24	71
2e	2	87	22	88
2f	4	75	23	68
2g	1.5	90	8	59
2h	2	89	20	55

Acknowledgements

The authors wish to thank the Principal, Rev. Fr. Ioannis Warpakma, SDB for the facilities and Rev. Fr. Stephen Mavely, SDB and Rev. Fr. Joseph Nellanatt, SDB for their encouragement during the course of this investigation. The financial support from ICAR-NATP-PIU and UGC-New Delhi is gratefully acknowledged. Authors (KC & MCD) thank ICAR for Senior Research Fellowships. Thanks are also due to the Heads of RSIC-CDRI (Lucknow) and RSIC-NEHU (Shillong) for recording spectra.

References

1. Taylor E C and Patel M, *J Heterocyclic Chem.*, 1991, **28**, 1857.
2. Coates W J, *Comprehensive Heterocyclic Chemistry II*, Vol 6, Edited by Boulton A J, Pergamon, Oxford, 1996, 225-231.
3. Elslager E F, Hess C, Johnson J, Ortwine D, Chu V and Werbel L M, *J Med Chem.*, 1981, **24**, 127.
4. Rivero I A, Espinoza K and Somanathan R, *Molecules*, 2004, **9**, 609 and references cited therein.
5. Sarac S, Yarim M, Ertan M, Kilic F S and Erol K, *Arzneim Forsch Drug Res.*, 2002, **52**, 27.
6. Yarim M, Sarac S, Kilic F S and Erol K, *Il Farmaco*, 2003, **58**, 17.
7. Kidwai M, Saxena S, Khan M K R and Thukral S S, *Eur J Med Chem.*, 2005, **40**, 816.
8. Kantevari M, Bantu R and Nagarapu L, *ARKIVOC*, 2006, **16**, 136.
9. Hamama W S, Hammouda M and Afsah E M, *Z. Naturforsch.*, 1988, **43B**, 483; *Chem. Abstr.*, 1988, **109**, 170359k.
10. Chanda K, Dutta M C and Vishwakarma J N, *Indian J Chem.*, 2006, **45B**, 1076.
11. Vishwakarma J N, Mofizuddin M, Ila H and Junjappa H, *J Heterocyclic Chem.*, 1988, **25**, 1387.
12. Karim E, Kishore K and Vishwakarma J N, *J Heterocyclic Chem.*, 2003, **40**, 901.
13. Chanda K, Dutta M C, Karim E and Vishwakarma J N, *J Heterocyclic Chem.*, 2004, **41**, 627.
14. Dutta M C, Chanda K and Vishwakarma J N, *J Heterocyclic Chem.*, 2005, **42**, 121.
15. Chanda K, Dutta M C and Vishwakarma J N, *Indian J Chem.*, 2004, **43B**, 2475.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

