

RESEARCH

Open Access



A fitted numerical scheme for second order singularly perturbed delay differential equations via cubic spline in compression

Podila Pramod Chakravarthy^{1*}, S Dinesh Kumar¹, Ragi Nageshwar Rao² and Devendra P Ghate³

*Correspondence:

pramodpodila@yahoo.co.in

¹Department of Mathematics,
Visvesvaraya National Institute of
Technology, Nagpur, 440010, India
Full list of author information is
available at the end of the article

Abstract

This paper deals with the singularly perturbed boundary value problem for the second order delay differential equation. Similar boundary value problems are associated with expected first-exit times of the membrane potential in models of neurons. An exponentially fitted difference scheme on a uniform mesh is accomplished by the method based on cubic spline in compression. The difference scheme is shown to converge to the continuous solution uniformly with respect to the perturbation parameter, which is illustrated with numerical results.

Keywords: singular perturbations; cubic spline in compression; boundary layer; delay differential equation; exponentially fitted finite difference method

1 Background

In the last few decades there has been a growing interest in the study of delay differential equations due to their occurrence in a wide variety of application fields such as bio-sciences, control theory, economics, material science, medicine, robotics *etc.* Any system involving a feedback control will almost always involve time delays. These arise because a finite time is required to sense the information and then to react to it. The delays or lags can represent gestation times, incubation periods, transport delays *etc.* Delay models are also prominent in describing several aspects of infectious disease dynamics such as primary infection, drug therapy, immune response *etc.* Delays have also appeared in the study of chemostat models, circadian rhythms, epidemiology, the respiratory system, tumor growth and neural networks. Statistical analysis of ecological data has shown that there is evidence of delay effects in the population dynamics of many species.

The details of the theory and applications of differential difference equations can be found in the collection of books, to name a few, Bellman and Cooke [1], Driver [2], El'sgol'ts and Norkin [3], Erneux [4], Gopalsamy [5], Györi and Ladas [6], Halanay [7], Kuang [8] and Smith [9]. In recent years there has been a growing interest in the numerical study of differential difference equations. However, the first discrete solution to delay differential equations was given by Feldstein [10], which became a landmark work to most of the researchers working in numerical analysis of delay differential equations. Bellen and Zennaro [11] gave the theoretical aspects of numerical methods for ordinary and delay differential equations, and suitable techniques for solving numerically such type of equations.

A singularly perturbed delay differential equation is a differential equation in which the highest derivative is multiplied by a small parameter and which involves at least one shift term. Such problems arise frequently in the mathematical modeling of various physical and biological phenomena like optically bistable devices [12, 13], description of the human pupil reflex [14], a variety of models for physiological processes or diseases [15], variational problems in control theory [16, 17] and the first-exit time problem in the modeling of the activation of neuronal variability [18].

Singularly perturbed delay differential equations have up to now not been satisfactorily discussed in numerical analysis literature; however, in recent years there has been a growing interest in the numerical study of such problems. Most of the previous works have been centered on the existence and uniqueness of solutions for initial value problems in differential difference equations and very little attention has been paid to construction of approximate solutions. The computation of the solution of delay differential equations has been a great challenge and great importance due to the appearance of such equations in mathematical modeling of biological problems. Stein [18] approximated the solution of his model of the activation of neuronal variability, which was studied by Tuckwell [19–21] and by Wilbur and Rinzel [22]. Lange and Miura [23–28] gave a series of papers on singularly perturbed differential difference equations by extending the matched asymptotic expansion approach developed for ordinary differential equations to obtain the approximate solution of these differential difference equations. An extensive numerical work has been initiated by Kadalbajoo and Sharma in their papers [29–38], Kadalbajoo and Kumar [39], Kadalbajoo and Ramesh [40]. Gulsu and Sezer [41] proposed a Taylor polynomial approach for solving m th order linear differential difference equations with mixed conditions. This method is based on first taking the truncated Taylor's expansions of the functions in the differential difference equations and then substituting their matrix forms into the equation. Hence the resultant matrix equation can be solved and the unknown Taylor coefficients can be found approximately.

It is well known that standard discretization methods for solving singular perturbation problems are unstable and fail to give accurate results when the perturbation parameter ε is small. Therefore it is important to develop suitable numerical methods to deal with these problems whose accuracy does not depend on the parameter value ε . So the method should be uniformly convergent with respect to the perturbation parameter, and various approaches for the numerical methods to solve singularly perturbed differential equations are given in [42–45]. The use of cubic splines for the solution of linear two point boundary value problems was suggested by Bickley [46]. Aziz and Khan [47] proposed a method based on cubic spline in compression for the linear second order singularly perturbed problems which have second and fourth order convergence depending on the choice of the parameters λ_1 and λ_2 involved in the method.

The analytical and numerical solution of singularly perturbed delay differential equations with large delays can be found in Amiraliyev and Erdogan [48], Amiraliyev and Cimen [49], Amiraliyeva *et al.* [50], Erdogan and Amiraliyev [51]. Subburayan and Ramanujam [52] gave an initial value technique to solve the singularly perturbed boundary value problem for the second order ordinary differential equations of convection-diffusion type with delay. Ghomanjani *et al.* [53] presented the Bezier curves to solve the optimal control problem with pantograph delays. A direct algorithm for solving this problem was given. Ghomanjani *et al.* [54] applied, for the first time, Bernstein's approximation on delay

differential equations and delay systems with inverse delay that models these problems. A direct algorithm is given for solving this problem. The delay function and inverse time function are expanded by the Bezier curves. The Bezier curves are chosen as piecewise polynomials of degree n , and the Bezier curves are determined on any subinterval by $n + 1$ control points. The approximated solution of delay systems containing inverse time is derived.

In this paper, we propose a scheme based on cubic spline in compression which comprises an exponentially fitted difference scheme on a uniform mesh. In Section 2, we state some important properties of the exact solution. In Section 3, we describe a difference scheme based on cubic spline in compression for a second order singularly perturbed delay differential equation. In Section 4, we give the numerical algorithm to solve a singularly perturbed delay differential equation. Some numerical results are presented in Section 5, and conclusions are given in Section 6.

2 Statement of the problem

We consider the following boundary value problem (BVP) for the delay differential equation (DDE):

$$\varepsilon y''(x) + a(x)y'(x) + b(x)y(x - 1) = f(x), \quad 0 < x < 2, \tag{1}$$

subject to the interval and boundary conditions

$$\begin{aligned} y(x) &= \phi(x), \quad x \in [-1, 0]; \\ y(2) &= \beta, \end{aligned} \tag{2}$$

where $0 < \varepsilon \ll 1$ and $a(x) \geq \alpha > 0$, $a(x), b(x), f(x)$ are given sufficiently smooth functions on $[0, 2]$, $\phi(x)$ is a smooth function on $[-1, 0]$ and β is a given constant which is independent of ε , the boundary value problem (1) along with (2) exhibits a strong boundary layer at $x = 0$ (cf. [49], p.2351).

If $a(x) < 0$, $a(x), b(x), f(x)$ are given sufficiently smooth functions on $[0, 2]$, $\phi(x)$ is a smooth function on $[-1, 0]$ and β is a given constant which is independent of ε , then the boundary value problem (1) along with (2) exhibits a strong boundary layer at $x = 2$ (cf. [52], p.236).

2.1 Stability result

Here we show some properties of the solution of (1) and (2). We use the following convention:

$$\begin{aligned} \|g\|_\infty &= \max_{0 \leq x \leq 2} |g(x)|, & \|g\|_1 &= \int_0^2 |g(x)| \, dx, & \|g\|_{\infty,1} &= \int_0^1 |g(x)| \, dx, \\ \|g\|_{\infty,2} &= \int_1^2 |g(x)| \, dx & \text{and} & \|g\|_0 &= \int_{-1}^0 |g(x)| \, dx. \end{aligned}$$

Lemma *If $a(x), b(x), f(x) \in C[0, 2]$ and $\phi(x) \in C[-1, 0]$ and $\rho = \alpha^{-1} \|b\|_{\infty,2} < 1$, then the solution $y(x)$ of problem (1) and (2) follows the estimates*

$$\|y\|_\infty \leq C_0, \tag{3}$$

$$|y'(x)| \leq C_2 \left(1 + \frac{1}{\varepsilon} e^{-\frac{\alpha x}{\varepsilon}} \right), \quad 0 \leq x \leq 2, \tag{4}$$

where

$$C_0 = (|\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0)(1 - \rho)^{-1},$$

$$C_2 = \frac{(|\beta| + |\phi(0)| + C_1)}{c_0},$$

$$C_1 = \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0 + \alpha^{-1} \|b\|_{\infty,2} C_0,$$

$$c_0 = \frac{1}{a^*} \left(1 - e^{-\frac{2a^*}{\varepsilon}} \right),$$

and $a^* = \|a\|_{\infty}$.

Proof From (1) we have

$$y'(x) = y'(0) e^{-\frac{1}{\varepsilon} \int_0^x a(\eta) d\eta} - \frac{1}{\varepsilon} \int_0^x F(\xi) e^{-\frac{1}{\varepsilon} \int_{\xi}^x a(\eta) d\eta} d\xi \tag{5}$$

with $F(x) = -f(x) + b(x)y(x-1)$.

Integrating (5) over $(0, x)$ we get

$$y(x) - y(0) = y'(0) \int_0^x e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau - \frac{1}{\varepsilon} \int_0^x d\tau \int_0^{\tau} F(\xi) e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\xi.$$

Since $y(0) = \phi(0)$, we have

$$y(x) = \phi(0) + y'(0) \int_0^x e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau - \frac{1}{\varepsilon} \int_0^x F(\xi) \left(\int_{\xi}^x e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau \right) d\xi. \tag{6}$$

Using the condition $y(2) = \beta$, we have

$$y'(0) = \frac{\beta - \phi(0) + \frac{1}{\varepsilon} \int_0^2 F(\xi) d\xi \int_{\xi}^2 e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau}. \tag{7}$$

Substituting (7) in (6), we get

$$y(x) = \phi(0) + \left[\beta - \phi(0) + \frac{1}{\varepsilon} \int_0^2 F(\xi) d\xi \int_{\xi}^2 e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau \right] \times \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau} - \frac{1}{\varepsilon} \int_0^x \left(F(\xi) \int_{\xi}^x e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau \right) d\xi. \tag{8}$$

Using Green's function,

$$G(x, \xi) = \frac{1}{\varepsilon} \int_{\xi}^2 e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau \cdot \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^s a(\eta) d\eta} ds}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau} - \frac{1}{\varepsilon} T_0(x - \xi) \int_{\xi}^x e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau, \tag{9}$$

equation (8) can be rewritten as

$$y(x) = \left(1 - \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^s a(\eta) d\eta} ds}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^\tau a(\eta) d\eta} d\tau} \right) \phi(0) + \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^s a(\eta) d\eta} ds}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^\tau a(\eta) d\eta} d\tau} \beta + \int_0^2 G(x, \xi) F(\xi) d\xi, \tag{10}$$

where $T_0(\lambda) = 1, \lambda \geq 0; T_0(\lambda) = 0, \lambda < 0$.

Alternatively the Green's function of the operator

$$Ly = -\varepsilon y''(x) - a(x)y'(x), \quad 0 < x < 2, \quad y(0) = 0 \quad \text{and} \quad y(2) = 0$$

can be expressed as

$$G(x, \xi) = \frac{1}{\varepsilon w(\xi)} \begin{cases} \varphi_1(\xi)\varphi_2(x), & 0 \leq \xi \leq x \leq 2, \\ \varphi_1(x)\varphi_2(\xi), & 0 \leq x \leq \xi \leq 2, \end{cases} \tag{11}$$

where the functions $\varphi_1(x)$ and $\varphi_2(x)$ are solutions of the problems

$$\begin{aligned} L\varphi_1 &= 0, & \varphi_1(0) &= 0, & \varphi_1(2) &= 1, \\ L\varphi_2 &= 0, & \varphi_2(0) &= 1, & \varphi_2(2) &= 0 \end{aligned}$$

and $w(\xi) = \frac{\varphi(\xi)}{Q(2)}$,

$$Q(x) = \int_0^x \varphi(s) ds, \quad \varphi(\xi) = \exp\left[\frac{-1}{\varepsilon} \int_0^\xi a(\tau) d\tau\right].$$

Formula (11) means that $G(x, \xi) \geq 0$, and it follows from (9) that

$$\begin{aligned} \max_{x, \xi \in [0, 2]} G(x, \xi) &= \max_{x, \xi \in [0, 2]} \left(\frac{1}{\varepsilon} \int_\xi^2 e^{-\frac{1}{\varepsilon} \int_\xi^\tau a(\eta) d\eta} d\tau \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^s a(\eta) d\eta} ds}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^\tau a(\eta) d\eta} d\tau} \right. \\ &\quad \left. - \frac{1}{\varepsilon} T_0(x - \xi) \int_\xi^x e^{-\frac{1}{\varepsilon} \int_\xi^\tau a(\eta) d\eta} d\tau \right) \\ &\leq \max_{x, \xi \in [0, 2]} \left(\frac{1}{\varepsilon} \int_\xi^2 e^{-\frac{1}{\varepsilon} \int_\xi^\tau a(\eta) d\eta} d\tau \cdot \frac{\int_0^x e^{-\frac{1}{\varepsilon} \int_0^s a(\eta) d\eta} ds}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^\tau a(\eta) d\eta} d\tau} \right) \\ &\leq \max_{x, \xi \in [0, 2]} \left(\frac{1}{\varepsilon} \{-\alpha^{-1} \varepsilon [e^{-\frac{\alpha}{\varepsilon}(2-\xi)} - 1]\} \right) \\ &\leq \max_{x, \xi \in [0, 2]} \left(\frac{1}{\varepsilon} \{\alpha^{-1} \varepsilon [1 - e^{-\frac{\alpha}{\varepsilon}(2-\xi)}]\} \right) \\ &\leq \alpha^{-1}. \end{aligned}$$

Hence $G(x, \xi) \leq \alpha^{-1}$. Using this inequality in (10), we obtain

$$\begin{aligned} \|y(x)\|_\infty &\leq |\phi(0)| + |\beta| + \int_0^2 |G(x, \xi)| |F(\xi)| d\xi \\ &\leq |\phi(0)| + |\beta| + \max_{x, \xi \in [0, 2]} |G(x, \xi)| \int_0^2 |F(\xi)| d\xi \end{aligned}$$

$$\begin{aligned} &\leq |\phi(0)| + |\beta| + \alpha^{-1} \int_0^2 |F(\xi)| d\xi \\ &\leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \int_0^2 |b(\xi)y(\xi - 1)| d\xi. \end{aligned}$$

Replacing $\xi = 1 + s$, we find that

$$\begin{aligned} &\leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \int_{-1}^1 |b(1 + s)y(s)| ds \\ &\leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \int_{-1}^0 |b(1 + s)\phi(s)| ds + \alpha^{-1} \int_0^1 |b(1 + s)y(s)| ds \\ &\leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \int_{-1}^0 |b(1 + s)\phi(s)| ds + \alpha^{-1} \|y\|_\infty \int_1^2 |b(s)| ds \\ &\leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0 + \alpha^{-1} \|b\|_{\infty,2} \|y\|_\infty. \end{aligned}$$

Therefore

$$(1 - \alpha^{-1} \|b\|_{\infty,2}) \|y\|_\infty \leq |\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0,$$

which implies

$$\|y(x)\|_\infty \leq (|\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0)(1 - \rho)^{-1}$$

since $\rho = \alpha^{-1} \|b\|_{\infty,2} < 1$.

Hence we have $\|y(x)\|_\infty \leq C_0$, where $C_0 = (|\phi(0)| + |\beta| + \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0)(1 - \rho)^{-1}$.

Now we prove estimate (4).

Since

$$\begin{aligned} \int_0^2 e^{\frac{-1}{\varepsilon} \int_0^\tau a(\eta) d\eta} d\tau &\geq \int_0^2 e^{\frac{-a^* \tau}{\varepsilon}} d\tau = \frac{\varepsilon}{a^*} (1 - e^{\frac{-2a^*}{\varepsilon}}) \\ &\geq \frac{\varepsilon}{a^*} (1 - e^{\frac{-2a^*}{\varepsilon}}) \\ &\equiv c_0 \varepsilon, \end{aligned}$$

where $c_0 = \frac{1}{a^*} (1 - e^{\frac{-2a^*}{\varepsilon}}) < 1$, $a^* = \|a\|_\infty$.

Consider from (7) that we have

$$\begin{aligned} &\frac{1}{\varepsilon} \int_0^2 F(\xi) d\xi \int_\xi^2 e^{\frac{-1}{\varepsilon} \int_\xi^\tau a(\eta) d\eta} d\tau \\ &\leq \frac{1}{\varepsilon} \int_0^2 |F(\xi)| d\xi \int_\xi^2 e^{\frac{-\alpha(\tau-\xi)}{\varepsilon}} d\tau \\ &\leq \frac{1}{\varepsilon} \int_0^2 |F(\xi)| d\xi \{ \alpha^{-1} \varepsilon [1 - e^{\frac{-\alpha}{\varepsilon} (2-\xi)}] \} \\ &\leq \alpha^{-1} \int_0^2 |F(\xi)| d\xi \end{aligned}$$

$$\leq \alpha^{-1} \|f\|_1 + \alpha^{-1} \|b\|_{\infty,1} \|\phi\|_0 + \alpha^{-1} \|b\|_{\infty,2} C_0 \equiv C_1.$$

Substituting this in (7), we get

$$\begin{aligned} |y'(0)| &\leq \frac{|\beta| + |\phi(0)| + \frac{1}{\varepsilon} \int_0^2 |F(\xi)| d\xi \int_{\xi}^2 e^{-\frac{1}{\varepsilon} \int_{\xi}^{\tau} a(\eta) d\eta} d\tau}{\int_0^2 e^{-\frac{1}{\varepsilon} \int_0^{\tau} a(\eta) d\eta} d\tau} \\ &\leq \frac{(|\beta| + |\phi(0)| + C_1)}{c_0 \varepsilon} \equiv \frac{C_2}{\varepsilon}, \quad \text{where } C_2 = \frac{(|\beta| + |\phi(0)| + C_1)}{c_0}. \end{aligned}$$

Using the procedure in (5), we get

$$\begin{aligned} |y'(x)| &\leq \frac{C_2}{\varepsilon} e^{-\frac{1}{\varepsilon} \int_0^x \alpha d\eta} + \frac{1}{\varepsilon} \int_0^x |F(\xi)| e^{-\frac{1}{\varepsilon} \int_{\xi}^x \alpha d\eta} d\xi \\ &\leq \frac{C_2}{\varepsilon} e^{-\frac{\alpha x}{\varepsilon}} + C_1 \\ &\leq \frac{C_2}{\varepsilon} e^{-\frac{\alpha x}{\varepsilon}} + C_2 c_0 - |\beta| - |\phi(0)| \\ &\leq \frac{C_2}{\varepsilon} e^{-\frac{\alpha x}{\varepsilon}} + C_2, \quad \text{since } c_0 < 1. \end{aligned}$$

Therefore we have

$$|y'(x)| \leq C_2 \left(1 + \frac{1}{\varepsilon} e^{-\alpha \frac{x}{\varepsilon}} \right), \quad 0 \leq x \leq 2. \quad \square$$

3 Derivation of the method

Let $x_0 = 0, x_N = 2, x_i = ih, h = 2/N$.

A function $s(x, \tau) = s(x)$ satisfying in $[x_i, x_{i+1}]$ the differential equations is

$$s''(x) + \tau s(x) = \left[s''(x_i) + \tau s(x_i) \right] \frac{(x_{i+1} - x)}{h} + \left[s''(x_{i+1}) + \tau s(x_{i+1}) \right] \frac{(x - x_i)}{h}, \tag{12}$$

where $s(x_i) = y_i$ and $\tau > 0$ is termed cubic spline in compression.

Solving (12) as a linear second order differential equation, we get

$$s(x) = A \cos \frac{\lambda}{h} x_i + B \sin \frac{\lambda}{h} x_i + \left(\frac{M_i + \tau y_i}{\tau} \right) \left(\frac{x_{i+1} - x}{h} \right) \left(\frac{M_{i+1} + \tau y_{i+1}}{\tau} \right) \left(\frac{x - x_i}{h} \right).$$

We can find the arbitrary constants A and B by using interpolatory conditions

$$s(x_{i+1}) = y_{i+1}, \quad s(x_i) = y_i.$$

Writing $\lambda = h\tau^{1/2}$ and $M_i = s''(x_i)$, we get

$$\begin{aligned} s(x) &= \frac{-h^2}{\lambda^2 \sin \lambda} \left[M_{i+1} \sin \frac{\lambda(x - x_i)}{h} + M_i \sin \frac{\lambda(x_{i+1} - x)}{h} \right] \\ &\quad + \frac{h^2}{\lambda^2} \left[\frac{(x - x_i)}{h} \left(M_{i+1} + \frac{\lambda^2}{h^2} y_{i+1} \right) + \frac{(x_{i+1} - x)}{h} \left(M_i + \frac{\lambda^2}{h^2} y_i \right) \right]. \end{aligned} \tag{13}$$

Differentiating equation (13) and equating the left- and right-hand derivatives at x_i , we have

$$\begin{aligned} & \frac{y_i - y_{i-1}}{h} + \frac{h}{\lambda^2} \left[(1 - \lambda \cot \lambda)M_i - \left(1 - \frac{\lambda}{\sin \lambda}\right)M_{i-1} \right] \\ &= \frac{y_{i+1} - y_i}{h} + \frac{h}{\lambda^2} \left[\left(1 - \frac{\lambda}{\sin \lambda}\right)M_{i+1} - (1 - \lambda \cot \lambda)M_i \right]. \end{aligned} \tag{14}$$

This leads to a tridiagonal system

$$h^2(\lambda_1 M_{i-1} + 2\lambda_2 M_i + \lambda_1 M_{i+1}) = y_{i+1} - 2y_i + y_{i-1}, \quad i = 1, 2, \dots, 2N - 1, \tag{15}$$

where

$$\lambda_1 = \frac{1}{\lambda^2} \left(\frac{\lambda}{\sin \lambda} - 1 \right), \quad \lambda_2 = \frac{1}{\lambda^2} (1 - \lambda \cot \lambda).$$

The condition of continuity given by (15) ensures the continuity of first order derivatives of the spline $s(x, \tau)$ at interior points.

Substituting, $\varepsilon M_i = -a(x_i)y'_i - b(x_i)y(x_i - 1) + f(x_i)$ in equation (15) and using the following approximations for first order derivative of y :

$$\begin{aligned} y'_i &\cong \frac{(y_{i+1} - y_{i-1})}{2h}, \\ y'_{i+1} &\cong \frac{(3y_{i+1} - 4y_i + y_{i-1})}{2h}, \\ y'_{i-1} &\cong \frac{(-y_{i+1} + 4y_i - 3y_{i-1})}{2h}, \end{aligned}$$

we get the following tridiagonal linear system:

$$\begin{aligned} & \left(-\varepsilon + \frac{3}{2}\lambda_1 h a_{i-1} + \lambda_2 h a_i - \frac{\lambda_1}{2} h a_{i+1} \right) y_{i-1} + (2\varepsilon - 2\lambda_1 h a_{i-1} + 2\lambda_1 h a_{i+1}) y_i \\ &+ \left(-\varepsilon + \frac{\lambda_1}{2} h a_{i-1} - \lambda_2 h a_i - \frac{3}{2}\lambda_1 h a_{i+1} \right) y_{i+1} \\ &= -h^2 \left[\lambda_1 (f_{i-1} - b_{i-1} y_{i-1-N}) + 2\lambda_2 (f_i - b_i y_{i-N}) \right. \\ &\quad \left. + \lambda_1 (f_{i+1} - b_{i+1} y_{i+1-N}) \right], \quad i = 1, 2, \dots, 2N - 1. \end{aligned} \tag{16}$$

4 Numerical algorithm

Step 1. We obtain the reduced problem by setting $\varepsilon = 0$ in equation (1) with an appropriate interval condition. Let $y_0(x)$ be the solution of the reduced problem of (1) and (2), i.e.,

$$a(x)y'_0(x) + b(x)y_0(x - 1) = f(x) \tag{17}$$

with

$$y_0(x) = \phi(x), \quad -1 \leq x \leq 0. \tag{18}$$

We solve (17) and (18) by using the classical Runge-Kutta method of order four in $0 \leq x \leq 1$.

We consider $y_0(1) = \gamma$.

Step 2. To obtain the solution in $0 < x < 1$, we consider the numerical scheme from (16) with a fitting factor

$$\sigma(\rho) = \frac{a_i \rho}{2} \operatorname{Coth}\left(\frac{a_i \rho}{2}\right), \quad \text{where } \rho = \frac{h}{\varepsilon} \text{ (cf. [42]).}$$

Scheme (16) with a fitting factor can be written as

$$E_i y_{i-1} - F_i y_i + G_i y_{i+1} = H_i, \quad 1 < i < N - 1,$$

where

$$\begin{aligned} E_i &= -\varepsilon \sigma + \frac{3}{2} \lambda_1 h a_{i-1} + \lambda_2 h a_i - \frac{\lambda_1}{2} h a_{i+1}, & F_i &= -(2\varepsilon \sigma - 2\lambda_1 h a_{i-1} + 2\lambda_1 h a_{i+1}), \\ G_i &= -\varepsilon \sigma + \frac{\lambda_1}{2} h a_{i-1} - \lambda_2 h a_i - \frac{3}{2} \lambda_1 h a_{i+1} \quad \text{and} \\ H_i &= -h^2 (\lambda_1 (f_{i-1} - b_{i-1} \phi_{i-1-N}) + 2\lambda_2 (f_i - b_i \phi_{i-N}) + \lambda_1 (f_{i+1} - b_{i+1} \phi_{i+1-N})). \end{aligned}$$

We solve this system by Thomas algorithm with the boundary conditions

$$y_0 = \phi(0) \quad \text{and} \quad y_N = \gamma.$$

Similarly, to obtain the solution in $1 < x < 2$, we rewrite the numerical scheme with the fitting factor as:

$$E_i y_{i-1} - F_i y_i + G_i y_{i+1} = H_i, \quad N + 1 < i < 2N - 1,$$

where

$$\begin{aligned} E_i &= -\varepsilon \sigma + \frac{3}{2} \lambda_1 h a_{i-1} + \lambda_2 h a_i - \frac{\lambda_1}{2} h a_{i+1}, & F_i &= -(2\varepsilon \sigma - 2\lambda_1 h a_{i-1} + 2\lambda_1 h a_{i+1}), \\ G_i &= -\varepsilon \sigma + \frac{\lambda_1}{2} h a_{i-1} - \lambda_2 h a_i - \frac{3}{2} \lambda_1 h a_{i+1} \quad \text{and} \\ H_i &= -h^2 (\lambda_1 (f_{i-1} - b_{i-1} y_{i-1-N}) + 2\lambda_2 (f_i - b_i y_{i-N}) + \lambda_1 (f_{i+1} - b_{i+1} y_{i+1-N})). \end{aligned}$$

We solve the system with the boundary conditions

$$y_N = \gamma \quad \text{and} \quad y_{2N} = \beta.$$

5 Numerical examples

To demonstrate the applicability of the method, we consider one boundary value problem of singularly perturbed linear differential difference equations exhibiting boundary layer at the left end of the interval $[0, 2]$ and four boundary value problems with right-end boundary layer. These problems were widely discussed in the literature. The numerical results are presented for $\lambda_1 = \frac{1}{18}, \lambda_2 = \frac{4}{9}$.

Since the exact solutions of the problems are not known, the maximum absolute errors for the examples are calculated using the following double mesh principle:

$$E_\varepsilon^N = \max_{0 \leq i \leq N} |y_i^N - y_{2i}^{2N}|.$$

For a value of N , the ϵ -uniform maximum absolute error is calculated by the formula $E^N = \max_{\epsilon} E_{\epsilon}^N$.

The numerical rate of convergence for all the examples has been calculated by the formula

$$R^N = \frac{\log |E_{\epsilon}^N / E_{\epsilon}^{2N}|}{\log 2}.$$

Example 1 ([24], p.2357) $\epsilon y''(x) + 128y'(x) + 0.25y(x - 1) = 0.25(x - 1)$, $0 < x < \frac{3}{2}$, $y(x) = x$, $-1 \leq x \leq 0$, $y(\frac{3}{2}) = 2$.

The numerical results are presented in Table 1 for different vales of perturbation parameter ϵ .

Example 2 ([23], p.247) $\epsilon y''(x) - 3y'(x) + y(x - 1) = 0$, $y(x) = 1$, $-1 \leq x \leq 0$, $y(2) = 2$.

Table 1 The maximum absolute errors for Example 1 for different values of ϵ

$\epsilon \downarrow \setminus N \rightarrow$	32	64	128	256	512	1,024
2^{-5}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	1.967E-06
2^{-6}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	7.480E-07
2^{-7}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.579E-07
2^{-8}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-9}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-10}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-11}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-12}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-13}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-14}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-15}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-16}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-17}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-18}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-19}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
2^{-20}	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
E^N	1.017E-05	5.086E-06	2.543E-06	1.272E-06	6.358E-07	5.544E-07
R^N	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.976E-01	1.284E+00

Table 2 The maximum absolute errors for Example 2 for different values of ϵ

$\epsilon \downarrow \setminus N \rightarrow$	32	64	128	256	512	1,024
2^{-5}	5.155E-04	1.465E-04	3.789E-05	9.552E-06	2.403E-06	8.200E-07
2^{-6}	7.592E-04	2.664E-04	7.515E-05	1.945E-05	4.911E-06	1.202E-06
2^{-7}	8.368E-04	3.857E-04	1.353E-04	3.818E-05	9.876E-06	2.470E-06
2^{-8}	8.409E-04	4.251E-04	1.944E-04	6.821E-05	1.924E-05	4.963E-06
2^{-9}	8.409E-04	4.272E-04	2.143E-04	9.756E-05	3.425E-05	9.658E-06
2^{-10}	8.409E-04	4.272E-04	2.153E-04	1.075E-04	4.892E-05	1.717E-05
2^{-11}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.387E-05	2.450E-05
2^{-12}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.694E-05
2^{-13}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-14}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-15}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-16}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-17}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-18}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-19}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
2^{-20}	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
E^N	8.409E-04	4.272E-04	2.153E-04	1.081E-04	5.414E-05	2.707E-05
R^N	9.769E-01	9.885E-01	9.947E-01	9.969E-01	9.998E-01	8.436E-01

Table 3 The maximum absolute errors for Example 3 for different values of ϵ

$\epsilon \downarrow \ N \rightarrow$	32	64	128	256	512	1,024
2^{-5}	2.074E-02	5.496E-03	1.395E-03	3.503E-04	8.744E-05	4.587E-05
2^{-6}	3.534E-02	1.073E-02	2.853E-03	7.243E-04	1.816E-04	6.401E-05
2^{-7}	4.558E-02	1.801E-02	5.495E-03	1.456E-03	3.697E-04	1.040E-04
2^{-8}	4.727E-02	2.316E-02	9.148E-03	2.781E-03	7.368E-04	1.916E-04
2^{-9}	4.730E-02	2.402E-02	1.167E-02	4.611E-03	1.399E-03	3.722E-04
2^{-10}	4.730E-02	2.403E-02	1.210E-02	5.860E-03	2.314E-03	7.028E-04
2^{-11}	4.730E-02	2.403E-02	1.211E-02	6.076E-03	2.935E-03	1.160E-03
2^{-12}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.044E-03	1.473E-03
2^{-13}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.526E-03
2^{-14}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-15}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-16}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-17}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-18}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-19}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
2^{-20}	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
E^N	4.730E-02	2.403E-02	1.211E-02	6.080E-03	3.046E-03	1.527E-03
R^N	9.769E-01	9.886E-01	9.941E-01	9.972E-01	9.965E-01	1.002E+00

Table 4 The maximum absolute errors for Example 4 for different values of ϵ

$\epsilon \downarrow \ N \rightarrow$	32	64	128	256	512	1,024
2^{-5}	1.299E-03	6.461E-04	2.673E-04	9.036E-05	2.661E-05	7.234E-06
2^{-6}	1.273E-03	6.528E-04	3.238E-04	1.339E-04	4.524E-05	1.331E-05
2^{-7}	1.269E-03	6.381E-04	3.272E-04	1.621E-04	6.701E-05	2.263E-05
2^{-8}	1.269E-03	6.362E-04	3.195E-04	1.638E-04	8.113E-05	3.352E-05
2^{-9}	1.269E-03	6.362E-04	3.186E-04	1.598E-04	8.194E-05	4.058E-05
2^{-10}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.994E-05	4.098E-05
2^{-11}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.998E-05
2^{-12}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-13}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-14}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-15}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-16}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-17}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-18}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-19}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
2^{-20}	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
E^N	1.269E-03	6.362E-04	3.186E-04	1.593E-04	7.971E-05	3.986E-05
R^N	9.958E-01	9.978E-01	9.996E-01	9.991E-01	9.996E-01	7.166E-01

The numerical results are presented in Table 2 for different vales of perturbation parameter ϵ .

Example 3 ([23], p.247) $\epsilon y''(x) - 2y'(x) + 5y(x - 1) = 0, y(x) = 1, -1 \leq x \leq 0, y(2) = 2.$

The numerical results are presented in Table 3 for different vales of perturbation parameter ϵ .

Example 4 ([23], p.247) $\epsilon y''(x) - 5y'(x) + \frac{1}{2}y(x - 1) = \begin{cases} -1, & 0 \leq x \leq 1, \\ 1, & 1 \leq x \leq 2, \end{cases} y(x) = 1, -1 \leq x \leq 0, y(2) = 2.$

The numerical results are presented in Table 4 for different vales of perturbation parameter ϵ .

Example 5 ([23], p.247) $\epsilon y''(x) - (x + 10)y'(x) + y(x - 1) = -x, y(x) = x, -1 \leq x \leq 0, y(2) = 2.$

Table 5 The maximum absolute errors for Example 5 for different values of ϵ

$\epsilon \downarrow \ N \rightarrow$	32	64	128	256	512	1,024
2^{-5}	1.576E-03	9.904E-03	1.478E-02	1.051E-02	5.821E-03	3.000E-03
2^{-6}	2.743E-03	9.310E-04	1.032E-02	1.511E-02	1.067E-02	5.887E-03
2^{-7}	2.766E-03	1.378E-03	1.304E-03	1.053E-02	1.528E-02	1.075E-02
2^{-8}	2.766E-03	1.402E-03	6.817E-04	1.656E-03	1.063E-02	1.538E-02
2^{-9}	2.766E-03	1.402E-03	7.056E-04	3.299E-04	1.832E-03	1.074E-02
2^{-10}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.531E-04	1.920E-03
2^{-11}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	6.508E-05
2^{-12}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-13}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-14}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-15}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-16}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-17}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-18}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-19}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
2^{-20}	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
E^N	2.766E-03	1.402E-03	7.056E-04	3.540E-04	1.772E-04	8.922E-05
R^N	9.806E-01	9.904E-01	9.952E-01	9.979E-01	9.902E-01	9.943E-01

The numerical results are presented in Table 5 for different vales of perturbation parameter ϵ .

6 Discussion and conclusions

In this paper we present an exponentially fitted finite difference scheme to solve singularly perturbed delay differential equation of second order with large delay. The method is based on cubic spline in compression. We have implemented the present method on one linear example with left-end boundary layer and four examples with right-end boundary layer by taking different values of ϵ . Numerical results are presented in tables. From the results, it can be observed that as the grid size h decreases, the maximum absolute errors decrease, which shows the convergence to the computed solution. On the basis of the numerical results of a variety of examples, it is concluded that the present method offers significant advantage for the linear singularly perturbed delay differential equations with large delays.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Visvesvaraya National Institute of Technology, Nagpur, 440010, India. ²School of Advanced Sciences, VIT University, Vellore, Tamilnadu 632014, India. ³Department of Aeronautical Engineering, ADCET, Sangli, Ashta, India.

Authors' information

P Pramod Chakravarthy received MSc and PhD from National Institute of Technology, Warangal, India. He is currently working as an Associate Professor and Head of the Department of Mathematics at Visvesvaraya National Institute of Technology, Nagpur, India. Since 2006 he has been at Visvesvaraya National Institute of Technology, Nagpur. His research interests include numerical methods for singular perturbation problems in differential equations and delay differential equations. Mr Dinesh Kumar is presently pursuing his PhD in Visvesvaraya National Institute of Technology, Nagpur, under the supervision of P Pramod Chakravarthy. R Nageshwar Rao received MSc from National Institute of Technology, Warangal, India and PhD from Visvesvaraya National Institute of Technology, Nagpur, India, under the supervision of P Pramod Chakravarthy. Presently he is working as an Assistant Professor in Mathematics at VIT University, Vellore, India - 632014. His research interests include numerical methods for singularly perturbed differential-difference equations. Devendra P Ghate received his PhD from Oxford University. Presently he is working as an Associate Professor in Aeronautical Engineering Department at ADCET, Ashta, Sangle, India.

Acknowledgements

The authors wish to thank the National Board for Higher Mathematics, Department of Atomic Energy, Government of India, for their financial support under the project No. NBHM/R.P. 37/2012/Fresh/1742 dated 15th November, 2012. The authors are grateful to the referees for their careful reading of the manuscript and valuable comments. The authors thank for the help from the editor too.

Received: 22 May 2015 Accepted: 8 September 2015 Published online: 21 September 2015

References

- Bellman, R, Cooke, KL: *Differential-Difference Equations*. Academic Press, New York (1963)
- Driver, RD: *Ordinary and Delay Differential Equations*. Springer, New York (1977)
- El'sgol'ts, LE, Norkin, SB: *Introduction to the Theory and Application of Differential Equations with Deviating Arguments*. Mathematics in Science and Engineering. Academic Press, San Diego (1973)
- Erneux, T: *Applied Delay Differential Equation*. Springer, New York (2009)
- Gopalsamy, K: *Stability and Oscillations in Delay Differential Equations of Population Dynamics*. Kluwer Academic, Dordrecht (1992)
- Györi, I, Ladas, G: *Oscillation Theory of Delay Equations with Applications*. Clarendon, Oxford (1991)
- Halanay, A: *Differential Equations: Stability, Oscillations, Time Lags*. Mathematics in Science and Engineering. Academic Press, San Diego (1996)
- Kuang, Y: *Delay Differential Equations with Applications in Population Dynamics*. Academic Press, New York (1993)
- Smith, H: *An Introduction to Delay Differential Equations with Applications to the Life Sciences*. Springer, Berlin (2010)
- Feldstein, MA, Stetter, HJ: *Simplified predictor-corrector methods*. Numerical analysis research technical report, University of California, Los Angeles (1963)
- Bellen, A, Zennaro, M: *Numerical Methods for Delay Differential Equations*. Oxford Science Publications, New York (2003)
- Derstine, MW, Gibbs, FAHMM, Kaplan, DL: Bifurcation gap in a hybrid optical system. *Phys. Rev. A* **26**, 3720-3722 (1982)
- Mallet-Paret, J, Nussbaum, RD: A differential-delay equations arising in optics and physiology. *SIAM J. Math. Anal.* **20**, 249-292 (1989)
- Longtin, A, Milton, J: Complex oscillations in the human pupil light reflex with mixed and delayed feedback. *Math. Biosci.* **90**, 183-199 (1988)
- Mackey, MC, Glass, L: Oscillation and chaos in physiological control systems. *Science* **197**, 287-289 (1977)
- Glizer, VY: Asymptotic solution of a boundary-value problem for linear singularly-perturbed functional differential equations arising in optimal control theory. *J. Optim. Theory Appl.* **106**, 309-335 (2000)
- Glizer, VY: Blockwise estimate of the fundamental matrix of linear singularly perturbed differential systems with small delay and its application to uniform asymptotic solution. *J. Math. Anal. Appl.* **278**, 409-433 (2003)
- Stein, RB: A theoretical analysis of neuronal variability. *Biophys. J.* **5**, 173-194 (1965)
- Tuckwell, HC: On the first-exit time problem for temporally homogeneous Markov processes. *J. Appl. Probab.* **13**, 39-48 (1976)
- Tuckwell, HC: *Introduction to Theoretical Neurobiology*, vol. 1. Cambridge University Press, Cambridge (1988)
- Tuckwell, HC: *Introduction to Theoretical Neurobiology*, vol. 2. Cambridge University Press, Cambridge (1988)
- Wilbur, WJ, Rinzel, J: An analysis of Stein's model for stochastic neuronal excitation. *Biol. Cybern.* **45**, 107-114 (1982)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary-value problems for differential-difference equations. *SIAM J. Appl. Math.* **42**, 502-531 (1982)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary-value problems for differential-difference equations. II. Rapid oscillations and resonances. *SIAM J. Appl. Math.* **45**, 687-707 (1985)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary-value problems for differential-difference equations. III. Turning point problems. *SIAM J. Appl. Math.* **45**, 708-734 (1985)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary value problems for differential difference equations. IV. A nonlinear example with layer behavior. *Stud. Appl. Math.* **84**, 231-273 (1991)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary-value problems for differential-difference equations. V. Small shifts with layer behavior. *SIAM J. Appl. Math.* **54**, 249-272 (1994)
- Lange, CG, Miura, RM: Singular perturbation analysis of boundary-value problems for differential-difference equations. VI. Small shifts with rapid oscillations. *SIAM J. Appl. Math.* **54**, 273-283 (1994)
- Kadalbajoo, MK, Sharma, KK: Numerical analysis of boundary value problems for singularly perturbed differential difference equations with small shifts of mixed type. *J. Optim. Theory Appl.* **115**(1), 145-163 (2002)
- Kadalbajoo, MK, Sharma, KK: Numerical analysis of singularly perturbed delay differential equations with layer behavior. *Appl. Math. Comput.* **157**, 11-28 (2004)
- Kadalbajoo, MK, Sharma, KK: Numerical analysis of boundary value problems for singularly perturbed differential difference equations: small shifts of mixed type with rapid oscillations. *Commun. Numer. Methods Eng.* **20**, 167-182 (2004)
- Kadalbajoo, MK, Sharma, KK: ε -Uniformly fitted mesh method for singularly perturbed differential difference equations: mixed type of shifts with layer behavior. *Int. J. Comput. Math.* **81**(1), 49-62 (2004)
- Kadalbajoo, MK, Sharma, KK: Numerical treatment of a mathematical model arising from a model of neuronal variability. *J. Math. Anal. Appl.* **307**, 606-627 (2005)
- Kadalbajoo, MK, Sharma, KK: Numerical treatment of boundary value problems for second order singularly perturbed delay differential equations. *Comput. Appl. Math.* **24**(2), 151-172 (2005)
- Kadalbajoo, MK, Sharma, KK: Numerical treatment for singularly perturbed nonlinear differential difference equations with negative shift. *Nonlinear Anal.* **63**, e1909-e1924 (2005)
- Kadalbajoo, MK, Sharma, KK: Parameter uniform fitted mesh method for singularly perturbed delay differential equations with layer behavior. *Electron. Trans. Numer. Anal.* **23**, 180-201 (2006)

37. Kadalbajoo, MK, Sharma, KK: An ε -uniform convergent method for a general boundary value problem for singularly perturbed differential difference equations: small shifts of mixed type with layer behavior. *J. Comput. Methods Sci. Eng.* **6**(1), 39-55 (2006)
38. Kadalbajoo, MK, Sharma, KK: A numerical method based on finite difference for boundary value problems for singularly perturbed delay differential equations. *Appl. Math. Comput.* **197**, 692-707 (2008)
39. Kadalbajoo, MK, Kumar, D: A computational method for singularly perturbed nonlinear differential-difference equations with small shift. *Appl. Math. Model.* **34**, 2584-2596 (2010)
40. Kadalbajoo, MK, Ramesh, VP: Hybrid method for numerical solution of singularly perturbed delay differential equations. *Appl. Math. Comput.* **187**, 797-814 (2007)
41. Gulsu, M, Sezer, M: A Taylor polynomial approach for solving differential-difference equations. *J. Comput. Appl. Math.* **186**, 349-364 (2006)
42. Doolan, ER, Miller, JJH, Schilders, WHA: *Uniform Numerical Methods for Problems with Initial and Boundary Layers*. Boole, Dublin (1980)
43. Miller, JJH, O'Riordan, E, Shishkin, GI: *Fitted Numerical Methods for Singular Perturbation Problems*. World Scientific, Singapore (1996)
44. Roos, HG, Stynes, M, Tobiska, L: *Numerical Methods for Singularly Perturbed Differential Equations*. Springer, Berlin (1996)
45. Farrell, PA, Hegarty, AF, Miller, JJH, O'Riordan, E, Shishkin, GI: *Robust Computational Techniques for Boundary Layers*. Chapman & Hall/CRC, New York (2000)
46. Bickley, WG: Piecewise cubic interpolation and two point boundary value problems. *Comput. J.* **11**, 206-208 (1968)
47. Aziz, T, Khan, A: A spline method for second order singularly perturbed boundary value problems. *J. Comput. Appl. Math.* **147**, 445-452 (2002)
48. Amiraliyev, GM, Erdogan, F: Uniform numerical method for singularly perturbed delay differential equations. *Comput. Math. Appl.* **53**, 1251-1259 (2007)
49. Amiraliyev, GM, Cimen, E: Numerical method for a singularly perturbed convection-diffusion problem with delay. *Appl. Math. Comput.* **216**, 2351-2359 (2010)
50. Amiraliyeva, IG, Erdogan, F, Amiraliyev, GM: A uniform numerical method for dealing with a singularly perturbed delay initial value problem. *Appl. Math. Lett.* **23**, 1221-1225 (2010)
51. Erdogan, F, Amiraliyev, GM: Fitted finite difference method for singularly perturbed delay differential equations. *Numer. Algorithms* **59**, 131-145 (2012)
52. Subburayan, V, Ramanujam, N: An initial value technique for singularly perturbed convection - diffusion problems with a negative shift. *J. Optim. Theory Appl.* **158**, 234-250 (2013)
53. Ghomanjani, F, Farahi, MH, Kamyad, AV: Numerical solution of some linear optimal control systems with pantograph delays. *IMA J. Math. Control Inf.* **32**(2), 225-243 (2015). doi:10.1093/imamci/dnt037
54. Ghomanjani, F, Farahi, MH, Kılıçman, A, Kamyad, AV, Pariz, N: Bezier curves based numerical solutions of delay systems with inverse time. *Math. Probl. Eng.* (2014). doi:10.1155/2014/602641

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
