

A Kinetic Database for Astrochemistry (KIDA)

V. Wakelam^{1,2}, E. Herbst^{3,4}, J.-C. Loison^{5,6}, I. W. M. Smith⁷, V. Chandrasekaran^{5,6}, B. Pavone^{1,2}, N. G. Adams⁸, M.-C. Bacchus-Montabonel⁹, A. Bergeat^{5,6}, K. Béroff¹⁰, V. M. Bierbaum¹¹, M. Chabot¹², A. Dalgarno¹³, E. F. van Dishoeck¹⁴, A. Faure¹⁵, W. D. Geppert¹⁶, D. Gerlich¹⁷, D. Galli¹⁸, E. Hébrard^{1,2}, F. Hersant^{1,2}, K. M. Hickson^{5,6}, P. Honvault^{19,20}, S. J. Klippenstein²¹, S. Le Picard²², G. Nyman²³, P. Pernot²⁴, S. Schlemmer²⁵, F. Selsis^{1,2}, I. R. Sims²², D. Talbi²⁶, J. Tennyson²⁷, J. Troe^{28,29}, R. Wester³⁰, L. Wiesenfeld¹⁵

ABSTRACT

We present a novel chemical database for gas-phase astrochemistry. Named the Kinetic Database for Astrochemistry (KIDA), this database consists of gas-phase reactions with rate coefficients and uncertainties that will be vetted to the greatest extent possible. Submissions of measured and calculated rate coefficients are welcome, and will be studied by experts before inclusion into the database. Besides providing kinetic information for the interstellar medium, KIDA is planned to contain such data for planetary atmospheres and for circumstellar envelopes. Each year, a subset of the reactions in the database (*kida.uva*) will be provided as a network for the simulation of the chemistry of dense interstellar clouds with temperatures between 10 K and 300 K. We also provide a code, named Nahoon, to study the time-dependent gas-phase chemistry of 0D and 1D interstellar sources.

Subject headings: Physical data and processes: astrochemistry — Astronomical databases: miscellaneous — ISM: abundances, molecules

¹Univ. Bordeaux, LAB, UMR 5804, F-33270, Floirac, France

²CNRS, LAB, UMR 5804, F-33270, Floirac, France

³Departments of Physics, Astronomy, and Chemistry, The Ohio State University, Columbus, OH 43210 USA

⁴Departments of Chemistry, Astronomy, and Physics, University of Virginia, Charlottesville, VA 22904 USA

⁵Univ. Bordeaux, ISM, CNRS UMR 5255, F-33400 Talence, France

⁶CNRS, ISM, CNRS UMR 5255, F-33400 Talence, France

⁷University Chemical Laboratories, Lensfield Road, Cambridge CB2 1EW, UK

⁸Department of Chemistry, University of Georgia, Athens, GA 30602, USA

⁹Université de Lyon (Lyon I), LASIM, CNRS-UMR5579, 43 Bd 11 Novembre 1918, 69622 Villeurbanne Cedex, France

¹⁰Institut des Sciences Moléculaires d'Orsay, CNRS and Université Paris-Sud, 91405 Orsay cedex, France

¹¹Department of Chemistry and Biochemistry, Center for Astrophysics and Space Astronomy, University of Colorado, Boulder, CO 80309, United States

¹²Intitut de Physique Nucléaire d'Orsay, IN2P3-CNRS and Université Paris-Sud, 91406 Orsay cedex, France

¹³Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

¹⁴Leiden Observatory, Leiden University, P.O. Box 9513, 2300

RA Leiden, The Netherlands

¹⁵UJF-Grenoble 1/CNRS, Institut de Planétologie et d'Astrophysique de Grenoble (IPAG) UMR 5274, Grenoble, F-38041, France

¹⁶Department of Physics, University of Stockholm, Roslagstullbacken 21, S-10691 Stockholm

¹⁷Technische Universität Chemnitz, Department of Physics, 09107 Chemnitz, Germany

¹⁸INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125, Firenze, Italy

¹⁹Laboratoire Interdisciplinaire Carnot de Bourgogne, UMR CNRS 5209, Université de Bourgogne, 21078 Dijon Cedex, France

²⁰UFR Sciences et Techniques, Université de Franche-Comté, 25030 Besançon Cedex, France

²¹Chemical Sciences and Engineering Division, Argonne National Laboratory, Argonne, IL, 60439, USA

²²Institut de Physique de Rennes, Equipe Astrochimie Expérimentale UMR 6251 du CNRS - Université de Rennes 1, Bât. 11c, Campus de Beaulieu, 35042 Rennes Cedex, France

²³Department of Chemistry, University of Gothenburg, 41296 Gothenburg, Sweden

²⁴Laboratoire de Chimie Physique, UMR 8000, CNRS, Univ. Paris-Sud, F-91405 Orsay

²⁵I. Physikalisches Institut, University of Cologne, Zlpicher Str.

1. Introduction

The interstellar medium (ISM) is composed of gas and dust under a wide range of physical conditions. The diffuse interstellar medium is characterized by very low gas densities ($\leq 10^3 \text{ cm}^{-3}$), temperatures between 30 and 100 K, high UV photon fluxes and chemical species mainly in atomic form (except for molecular hydrogen). Due to gravitational interactions and turbulence, this material slowly forms denser ($\sim 10^4 \text{ cm}^{-3}$) and colder (10 K) regions through which ionizing cosmic rays can penetrate but UV photons cannot. Over a period of about 10^5 - 10^6 yr, these atoms and molecular hydrogen start a series of chemical reactions that synthesize molecules in the gas and on dust particles. Some of these molecules are large species, mainly in the form of unsaturated (H-poor) carbon-chains. Due to perturbations, these cold and dense regions, called dark molecular clouds or cold cores, collapse to begin the different stages of star and planet formation; these stages comprise a prestellar core, a protostar, a star with a protoplanetary disk, and then possibly a planetary system. As the physical evolution occurs, a chemical evolution also occurs in which the dominant larger molecules are no longer unsaturated carbon chains but more saturated organic and prebiotic species quite common on the earth. At the end of their lives, stars re-inject their material into the ISM, now enriched in heavy elements. This set of events is the life cycle of stars and interstellar matter.

The study of the composition and evolution of interstellar material is part of what we call astrochemistry. During the next ten years, many fascinating discoveries are expected in this field of research thanks in part to a number of new observational instruments including the Herschel Space Observatory (HSO), the Atacama Large Millimeter/Submillimeter Array (ALMA), as well as the expanded Very Large Array (eVLA), and the Stratospheric Observatory for Infrared Astronomy (SOFIA). HSO and SOFIA are currently observing

molecular spectra in the far-infrared, a frequency range not accessible from the ground, whereas the ALMA interferometer, now only in its early science stage, will give access to very high spectral resolution and sensitivity, with broadband spectral coverage as well as spatial resolution never achieved before. Many new molecules have been and will be detected, increasing our knowledge of the complexity of the ISM. Moreover, astrochemistry is important for the understanding of Sun-like star and planetary formation. Indeed, protostellar collapse is governed by quantities such as the ionization fraction (directly related to the chemical composition), which couples the gas to the magnetic field. Finally, the evolution of the chemical composition of the gas, dust, and ice in the protoplanetary disk is a key process for the understanding of planetary formation, the fate of complex organics acquired in earlier stages, as well as the emergence of life.

In order to understand the formation and destruction of molecules observed in assorted sources in the ISM, astrochemists model the chemical evolution through reaction networks. In these models, the chemistry is described by processes of various types: ion-neutral and neutral-neutral reactions, dissociation and ionization by UV photons and cosmic ray particles, dissociative recombination, etc. The models compute the abundances as a function of time for up to four hundred gas phase species using more than four thousand reactions, starting from an initial composition. Extensions that include isotopologues, anions, and large molecules contain many more reactions and molecules. For dark clouds, the initial composition can consist of abundances measured in more diffuse regions, mainly atoms and molecular hydrogen. Two main chemical databases of gas-phase chemistry currently exist in astrochemistry for the interstellar medium. The better known is the udfa database (previously called the UMIST database), created by Tom Millar (Queen's University of Belfast, Northern Ireland), first released in 1991 (Millar et al. 1991, 1997; Le Teuff et al. 2000; Woodall et al. 2007). Updated versions of the database have been released in 1995, 1999, and 2006 (<http://www.udfa.net/>). The second one is the OSU database, initially developed by Prasad and Huntress (1980), updated by Chun Leung and Eric Herbst, and then revised over the last twenty years by Herbst and collaborators. Several versions dating back to 2003 are available at <http://www.physics.ohio-state.edu/~eric/research.html>. Although the two databases are similar, some significant differences exist because

77, 50937 Cologne, Germany

²⁶Université Montpellier II - GRAAL, CNRS - UMR 5024, place Eugène Bataillon, 34095 Montpellier, France

²⁷Department of Physics and Astronomy, University College London, Gower St., London WC1E 6BT, UK

²⁸Institut für Physikalische Chemie, Universität Göttingen, Tammannstr. 6, D-37077 Göttingen, Germany

²⁹Max-Planck-Institut für Biophysikalische Chemie, Am Fassberg 11, D-37077 Göttingen, Germany

³⁰Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Technikerstraße 25, A-6020 Innsbruck, Austria

of occasionally different choices of which reactions to include; udfa includes more studied reactions whereas OSU relies more heavily on unstudied radical-neutral reactions. In addition, the two networks differ partially because of

1. Different approaches to estimate the rate coefficients of reactions not studied in the laboratory,
2. Different choices of experimental rate coefficients from uncritical compilations (the NIST database for instance ¹, and
3. Different approximations regarding the temperature dependence of ion-polar neutral reactions.

Until its most recent version, which is useable up to 800 K, the OSU database had been more relevant for cold sources (~ 10 K) whereas udfa includes many reactions with energy barriers for hotter conditions (> 100 K). Other accessible databases exist for particular cases such as Srates² for sulfur chemistry (Wakelam et al. 2004, 2011), a database for shock models (Pineau des Forets et al. 1993) and one for photon-dominated regions³ (Le Petit et al. 2006). The existence of these diverse databases, which contain many unstudied reactions, and often lack references, is a barrier to the improvement of chemical models. We have therefore initiated a project of constructing a unique and critical database for gas-phase astrochemistry named the Kinetic Database for Astrochemistry⁴ (KIDA). This database will include as many of the important gas-phase reactions for modeling various astronomical environments and planetary atmospheres as possible. Indeed, reaction networks relevant for planetary atmospheres require many of the same reactions as those in interstellar modeling, but over a different range of temperatures.

2. General presentation of the database

KIDA is designed in order to allow the efficient managing of data, user-friendly visualization of data and flexible downloading of lists of reactions. In addition to being able to consult the database, users can submit data to KIDA via an online interface.

¹<http://kinetics.nist.gov/kinetics/index.jsp>

²http://kida.obs.u-bordeaux1.fr/uploads/models/Srates_update.txt

³<http://pdr.obspm.fr/PDRDB/>

⁴<http://kida.obs.u-bordeaux1.fr>

2.1. Data on chemical species

Users can find two different types of data in KIDA. Some data are related to chemical species and others are related to chemical reactions. In the first type, one can find InChI codes, CAS registry numbers, electronic states, polarizabilities, dipole moments, and enthalpies of formation. InChI codes⁵ (for International Chemical Identifier) are character chains defined by IUPAC to identify any chemical species. The InChI code of OCS for instance is InChI=1S/COS/c2-1-3. These codes are needed for clear identification, especially in the context of interoperability of databases. Free software is available to draw the structure of the molecule from the InChI code. CAS registry numbers are numeric identifiers from the American Chemical Society. These numbers exist, however, only for chemical species used in industry. Most of these data in KIDA have been collected from other databases such as the NIST Chemistry Webbook⁶, CCCBDB⁷, and Woon & Herbst (2009). JPEG images of the structures of the species, drawn with the JMOL free software⁸, are also displayed.

2.2. Rate coefficients

For chemical reactions, rate coefficients are stored in the form of three parameters (A, B and C), which can be used to compute the rate coefficients using different formulae depending on the considered process and the range of temperatures desired. Rate coefficients define the intrinsic velocity of chemical reactions; the units in which they are expressed depends on the order of the reaction. The overall rate of a reaction is given by the product of the rate coefficient and the densities of the reactants. There are nine classes of reactions in KIDA, which are listed and described in Table 1. The first three classes can be thought of as unimolecular processes induced by cosmic rays or photons; here the rate coefficient has units of s^{-1} , and reaction rates are computed by multiplying the rate coefficient by the density (in cm^{-3}) of the reactant. The next five classes involve bimolecular collisions; in these classes rate coefficients have units of $cm^3 s^{-1}$, and the rate of reaction is computed by multiplying the rate coefficient by the densities (in cm^{-3}) of

⁵<http://www.iupac.org/inchi/>

⁶<http://webbook.nist.gov/chemistry/form-ser.html>

⁷<http://cccbdb.nist.gov/>

⁸<http://jmol.sourceforge.net/index.fr.html>

the two reactants. The last type is composed of termolecular (three-body) association reactions, which are processes where two reactants associate to form a product that is stabilized by collision with a third species, which can be a reactant or an inert atom or molecule. In the low-density limit, the rate of reaction is given by a rate coefficient in units of $\text{cm}^6 \text{s}^{-1}$ multiplied by the product of the densities of the three species involved in the reaction. A more general rate law involves a more complex dependence on density, which eventually saturates into a bimolecular rate law (Anicich 1993). Three-body association reactions are effective in “very dense” sources by interstellar if not terrestrial standards ($> 10^{11} \text{ cm}^{-3}$) and are generally considered in the atmospheres of planets. This type of reaction will be included later into KIDA.

The rate coefficients for bimolecular processes involving two heavy species or one heavy species and an electron are each given a certain range of temperature in which the parameters are applicable. By default, the rate coefficients are valid between 10 and 300 K. For updated rate coefficients, the temperature range is given by the results of experiments or the estimates of experts. Extrapolations outside this range of temperature are not recommended unless the rate coefficient is predicted to be totally independent of temperature, as occurs for the Langevin model of exothermic ion-non-polar-neutral reactions. Temperature dependent rate coefficients, at a specific temperature T , can often be fit to the Arrhenius-Kooij formula:

$$k_{Kooij}(T) = A(T/300)^B \exp(-C/T) \text{ cm}^3 \text{ s}^{-1}, \quad (1)$$

where A , B , and C are parameters. Several sets of rate coefficients for the same reaction can exist for different ranges of temperature or even the same range. If this happens for identified key reactions, KIDA experts can give recommendations on which values to use (see section 2.3).

Rate coefficients for unmeasured reactions between ions and neutral species with a dipole moment are computed using the Su-Chesnavich capture approach as discussed in Woon & Herbst (2009) and Wakelam et al. (2010b). Here the rate coefficient k_D is expressed in terms of the temperature-independent Langevin rate coefficient k_L using two formulas, one for lower and one for higher temperatures. The temperature ranges are divided by that temperature for which a parameter x , defined by the relation,

$$x = \frac{\mu_D}{(2\alpha kT)^{1/2}}, \quad (2)$$

where (in cgs-esu units) μ_D is the dipole moment and α the dipole polarizability, is equal to two. The formulae for k_D/k_L at the lower and higher temperature ranges are respectively

$$k_D/k_L = 0.4767x + 0.6200 \quad (3)$$

and

$$k_D/k_L = (x + 0.5090)^2/10.526 + 0.9754. \quad (4)$$

At the lowest temperatures in the region of validity for equation (3) ($T \geq 10 \text{ K}$), k_D goes as $T^{-1/2}$, whereas at the highest temperatures, k_D approaches the Langevin value. The rate coefficients are valid in the so-called semiclassical region, before they level off with decreasing temperature near 10 K or below (Troe 1996).

Rate coefficients for photo-processes by UV photons, k_{photo} , are normally computed as a function of the visual extinction A_V :

$$k_{photo} = A \exp(-CA_V) \text{ s}^{-1}. \quad (5)$$

The rate is obtained by multiplying k_{photo} by the density of the reactant. The parameter A for k_{photo} represents the unattenuated photodissociation rate in a reference radiation field whereas $\exp(-CA_V)$ takes into account the continuum attenuation from the dust. Since dust particles tend to attenuate light at shorter wavelengths more strongly, values of C are larger for species that have thresholds for photodissociation/photoionization farther into the UV (van Dishoeck 1988). Photoinduced rates depend on the external UV radiation field assumed. For example, the so-called average interstellar radiation field differs from the radiation field around a specific type of star. It is possible in KIDA to specify which field will be used. The data included in KIDA have been computed for the interstellar radiation field (Draine 1978). Data computed with other field will be included in the future.

Cosmic rays, chiefly protons and alpha particles with energies above 1 MeV, interact with gas-phase atoms and molecules in two ways. First, these high-energy particles can directly ionize a species. This direct process is dominant for H_2 , H, O, N, He and CO. Electrons produced in the direct process can cause secondary ionization before being thermalized. The secondary process is taken into account in most treatments by a factor of 5/3 (Dalgarno et al. 1999; Rimmer et al. 2012) while the effect of ionization by alpha particles

is taken account of by a separate factor of 1.8. The total ionization rate coefficient including these two factors is computed by

$$k_{CR,i} = A_i \zeta \quad (6)$$

where ζ (s^{-1}) incorporates the flux of cosmic rays and secondary electrons and normally refers to atomic hydrogen (Cravens & Dalgarno 1978). In the udfa, OSU, and KIDA networks, however, ζ refers to *molecular* hydrogen (ζ_{H_2}), and is approximately twice the value for atomic hydrogen. The rate coefficients for different species can be obtained by including a factor A_i .

In addition to secondary ionization, the electrons produced in direct cosmic ray ionization can electronically excite both molecular and atomic hydrogen. The radiative relaxation of H_2 (and H) produces UV photons that ionize and dissociate molecules. This indirect process is known as the Prasad-Tarafdar mechanism (Prasad & Tarafdar 1983). It represents a source of UV photons inside dense clouds. The rate coefficients of dissociation or ionization that proceed via excitation of molecular hydrogen are given by

$$k_{CRP,i} = \frac{A_i}{(1 - \omega)} \frac{n(H_2)}{(n(H) + 2n(H_2))} \zeta_{H_2} \quad (7)$$

with ζ_{H_2} the total ionization rate for H_2 , ω the albedo of the grains in the far ultraviolet, $n(H)$ the density of atomic hydrogen and $n(H_2)$ the density of molecular hydrogen (see also Gredel et al. 1989). In the OSU network (and subsequently in KIDA), it is assumed that ω is 0.5 and that the abundance of atomic hydrogen is negligible compared to molecular hydrogen. The rate then simplifies to $k_{CRP,i} = A_i \zeta_{H_2}$. This of course is only valid in dense sources. Unlike the case for direct cosmic ray ionization, the factor A_i can be quite large, often in the range $10^3 - 10^4$. The case of CO must be treated individually because of self-shielding (Gredel et al. 1987).

Any data provided in KIDA is associated with a reference. This reference can be bibliographic, another database, or a datasheet, as discussed in Section 2.3. All the reactions from the low-temperature OSU database osu-01-2009 have been included, even reactions involving anionic species (Harada & Herbst 2008). Many updates concerning photoinduced rates are currently being undertaken based on results from van Dishoeck et al. (2006) and van Hemert & van Dishoeck (2008). Branching ratios for the dissociation of carbon clusters by both external UV radiation and cos-

mic ray-induced photons have been added based on Chabot et al. (2010). Reactions necessary for primordial chemistry, collected by D. Galli, will be included before the end of 2012. Finally, many other additions have been made, in particular for neutral-neutral reactions, based on new experimental and/or theoretical results. All the original bibliographic references listed in KIDA are given in an online appendix to this paper.

2.3. Recommendations concerning rate coefficients

For all the rate coefficients in KIDA, a status is indicated. There are four possible recommendations, which are listed below:

0. Not recommended: This status suggests strongly that the reaction not be included.
1. Unknown: This status is mainly for reactions from other databases that have not been looked at by KIDA experts.
2. Valid: This is a status meant for reactions that are associated with a bibliographic reference.
3. Recommended: Recommended rate coefficients are mainly deduced from a variety of sources (experimental or theoretical) but can also just be educated guesses. The recommended status must be justified by a datasheet and can evolve with time when new data are available. Datasheets are pdf files stored in KIDA⁹ summarizing the information available on a reaction. These datasheets are inspired from the work done by the IUPAC subcommittee for Gas Kinetic Data Evaluation¹⁰. Datasheets, of which there are currently more than 40, are signed by one or more authors and are reviewed internally by KIDA experts.

2.4. Uncertainties for rate coefficients

The reliability of any reaction rate coefficient stored in KIDA is expressed in terms of an estimated uncertainty over the temperature range covered by the recommendation. This uncertainty is specified by three fields, two of which are shown in Table 2:

⁹http://kida.obs.u-bordeaux1.fr/publications#span_ datasheets

¹⁰<http://www.iupac-kinetic.ch.cam.ac.uk/>

1. The type of the statistical distribution representing the uncertainty. The preferred distribution, and the default, is the Lognormal (default = logn). This choice enforces the positivity of the rate coefficient, even for large uncertainty factors ($F_0 \geq 2$) (Thompson & Stewart 1991; Hébrard et al. 2009). Other distributions are implemented to accommodate the most common uncertainty representations: Normal (norm), Uniform (unif), and Loguniform (logu) (Table 2);
2. The uncertainty parameter F_0 . By default, it is the standard geometric deviation for the Lognormal distribution, but has no default value. It defines a standard “ 1σ ” confidence interval around the reference value k_0 (Sander et al. 20011), with the probability of assertion (Pr) defined by $\text{Pr}(k_0/F_0 \leq k \leq k_0 * F_0) \approx 0.68$. For other distributions than the Lognormal, F_0 has different meanings and units, as specified in Table 2;
3. The expansion parameter g (default = 0, units: Kelvin). It is used to parameterize a possible temperature-dependence of the uncertainty, according to the formula $F(T) = F_0 * \exp(g|1/T - 1/T_0|)$ (Sander et al. 20011). In KIDA, the reference temperature T_0 is fixed at 300 K (Hébrard et al. 2009).

More details on how to use, choose, and specify uncertainty representations in KIDA, notably for data providers, is given in <http://kida.obs.u-bordeaux1.fr/help/helpUncert.html>.

2.5. Member Area, Submitting Data to KIDA, and Downloading Lists of Reactions

In order to be able to interact effectively with KIDA, users need to first register by providing an email address and a password. They will then have access to a member area where they can submit rate coefficients to KIDA experts and see the history of their downloads. To improve the updating of KIDA, we have designed an online interface to upload data. Any user registered with KIDA can submit rate coefficients (with a temperature range of validity, an uncertainty and a reference). These data will then be sent to the “Super-Expert” of the type of reaction, who will then consult his/her experts to see if the data have

been correctly included or not. Lists of KIDA super-experts and experts with terms until 2013 are given in <http://kida.obs.u-bordeaux1.fr/contact>.

It is also possible to download lists of reactions from KIDA using an online form¹¹. In this form, the user can select 1) the elements and/or species he is interested in, 2) the types of reactions and 3) the range of temperature. This last choice can be used, for instance, as a filter to remove some of the high temperature reactions if one is interested in modeling the cold interstellar medium. If no recommended rate coefficients exist in the full range of temperature asked by the user, several rate coefficients for the same reaction may be available in the downloaded file. These reactions are included in the readme file downloaded at the same time, and the user has to deal with them. If rate coefficients are defined over complementary ranges of temperature, the user may want to keep them and take into account this information in the model (see section 3.1). Temperature ranges are indicated in the network along with the status of the KIDA recommendation (see section 2.3). In any case, we do not recommend the extrapolation of rate coefficients outside this range. If this procedure must be undertaken, we recommend instead using the results at the relevant extremum of temperature or consulting an expert or super expert from the KIDA lists.

For each reaction, KIDA stores up to three reactants, up to five products, the values of the kinetic parameters A, B and C, the uncertainty parameters F_0 , g and the type of uncertainty, the temperatures T_{min} and T_{max} , which define the range of temperatures over which the rate coefficient is defined, the name of the formula used to compute the rate coefficient or the external UV field used to compute the photoinduced rates, the number of the reaction in the network, another number that represents a particular choice of A, B and C when more than one possibility has been included, and a numerical recommendation given by KIDA experts and discussed in the next section. When there is no relevant temperature range, -9999 and 9999 default values are used for the minimum and maximum temperatures.

¹¹<http://kida.obs.u-bordeaux1.fr/astrophysicist>

3. New Model for the Cold Dense Interstellar Medium

3.1. The Nahoon Public Code

The numerical code Nahoon is publicly available on KIDA to compute the gas-phase chemistry for astrophysical objects (<http://kida.obs.u-bordeaux1.fr/models/>). The program is written in fortran 90 and uses the DL-SODES (double precision) solver from the ODEPACK package (<http://www.netlib.org/odepack/opkd-sum>) to solve the coupled stiff differential equations. The solver computes the chemical evolution of gas-phase species at a fixed temperature and density (commonly referred to as zero-dimension, or 0D) but can be used in 1D if a grid of temperature, density, and visual extinction is provided. Nahoon is formatted to read chemical networks downloaded from KIDA. It contains a test to check the temperature range of the validity of the rate coefficients and avoid extrapolations outside this range. A test is also included to check for duplication of chemical reactions, defined over complementary ranges of temperature. In that case, Nahoon chooses the recommended rate coefficient. Some reactions, although important for the interstellar medium, are not included in KIDA such as H₂ formation on grains or recombination of cations with negatively charged grains. These reactions are, however, provided in a set of user instructions associated with Nahoon. This code represents a new version of the one that was put online in 2007 (Wakelam & Herbst 2008). Grains, both neutral and negatively charged, and electrons are considered as chemical species and their concentrations are computed at the same time as those of the other species.

3.2. Gas-phase network for cold dense molecular clouds

KIDA is a non-selective database. It may not be easy for all users to select the most appropriate list of reactions. For this reason, we will release every year a standard list of reactions for the dense interstellar medium between 10 and 300 K. This network will be available on the same page as Nahoon (<http://kida.obs.u-bordeaux1.fr/models/>). To treat the formation of molecular hydrogen on granular surfaces, the OSU database uses the approximation from Hollenbach & Salpeter (1971), which neglects any details of the process, and contains the assumption that the formation rate of H₂ is equal to 1/2 the collision rate of H atoms with grains (see also Leung et al.

1984). This approximation can induce some numerical problems during the computation because the elemental abundance of hydrogen is not rigorously conserved by the rather complex procedure used by the authors of the code. Rather than improving the procedure, we use a simplified version of the more detailed approach in gas-grain models (Hasegawa et al. 1992). In particular, we have divided the H₂ formation into two reactions and included another species JH, which refers to hydrogen atoms on grains. The first reaction is the sticking of atomic hydrogen onto grains (neutral or negatively charged): H → JH.

The rate of this reaction (in cgs units of cm⁻³ s⁻¹) is

$$R_{H,acc} = \pi a^2 \sqrt{\frac{8kT}{\pi m_H}} n_d n(H) \quad (8)$$

with a the radius of grains (in cm), k the Boltzmann constant, m_H the atomic mass of hydrogen, n_d the total density of grains, $T(K)$ the temperature, and $n(H)$ the density of atomic hydrogen. For 0.1 μm radius grains, the rate is $7.92 \times 10^{-5} \sqrt{T/300} n_d n(H)$.

The second reaction is the formation of H₂ on grains: JH + JH → H₂. We assume here that H₂ formed on grains is directly released in the gas phase. Since desorption of JH is not included, the process has an efficiency of 100%. The rate of this reaction is given by the expression

$$R_{H_2,form} = \frac{\nu_0 \exp(-E_b/kT)}{N_s n_d} n(JH)^2 \quad (9)$$

with N_s the number of binding sites on a grain, ν_0 the characteristic or trial vibrational frequency, and E_b the potential energy barrier between two adjacent surface potential wells.

A useful approximate formula is $\nu_0 = \sqrt{2n_s E_D k_B / (\pi m_H)}$ (Tielens & Allamandola 1987), where n_s is the surface density of sites on a grain in cm⁻², and E_D the adsorption energy of H in erg. The reader can find a detailed explanation in Hasegawa et al. (1992) and Semenov et al. (2010). Assuming grains of 0.1 μm radius, the number of sites per grain of 10⁶ (Hasegawa et al. 1992), $E_D/k = 450$ K, and $E_b/k = 225$ K, n_s is 7.96×10^{14} cm⁻², ν_0 is 4.32×10^{12} s⁻¹ and the rate of H₂ formation is $8.64 \times 10^6 \exp(-225/T) n(JH)^2 / n_d$ cm³ s⁻¹. See Garrod & Herbst (2006) for a discussion concerning the values of E_D and E_b , which depend upon the surface. The value of n_d should obey the constraint that $\pi a^2 n_d \approx 1 - 2 \times 10^{-21} n_H$, so that here $n_d \approx 5 \times 10^{-12} n_H$, where

a is the granular radius and n_{H} is the total hydrogen nuclear density.

The second class of processes involves electron attachment to grains and the recombination of cations with negatively charged grains. Rate coefficients for such processes can be obtained in the ballistic approximation from the following equation:

$$k_{\text{coll}} = \pi a^2 \sqrt{\frac{8kT}{\pi m_i}} \quad (10)$$

where m_i is the mass (g) of the impacting species (electrons or ions). We make the approximation that the mass of the impactor is negligible compared to the grain mass, so that the reduced mass is then approximately the mass of the lighter collider. For collisions between two charged species (cations and negatively charged grains), this rate can be multiplied by $(1 + \frac{Ze^2}{akT})$ to account for the Coulomb attraction, with e the elemental electronic charge and Z the number of negative charges on the grain (Weisheit & Upham 1978). It is typically assumed, however, that, in dense clouds, grains can be neutral or singly negatively charged only, reflecting the possibility that the trajectory of an electron nearing a negatively-charged grain is repulsive. Following Leung et al. (1984), we neglect any surface electronic charge, based on the assumption that the negative charge/charges on a grain remain localized so the additional factor does not play an important role in the dynamics of the recombination with cations. If the negative charges on a grain were delocalized, then the Coulomb interaction should be included. We have listed in Table 3 the minimum set of reactions with grains that should be included in a network for dense clouds. The rate coefficients ($\text{cm}^3 \text{s}^{-1}$) for these processes are computed from the A and B listed in the table using the equation $k = A(T/300)^B$, where A is obtained for a grain radius of $0.1 \mu\text{m}$ and $B = 0.5$. For grains of different radii, the ballistic rate coefficients simply scale as the square of the radius. The referee has noted correctly that reactions involving gaseous ions and neutral grains are not included in the KIDA network. They are not included because the only reactive gas-grain processes in the network involve charge neutralization of negatively-charged grains.

In the interstellar medium, molecules can be dissociated by absorption of UV photons via electronic continua or through lines. The photodissociation of the dominant species H_2 and CO is dominated by line absorption (although the mechanisms are different) so that H_2 and CO self-shielding is an important process.

To avoid a detailed treatment of radiative transfer, we adopted the approximation from Lee et al. (1996), which gives H_2 and CO photodissociation rates as a function of A_{v} , as well as the H_2 and CO column densities. To include self-shielding, the user of Nahoon has to set the conditions at the border of the cloud, including the abundances of H_2 and CO even when working in 0D. In 1D, a plane parallel geometry is assumed and the UV photons enter the cloud from one direction. Photodissociation rates at a spatial point are then computed using the column densities of H_2 and CO according to the computed abundances and physical structure in front of this point. The visual extinction is then computed using the H_2 column density and the standard ratio between this column density and visual extinction. This geometry is not correct for clouds where UV photons can penetrate from both sides of the cloud. Note that new shielding functions have been published by Visser et al. (2009) and will be included in Nahoon in the future.

In addition to these processes, which are downloaded with the new Nahoon, we have downloaded our first annual network of gas-phase chemical reactions necessary to compute the chemistry in the interstellar medium where the temperature is between 10 and 300 K and where surface chemistry is not critical. Approximations for the depth-dependent photodissociation rates are valid for A_{v} above 5 (van Dishoeck 1988). The full network, designated *kida.uva.2011*, and the associated list of species are available at <http://kida.obs.u-bordeaux1.fr/models>. This list contains 6090 reactions involving 474 species, including neutral and negatively charged grains, as well as electrons. Elements that are included are H, He, C, N, O, Si, S, Fe, Na, Mg, Cl, P and F. Anionic chemistry from Harada & Herbst (2008) has also been included. For temperatures in the range 300 K - 800 K, modelers can use the high-temperature OSU network *osu-09-2010-ht*, found and discussed on the URL <http://www.physics.ohio-state.edu/~eric/research.html>.

3.3. Using uncertainties for rate coefficients in models

Using uncertainty information in chemical modeling is vital, notably for the extreme environments targeted by KIDA, where many reaction rate coefficients are known with little accuracy, either estimated or extrapolated. Uncertainty management has two goals:

1. Uncertainty Propagation: to estimate the preci-

sion of the model outputs; and

2. Sensitivity Analysis: to identify key reactions, i.e. those contributing notably to the uncertainty of model outputs and for which better experimental or theoretical estimations are needed (Dobrijevic et al. 2010).

Improvement of data in KIDA is based on an iterative strategy involving uncertainty propagation and sensitivity analysis (Wakelam et al. 2010b).

Monte Carlo Uncertainty Propagation (MCUP).

The simplest way to implement these methods is through Monte Carlo sampling (Thompson & Stewart 1991; Dobrijevic et al. 2008; Wakelam et al. 2005; Carrasco et al. 2007; Wakelam et al. 2010b). Random draws for a rate coefficient at any temperature are generated using the formulae in Table 4. For MCUP, the code is run for N random draws of the M rate constants of the chemical scheme $\{k_i^{(j)}; i = 1, M; j = 1, N\}$ (all parameters are perturbed simultaneously), and the N sets of outputs are stored for further statistical analysis (mean value, uncertainty factor, input/output correlation...).

Sensitivity Analysis. A convenient way to perform sensitivity analysis in the MCUP setup is to calculate the correlation coefficients between the inputs (k_i) and outputs of the model. Large absolute values of correlation coefficients reveal strong influences of inputs on outputs (Wakelam et al. 2010b; Dobrijevic et al. 2010). Variations of this method include using rank correlation coefficients, or the logarithm of inputs and/or outputs (Helton et al. 2006; Saltelli et al. 2004).

Monte Carlo simulations have been implemented in the Nahoon code bundle, and an IDL procedure has been provided to compute the standard deviation for the predicted abundances of species due to uncertainties in reaction rate coefficients.

3.4. Limitations

The interstellar chemical model defined by both the numerical program Nahoon and the osu.uva.2011 network takes into account a limited number of processes, which makes its use valid only over a limited range of temperature, density, and visual extinction. This model is meant to compute the gas-phase chemistry in regions where the visual extinction is larger than $\approx 3 - 4$. For smaller A_V , detailed PDR codes,

such as the Meudon code (Le Petit et al. 2009), are more appropriate. Also, this model does not take into account positively charged grains, which play a major role at low extinction. The temperature limit is set by the network to the range 10 to 300 K, as already discussed. For higher temperatures, one should adapt an expanded gas-phase network developed by Harada et al. (2010), which can be utilized up to gas temperatures of 800 K. The accretion time scale onto grains of radius 0.1μ is roughly $10^9/n_H(\text{cm}^{-3})$ yr. At times larger than this, the gas-phase can be significantly depleted of heavy species if the temperature lies below the the evaporation temperature of easily vaporized species such as CO and CH₄. At lower temperatures, non-thermal desorption mechanisms, such as photodesorption, can still maintain a residual gas phase. Models including interactions with grains and grain surface processes (Garrod et al. 2007) are, however, based on many uncertain processes. In addition their use and the analysis of the results are more difficult. For this reason, pure gas-phase models can still be useful even for dense cloud conditions but users have to be careful when they compare their results with observations.

The computation of H₂ formation on grains and the recombination of negative grains with cations, presented here, depend on the size of the grains. We have computed the rate coefficients for these processes for a single grain size of $0.1\mu\text{m}$, which with a typical grain density of 2.5 gm/cm^3 , can explain observed extinctions with a gas-to-dust mass ratio of about 0.01, a reasonable value considering heavy elemental depletions in diffuse clouds (Tielens 2005). The effects of using a detailed grain size distribution on the rates of surface processes and on the chemical results for a full gas-grain model can be found in Acharyya et al. (2011); these authors included grains large enough to have a constant rather than stochastic temperature and used a variety of size distributions (Weingartner & Draine 2001; Mathis et al. 1977). They found little effect on the overall chemical results for a cold core with the OSU gas-grain network. The relationship between granular size and rates of assorted processes is a complex one, because the rates can depend upon a number of parameters in addition to surface area, such as whether the system lies in the accretion or diffusion limit and, for cold dense cores, the depth of the ice mantle.

The temperature dependence of the rate of H₂ formation on grains is a rather complex issue, and

depends strongly on how rough the dust surface is and the availability of chemisorption sites, both of which can allow the process to continue up to much higher temperatures than the standardly-assumed Langmuir-Hinshelwood mechanism based on physisorption on smooth surfaces (Cuppen & Herbst 2005; Cazaux & Tielens 2010). The net result is that for systems with both physisorption and chemisorption sites, the efficiency of H₂ formation can remain high at all relevant interstellar dust temperatures, although this result is not guaranteed. For small particles, in which the temperature is stochastic, the situation is more complex but has also been studied via Monte Carlo techniques (Cuppen et al. 2006). For dust particles with well-defined temperatures up to 50 K, the physisorption mechanism assumed here should be adequate for rough particles.

Finally, despite their possible impact on the interstellar chemistry of cold dense cores, as shown by Wakelam & Herbst (2008), Polycyclic Aromatic Hydrocarbons (PAHs) are not included in this model because of the strong uncertainties that still remain (1) on the size and abundance of these species in dense regions and (2) on their reactions with other species.

3.5. Comparison with other networks

To compare the abundances computed with the `kida.uva.2011` network and other public networks, we have considered the case of dark molecular clouds using Nahoon in 0D. The temperature is set to 10 K, the proton density to $2 \times 10^4 \text{ cm}^{-3}$, the total H₂ cosmic ray ionization to $1.3 \times 10^{-17} \text{ s}^{-1}$ and the visual extinction to 30. Initial abundances are taken from Wakelam et al. (2010a). The results for selected molecular abundances, as computed with the `kida.uva.2011` list of reactions, are displayed in Fig. 1 with the results of the uncertainty propagation of rate coefficients. For comparison, abundances computed using the `udfa06` network (<http://www.udfa.net/>) and `osu-01-2009` network (<http://www.physics.ohio-state.edu/~eric/research.html>) networks are also included. Contrary to the publicly available `udfa06` network, the OSU network used also contains anions.

The particular molecular abundances plotted have been selected because they show some variation in their abundances with the database used. The OSU and `kida.uva` networks lead to similar results since KIDA is mainly based on the OSU compilation although a number of updates have been performed. Abundances

computed with `udfa06` on the contrary can be different, as shown especially for CS and the cations in Fig. 1. Model predictions done with the different networks are however within the 2σ error bars computed from the uncertainty propagation of rate coefficients, which are typically an order-of-magnitude in both directions.

4. Conclusions

We have presented in this article a new chemical database for gas-phase astrochemistry named KIDA (for KInetic Database for Astrochemistry). KIDA can become a useful tool for physical chemists to circulate their relevant work for astrochemistry, especially their most up-to-date data. In addition to kinetic data for the chemistry in the interstellar medium, KIDA will soon contain kinetic data for planetary atmospheres (other than the Earth). Kinetic data for carbon-rich and oxygen-rich circumstellar shells are also planned for the future. State-to-state reactions, excited species, isotopologues and ortho and para forms will be considered for addition in KIDA. The quality of the data present in KIDA will be improved over the years by KIDA experts and will be preserved thanks to the detailed information stored in KIDA. The visibility of KIDA will be improved by its interoperability with the Virtual Atomic and Molecular DataCenter (Dubernet et al. 2010, <http://www.vamdc.eu/>) and the Europlanet Research Infrastructure (<http://www.europlanet-eu.org/outreach/>). In addition to the rate coefficients, we are providing the community with a public fortran code, able to read networks downloaded from KIDA, called Nahoon, which can calculate the evolution of the gas-phase composition for 0D and 1D sources in the ISM and an extensive list of chemical reactions for the dense ISM with temperatures between 10 and 300 K. Neither KIDA nor Nahoon treats the interaction of the gas-phase with interstellar grains, except for the formation of molecular hydrogen and recombination of the major cations with negatively charged grains. Omission of these processes is not realistic in the context of interstellar chemistry but the numerical and chemical complexity, and additional uncertainties brought by the treatment of surface chemistry, make pure gas-phase models still needed. In addition, gas-grain models also rely on the quality of gas-phase rate coefficients.

The KIDA team acknowledges various sources of funding, which have allowed us to develop this

database. These sources include The University of Bordeaux, The “Institut de Physique Fondamentale de Bordeaux”, the CNRS/INSU (PCMI, PNP and ASOV), The Observatoire Aquitain des Sciences de l’Univers. We acknowledge support from the European Research Council (ERC Grant 209622: E₃ARTHs), the Agence Nationale de la Recherche (ANR-JC08_311018: EMA:INC), the European program Astronet (CATS project), and the European Seventh Framework Programme (FP7: VAMDC and EUROPLANET projects). VAMDC is funded under the Combination of Collaborative Projects and Coordination and Support Actions Funding Scheme of The Seventh Framework Program (Call topic: INFRA-2008-1.2.2 , Grant Agreement number: 239108). The EUROPLANET RI (Research Infrastructure) is funded under the Grant Agreement number: 228319. The International Space Science Institute provided some of the KIDA team members with the opportunity to start this project in the context of the international team with the title “New generation of databases for interstellar chemical modeling in preparation for HSO and ALMA”. VW and IWMS thank the Royal Society for its financial support of their collaboration. VW is grateful to all the participants of the project, who helped design the data model of the database. VW also thanks Franck Le Petit for helpful discussions on the rate coefficients for photodissociation induced by cosmic rays. EH thanks the National Science Foundation (US) for his program in astrochemistry and NASA for the study of pre-planetary matter. NGA thanks NASA for support of his experimental program. SJK acknowledges support through NASA - PATM grant number NNH09AK24I. The anonymous referee is thanked for his/her useful comments, which helped to improve the quality of the paper.

A. Appendix

A.1. KIDA references

Original references for kinetic data in KIDA are: Adams & Smith (1976, 1984); Amano (1990); Anicich & Huntress (1986); Anicich et al. (1976); Antipov et al. (2009); Atkinson et al. (2004); Chenel et al. (2010); Barckholtz et al. (2001); Barlow & Dunn (1987); Baulch et al. (2005); Becker et al. (1989, 2000); Bedjanian et al. (1999); Bergeat et al. (2009); Berkowitz et al. (1986); Berteloite et al. (2010b,a); Bettens et al. (1999); Bettens & Herbst (1995); Bohme et al. (1988); Brown et al. (1983); Brownsword et al. (1997, 1996); Canosa et al. (1997); Carty et al. (2006); Chabot et al. (2010); Costes et al. (2009); Daugey et al. (2005); Davidson & Hanson (1990); Deeyamulla & Husain (2006); Defrees & McLean (1989); Derkatch et al. (1999); Dorthe et al. (1991); Duff & Sharma (1996); Ehlerding et al. (2004); Eichelberger et al. (2007); Fehsenfeld et al. (1975); Gamallo et al. (2006); Geppert et al. (2004b,a, 2000); Grebe & Homann (1982); Gredel et al. (1989); Gu et al. (2006); Sun & Li (2004); Harada & Herbst (2008); Herbst et al. (1989a); Herbst (1985, 1987); Herbst et al. (1989b, 2000); Jensen et al. (2000); Kaiser et al. (2002); Kalhori et al. (2002); Lee et al. (1978); Lilenfeld & Richardson (1977); Lique et al. (2009); Loison & Bergeat (2009); Marquette et al. (1988); Martinet et al. (2004); Martinez et al. (2008, 2010); Mayer et al. (1967); McGowan et al. (1979); Mebel & Kaiser (2002); Messing et al. (1980); Millar & Herbst (1990); Montaigne et al. (2005); Neufeld et al. (2005); Ozeki et al. (2004); Parisel et al. (1996); Prasad & Huntress (1980); Huo et al. (2011); Sablier & Fujii (2002); Sander et al. (2001); Shin & Beauchamp (1986); Sims et al. (1994); Smith & Adams (1981); Smith (1989); Smith et al. (2001); Snow et al. (2009); Stevens et al. (2010); Talbi & Smith (2009); Thorne et al. (1983, 1984); Tuna et al. (2008); van Dishoeck et al. (2006); van Dishoeck (1988); Van Harrevelt et al. (2002); van Hemert & van Dishoeck (2008); Vazquez et al. (2009); Viggiano et al. (2005); Vinckier (1979); Vroom & de Heer (1969); Wakelam et al. (2009); Wennberg et al. (1994); Woon & Herbst (2009); Zabarnick et al. (1989).

REFERENCES

- Acharyya, K., Hassel, G. E., & Herbst, E. 2011, *ApJ*, 732, 73
- Adams, N. G. & Smith, D. 1976, *Journal of Physics B Atomic Molecular Physics*, 9
- . 1984, *Int. J. Mass Spectrom. Ion Proc.*, 61, 133
- Amano, T. 1990, *Journal of Chemical Physics*, 92, 6492
- Anicich, V. G. 1993, *Journal of Physical and Chemical Reference Data*, 22, 1469
- Anicich, V. G., Huntress, W. T., & Futrell, J. H. 1976, *Chemical Physics Letters*, 40, 233
- Anicich, V. G. & Huntress, Jr., W. T. 1986, *ApJ*, 62, 553
- Antipov, S. V., Sjölander, T., Nyman, G., & Gustafsson, M. 2009, *J. Chem. Phys.*, 131, 074302
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., & Troe, J. 2004, *Atmospheric Chemistry & Physics*, 4, 1461
- Barckholtz, C., Snow, T. P., & Bierbaum, V. M. 2001, *ApJ*, 547, L171
- Barlow, S. E. & Dunn, G. H. 1987, *Int. J. Mass Spectrom. Ion Proc.*, 80, 227
- Baulch, D. L., Bowman, C. T., Cobos, C. J., Cox, R. A., Just, T., Kerr, J. A., Pilling, M. J., Stocker, D., Troe, J., Tsang, W., Walker, R. W., & Warnatz, J. 2005, *Journal of Physical Chemistry Reference Data*, 34, 757
- Becker, K. H., Engelhardt, B., Wiesena, P., & Bayesb, K. D. 1989, *Chemical Physics Letters*, 154, 342
- Becker, K. H., Kurtenbach, R., Schmidt, F., & Wiesen, P. 2000, *Combust. Flame*, 120, 570
- Bedjanian, Y., Bras, G. L., & Poulet, G. 1999, *Journal of Chemical Physics*, 110, 7017
- Bergeat, A., Hickson, K. M., Daugey, N., Caubet, P., & Costes, M. 2009, *Phys. Chem. Chem. Phys.*, 11, 8149
- Berkowitz, J., Curtiss, L. A., Gibson, S. T., Greene, J. P., Hillhouse, G. L., & Pople, J. A. 1986, *J. Chem. Phys.*, 84, 375
- Berteloite, C., Le Picard, S. D., Balucani, N., Canosa, A., & Sims, I. R. 2010a, *Phys. Chem. Chem. Phys.*, 12, 3666
- . 2010b, *Phys. Chem. Chem. Phys.*, 12, 3677
- Bettens, R. P. A., Hansen, T. A., & Collins, M. A. 1999, *J. Chem. Phys.*, 111, 6322
- Bettens, R. P. A. & Herbst, E. 1995, *Int. J. Mass Spectrom. Ion Proc.*, 149, 321
- Bohme, D. K., Wlodek, S., & Fox, A. 1988, in *Rate Coefficients in Astrochemistry. Proceedings of a Conference held in UMIST, Manchester, United Kingdom, September 21-24, 1987.*, ed. T. J. Millar & D. A. Williams, p. 193
- Brown, R. D., Eastwood, F. W., Elmes, P. S., & Godfrey, P. D. 1983, *Journal of the American Chemical Society*, 105, 6496
- Brownsword, R. A., Gatenby, S. D., Herbert, L. B., Smith, I. W. M., Stewart, D. W. A., & Symonds, A. C. 1996, *J. Chem. Soc. Faraday Trans.*, 92, 723
- Brownsword, R. A., Hancock, G., & Heard, D. E. 1997, *J. Chem. Soc. Faraday Trans.*, 93, 2473
- Canosa, A., Sims, I. R., Travers, D., Smith, I. W. M., & Rowe, B. R. 1997, *A&A*, 323, 644
- Carrasco, N., Dutuit, O., Thissen, R., Banaszkiwicz, M., & Pernot, P. 2007, *Planet. Space Sci.*, 55, 141
- Carty, D., Goddard, A., Kahler, S. P. K., Sims, I. R., & Smith, I. W. M. 2006, *J. Phys. Chem. A*, 110, 3101
- Cazaux, S. & Tielens, A. G. G. M. 2010, *ApJ*, 715, 698
- Chabot, M., Tuna, T., Béroff, K., Pino, T., Le Padellec, A., Déséquelles, P., Martinet, G., Nguyen-Thi, V. O., Carpentier, Y., Le Petit, F., Roueff, E., & Wakelam, V. 2010, *A&A*, 524, 39
- Chenel, A., Mangaud, E., Justum, Y., Talbi, D., Bacchus-Montabonel, M.-C., & Desouter-Lecomte, M. 2010, *Journal of Physics B Atomic Molecular Physics*, 43, 245701
- Costes, M., Halvick, P., Hickson, K. M., Daugey, N., & Naulin, C. 2009, *ApJ*, 703, 1179

- Cravens, T. E. & Dalgarno, A. 1978, *ApJ*, 219, 750
- Cuppen, H. M. & Herbst, E. 2005, *MNRAS*, 361, 565
- Cuppen, H. M., Morata, O., & Herbst, E. 2006, *MNRAS*, 367, 1757
- Dalgarno, A., Yan, M., & Liu, W. 1999, *ApJS*, 125, 237
- Daugey, N., Caubet, P., Retail, B., Costes, M., Bergeat, A., & Dorthe, G. 2005, *Phys. Chem. Chem. Phys.*, 7, 2921
- Davidson, D. F. & Hanson, R. K. 1990, *International Journal of Chemical Kinetics*, 22, 843
- Deeyamulla, M. P. & Husain, D. 2006, *J. Photochem. Photobiol. A Chem.*, 184, 347
- Defrees, D. J. & McLean, A. D. 1989, *Chemical Physics Letters*, 158, 541
- Derkatch, A. M., Al-Khalili, A., Vikor, L., Neau, A., Shi, W., Danared, H., Af Ugglas, M. A., & Larsson, M. 1999, *Journal of Physics B Atomic Molecular Physics*, 32, 3391
- Dobrijevic, M., Cavalié, T., Hébrard, E., Billebaud, F., Hersant, F., & Selsis, F. 2010, *Planet. Space Sci.*, 58, 1555
- Dobrijevic, M., Claeys-Bruno, M., Sergent, M., & Phan-Tan-Luu, R. 2008, *Planet. Space Sci.*, 56, 519
- Dorthe, G., Caubet, P., Vias, T., Barrere, B., & Marchais, J. 1991, *J. Phys. Chem.*, 95, 5109
- Draine, B. T. 1978, *ApJ*, 36, 595
- Dubernet, M. L., Boudon, V., Culhane, J. L., Dimitrijevic, M. S., Fazliev, A. Z., Joblin, C., Kupka, F., Leto, G., Le Sidaner, P., Loboda, P. A., Mason, H. E., Mason, N. J., Mendoza, C., Mulas, G., Millar, T. J., Nuñez, L. A., Perevalov, V. I., Piskunov, N., Ralchenko, Y., Rixon, G., Rothman, L. S., Roueff, E., Ryabchikova, T. A., Ryabtsev, A., Sahal-Bréchet, S., Schmitt, B., Schlemmer, S., Tennyson, J., Tyuterev, V. G., Walton, N. A., Wakelam, V., & Zeippen, C. J. 2010, *J. Quant. Spectrosc. Radiat. Transfer*, 111, 2151
- Duff, J. W. & Sharma, R. D. 1996, *Geophysical Research Letters*, 23, 2777
- Ehlerding, A., Hellberg, F., Thomas, R., Kalhori, S., Viggiano, A. A., Arnold, S. T., Larsson, M., & Af Ugglas, M. 2004, *Phys. Chem. Chem. Phys.*, 6, 949
- Eichelberger, B., Snow, T. P., Barckholtz, C., & Bierbaum, V. M. 2007, *ApJ*, 667, 1283
- Fehsenfeld, F. C., Lindinger, W., Schmeltekopf, A. L., Albritton, D. L., & Ferguson, E. E. 1975, *J. Chem. Phys.*, 62, 2001
- Gamallo, P., Sayos, R., Gonzalez, M., Petrongolo, C., & Defazio, P. 2006, *J. Chem. Phys.*, 124, 174303
- Garrod, R. T. & Herbst, E. 2006, *A&A*, 457, 927
- Garrod, R. T., Wakelam, V., & Herbst, E. 2007, *A&A*, 467, 1103
- Geppert, W. D., Reignier, D., Stoecklin, T., Naulin, C., Costes, M., Chastaing, D., Le Picard, S. D., Sims, I. R., & Smith, I. W. M. 2000, *Phys. Chem. Chem. Phys.*, 2, 2873
- Geppert, W. D., Thomas, R., Ehlerding, A., Hellberg, F., Österdahl, F., Af Ugglas, M., & Larsson, M. 2004a, *Inter. J. Mass Spec.*, 237, 25
- Geppert, W. D., Thomas, R., Semaniak, J., Ehlerding, A., Millar, T. J., Österdahl, F., Af Ugglas, M., N., D., Paal, A., & Larsson, M. 2004b, *ApJ*, 609, 459
- Grebe, J. & Homann, K. 1982, *Ber. Bunsenges. Phys. Chem.*, 86, 587
- Gredel, R., Lepp, S., & Dalgarno, A. 1987, *ApJ*, 323, L137
- Gredel, R., Lepp, S., Dalgarno, A., & Herbst, E. 1989, *ApJ*, 347, 289
- Gu, X., Guo, Y., Mebel, A. M., & Kaiser, R. I. 2006, *J. Phys. Chem. A*, 110, 11265
- Harada, N. & Herbst, E. 2008, *ApJ*, 685, 272
- Harada, N., Herbst, E., & Wakelam, V. 2010, *ApJ*, 721, 1570
- Hasegawa, T. I., Herbst, E., & Leung, C. M. 1992, *ApJ*, 82, 167
- Hébrard, E., Dobrijevic, M., Pernot, P., Carrasco, N., Bergeat, A., Hickson, K. M., Canosa, A., Le Picard, S. D., & Sims, I. R. 2009, *Journal of Physical Chemistry A*, 1131, 11227

- Helton, J., Johnson, J., Sallaberry, C., & Storlie, C. 2006, *Reliability Engineering and System Safety*, 91, 1175
- Herbst, E. 1985, *ApJ*, 291, 226
- . 1987, *ApJ*, 313, 867
- Herbst, E., Millar, T. J., Wlodek, S., & Bohme, D. K. 1989a, *A&A*, 222, 205
- Herbst, E., Smith, D., Adams, N. G., & McIntosh, B. J. 1989b, *J. Chem. Soc., Faraday Trans. 2*, 85, 1655
- Herbst, E., Terzieva, R., & Talbi, D. 2000, *MNRAS*, 311, 869
- Hollenbach, D. & Salpeter, E. E. 1971, *ApJ*, 163, 155
- Huo, R.-P., Zhang, X., Huang, X.-R., Li, J.-L., & Sun, C.-C. 2011, *J. Phys. Chem. A*, 115, 3576
- Jensen, M. J., Bilodeau, R. C., Safvan, C. P., Seiersen, K., Andersen, L. H., Pedersen, H. B., & Heber, O. 2000, *ApJ*, 543, 764
- Kaiser, R. I., Yamada, M., & Osamura, Y. 2002, *J. Phys. Chem. A*, 106, 4825
- Kalhari, S., Viggiano, A. A., Arnold, S. T., Rosèn, S., Semaniak, J., Derkatch, A. M., Af Ugglas, M., & Larsson, M. 2002, *A&A*, 391, 1159
- Le Petit, F., Barzel, B., Biham, O., Roueff, E., & Le Bourlot, J. 2009, *A&A*, 505, 1153
- Le Petit, F., Nehmé, C., Le Bourlot, J., & Roueff, E. 2006, *ApJ*, 164, 506
- Le Teuff, Y. H., Millar, T. J., & Markwick, A. J. 2000, *A&A*, 146, 157
- Lee, H.-H., Herbst, E., Pineau des Forets, G., Roueff, E., & Le Bourlot, J. 1996, *A&A*, 311, 690
- Lee, J. H., Michael, J. V., Payne, W. A., & Stief, L. J. 1978, *J. Chem. Phys.*, 69, 3069
- Leung, C. M., Herbst, E., & Huebner, W. F. 1984, *ApJ*, 56, 231
- Lilenfeld, H. V. & Richardson, R. J. 1977, *J. Chem. Phys.*, 67, 3991
- Lique, F., Jorfi, M., Honvault, P., Halvick, P., Lin, S. Y., Guo, H., Xie, D. Q., Dagdigian, P. J., Klos, J., & Alexander, M. H. 2009, *J. Chem. Phys.*, 79, 221104
- Loison, J. C. & Bergeat, A. 2009, *Phys. Chem. Chem. Phys.*, 11, 655
- Marquette, J. B., Rebrion, C., & Rowe, B. R. 1988, *J. Chem. Phys.*, 89, 2041
- Martinet, G., Diaz-Tendero, S., Chabot, M., Wohrer, K., Della-Negra, S., Mezdari, F., Hamrita, H., Desesquelles, P., Le Padellec, A., Gardes, D., Lavergne, L., Lalu, G., Grave, X., Clavelin, J. F., Hervieux, P. A., Alcami, M., & Martin, F. 2004, *Physical Review Letters*, 93, 063401
- Martinez, Jr., O., Betts, N. B., Villano, S. M., Eyet, N., Snow, T. P., & Bierbaum, V. M. 2008, *ApJ*, 686, 1486
- Martinez, Jr., O., Yang, Z., Demarais, N. J., Snow, T. P., & Bierbaum, V. M. 2010, *ApJ*, 720, 173
- Mathis, J. S., Rimpl, W., & Nordsieck, K. H. 1977, *ApJ*, 217, 425
- Mayer, S. W., Schieler, L., & Johnston, H. S. 1967, *Symp. Int. Combust. Proc.*, 11, 837
- McGowan, J. W., Mul, P. M., D'Angelo, V. S., Mitchell, J. B. A., Defrance, P., & Froelich, H. R. 1979, *Physical Review Letters*, 42, 373
- Mebel, A. M. & Kaiser, R. I. 2002, *ApJ*, 564, 787
- Messing, I., Carrington, T., Filseth, S., & C.M., S. 1980, *Chemical Physics Letters*, 74, 56
- Millar, T. J., Bennett, A., Rawlings, J. M. C., Brown, P. D., & Charnley, S. B. 1991, *A&A*, 87, 585
- Millar, T. J., Farquhar, P. R. A., & Willacy, K. 1997, *A&A*, 121, 139
- Millar, T. J. & Herbst, E. 1990, *A&A*, 231, 466
- Montaigne, H., Geppert, W. D., Semaniak, J., Österdahl, F., Hellberg, F., Thomas, R. D., Af Ugglas, M., Roberts, H., Millar, T. J., Zhaunerchyk, V., Kamińska, M., Al-Khalili, A., Källberg, A., & Larsson, M. 2005, *ApJ*, 631, 653
- Neufeld, D. A., Wolfire, M. G., & Schilke, P. 2005, *ApJ*, 628, 260
- Ozeki, H., Hirao, T., Saito, S., & Yamamoto, S. 2004, *ApJ*, 617, 680
- Parisel, O., Hanus, M., & Ellinger, Y. 1996, *Chemical Physics*, 212, 331

- Pineau des Forets, G., Roueff, E., Schilke, P., & Flower, D. R. 1993, *MNRAS*, 262, 915
- Prasad, S. S. & Huntress, Jr., W. T. 1980, *ApJ*, 43, 1
- Prasad, S. S. & Tarafdar, S. P. 1983, *ApJ*, 267, 603
- Rimmer, P. B., Herbst, E., Morata, O., & Roueff, E. 2012, *A&A*, 537, A7
- Sablier, M. & Fujii, T. 2002, *Chemical Reviews*, 102, 2855
- Saltelli, A., Tarantola, S., Campolongo, F., & Ratto, M. 2004, *Sensitivity Analysis in Practice* (Wiley)
- Sander, S. P., Abbatt, J., Barker, J. R., Burkholder, J. B., Friedl, R. R., Golden, D. M., Huie, R. E., Kolb, C. E., Kurylo, M. J., Moortgat, G. K., Orkin, V. L., & Wine, P. H. 2001, *JPL Publication 10-6*, Jet Propulsion Laboratory, Pasadena
- Semenov, D., Hersant, F., Wakelam, V., Dutrey, A., Chapillon, E., Guilloteau, S., Henning, T., Launhardt, R., Piétu, V., & Schreyer, K. 2010, *A&A*, 522, A42+
- Shin, S. K. & Beauchamp, J. L. 1986, *J. Phys. Chem.*, 90, 1507
- Sims, I. R., Queffelec, J.-L., Defrance, A., Rebrion-Rowe, C., Travers, D., Bocherel, P., Rowe, B. R., & Smith, I. W. M. 1994, *J. Chem. Phys.*, 100, 4229
- Smith, I. W. M. 1989, *ApJ*, 347, 282
- Smith, I. W. M. & Adams, N. G. 1981, *MNRAS*, 197, 377
- Smith, T. C., Clouthier, D. J., & Steimle, T. C. 2001, *The Journal of Chemical Physics*, 115, 817
- Snow, T. P., Stepanovic, M., Betts, N. B., Eichelberger, B. R., Martinez, O., & Bierbaum, V. M. 2009, *Astrobiology*, 9, 1001
- Stevens, W. R., Ruscic, B., & Baer, T. 2010, *J. Phys. Chem. A*, 114, 1313413145
- Sun, H. & Li, Z. 2004, *Chemical Physics Letters*, 399, 33
- Talbi, D. & Smith, I. W. M. 2009, *Phys. Chem. Chem. Phys.*, Incorporating Faraday Transactions, 11, 8477
- Thompson, A. M. & Stewart, R. W. 1991, *J. Geophys. Res. - Atmos.*, 96, 13089
- Thorne, L. R., Anicich, V. G., & Huntress, W. T. 1983, *Chemical Physics Letters*, 98, 162
- Thorne, L. R., Anicich, V. G., Prasad, S. S., & Huntress, Jr., W. T. 1984, *ApJ*, 280, 139
- Tielens, A. G. G. M. 2005, *The Physics and Chemistry of the Interstellar Medium* (Cambridge, UK: Cambridge University Press)
- Tielens, A. G. G. M. & Allamandola, L. J. 1987, in *NATO ASIC Proc. 210: Physical Processes in Interstellar Clouds*, ed. G. E. Morfill & M. Scholer, 333–376
- Troe, J. 1996, *J. Chem. Phys.*, 105, 6249
- Tuna, T., Chabot, M., Pino, T., Désesquelles, P., LePardellec, A., Martinet, G., Barat, M., Lucas, B., Mezdari, F., Montagnon, L., Van-Oanh, N. T., Lavergne, L., Lachaize, A., Carpentier, Y., & Béroff, K. 2008, *J. Chem. Phys.*, 128, 124312
- van Dishoeck, E. F. 1988, in *Rate coefficients in astrochemistry. Proceedings of a Conference held in UMIST*. Ed. T. J. Millar and D. A. Williams (Kluwer Academic Publishers), 49–72
- van Dishoeck, E. F., Jonkheid, B., & van Hemert, M. C. 2006, *Faraday Discussions*, 133, 231
- Van Harrevelt, R., van Hemert, M. C., & Schatz, G. C. 2002, *J. Chemical Physics*, 116, 6002
- van Hemert, M. C. & van Dishoeck, E. F. 2008, *Chemical Physics*, 343, 292
- Vazquez, J., Harding, M. E., Gauss, J., & Stanton, J. F. 2009, *The journal of physical chemistry. A*, 113, 12447
- Viggiano, A. A., Ehlerding, A., Hellberg, F., Thomas, R. D., Zhaunerchyk, V., Geppert, W. D., Montaigne, H., Larsson, M., Kaminska, M., & Österdahl, F. 2005, *Journal of Chemical Physics*, 122, 226101
- Vinckier, C. 1979, *J. of Chem. Phys.*, 83, 1234
- Visser, R., van Dishoeck, E. F., & Black, J. H. 2009, *A&A*, 503, 323

- Vroom, D. A. & de Heer, F. J. 1969, *J. Chem. Phys.*, 50, 580
- Wakelam, V., Caselli, P., Ceccarelli, C., Herbst, E., & Castets, A. 2004, *A&A*, 422, 159
- Wakelam, V. & Herbst, E. 2008, *ApJ*, 680, 371
- Wakelam, V., Herbst, E., Le Bourlot, J., Hersant, F., Selsis, F., & Guilloteau, S. 2010a, *A&A*, 517, A21+
- Wakelam, V., Hersant, F., & Herpin, F. 2011, *A&A*, 529, A112+
- Wakelam, V., Loison, J. C., Herbst, E., Talbi, D., Quan, D., & Caralp, F. 2009, *A&A*, 495, 513
- Wakelam, V., Selsis, F., Herbst, E., & Caselli, P. 2005, *A&A*, 444, 883
- Wakelam, V., Smith, I. W. M., Herbst, E., Troe, J., Geppert, W. D., Linnartz, H., Öberg, K., Roueff, E., Agúndez, M., Pernot, P., Cuppen, H. M., Loison, J. C., & Talbi, D. 2010b, *Space Sci. Rev.*, 156, 13
- Weingartner, J. C. & Draine, B. T. 2001, *ApJ*, 548, 296
- Weisheit, J. C. & Upham, Jr., R. J. 1978, *MNRAS*, 184, 227
- Wennberg, P. O., Anderson, J. G., & Weisenstein, D. K. 1994, *Journal of Geophysical Research*, 99, 18839
- Woodall, J., Agúndez, M., Markwick-Kemper, A. J., & Millar, T. J. 2007, *A&A*, 466, 1197
- Woon, D. E. & Herbst, E. 2009, *ApJ*, 185, 273
- Zabarnick, S., Fleming, J. W., & Lin, M. C. 1989, *Int. J. Chem. Kinet.*, 21, 765

TABLE 1
TYPES OF REACTIONS DEFINED IN KIDA

Type of reaction	Abbreviation	Description
Direct cosmic ray processes	CR	Dissociation or ionization of species due to direct collision with cosmic ray particles.
Photo-processes induced by cosmic rays (secondary photons)	CRP	Dissociation or ionization of species due to UV photons emitted following H ₂ excitation.
Photo-processes	Phot	Dissociation or ionization of neutral species by UV photons with a standard interstellar UV field.
Bimolecular reactions	Bimo	Neutral-neutral ($A + B \rightarrow C + D$), ion-neutral ($A^+ + B \rightarrow C^+ + D$, $A^- + B \rightarrow C^- + D$) ^a , anion-cation ($A^+ + B^- \rightarrow C + D$) reactions and dissociative neutral attachment ($A + B \rightarrow AB^+ + e^-$)
Charge exchange reactions	CE	$A^+ + B \rightarrow A + B^+$ and $A^+ + B^- \rightarrow A + B$
Radiative associations	RA	Association reactions between two species (neutral or ionized) stabilized by the emission of a photon ($A + B \rightarrow AB + \text{photon}$ or $A^+ + B \rightarrow AB^+ + \text{photon}$).
Associative detachment	AD	Association of a neutral species and an anion, resulting in the ejection of the extra electron ($A^- + B \rightarrow AB + e^-$).
Electronic recombination and attachment	ER	Dissociative neutral recombination ($AB^+ + e^- \rightarrow A + B$) or the emission of a photon ($AB^+ + e^- \rightarrow AB + \text{photon}$) or the attachment of an electron ($A + e^- \rightarrow A^- + \text{photon}$).
Third-body assisted association	3-body	Association reactions between two species, stabilized by collision with a third body ($A + B + C \rightarrow AB + C$).

^aCharge exchange reactions have been defined as a separate type due to the particular microphysics behind this process.

Distribution	Formula
Lognormal	$k(T) = \exp(\ln k_0(T) + \ln F(T) * N(0, 1))$
Normal	$k(T) = k_0(T) + F(T) * N(0, 1)$
Uniform	$k(T) = k_0(T) + F(T) * 2 * (U(0, 1) - 0.5)$
Loguniform	$k(T) = \exp(\ln k_0(T) + \ln F(T) * 2 * (U(0, 1) - 0.5))$

Table 4: Generating random samples for MCUP. $U(0, 1)$ is a standard uniform random number generator (between 0 and 1); $N(0, 1)$ is a standard normal/gaussian random number generator (centered at 0; variance 1). In systems with a temperature gradient, and where multiple uncertain rate coefficients are managed simultaneously, care has to be taken that the same random number is used for the whole temperature range of a single reaction.

Table 2: The types of distributions and uncertainty factors used in KIDA.

Type	Distribution	F_0 meaning	F_0 unit
norm	Normal	stdev ¹ : $\text{Pr}^2(k_0 - F_0 \leq k \leq k_0 + F_0) \sim 68 \%$	as rate coefficient
unif	Uniform	half range: $\text{Pr}^2(k_0 - F_0 \leq k \leq k_0 + F_0) = 100 \%$	as rate coefficient
logn	Lognormal	geometric stdev ¹ : $\text{Pr}^2(k_0/F_0 \leq k \leq k_0 * F_0) \sim 68 \%$	no units
logu	Loguniform	geometric half range: $\text{Pr}^2(k_0/F_0 \leq k \leq k_0 * F_0) = 100 \%$	no units

¹ stdev = standard deviation

² Pr = Probability of assertion

Table 3: Reactions involving grains

Reaction	A ($\text{cm}^3 \text{s}^{-1}$)	B
$\text{C}^+ + \text{GRAIN}^- \rightarrow \text{C} + \text{GRAIN0}$	2.3×10^{-5}	0.5
$\text{Fe}^+ + \text{GRAIN}^- \rightarrow \text{Fe} + \text{GRAIN0}$	1.1×10^{-5}	0.5
$\text{H}^+ + \text{GRAIN}^- \rightarrow \text{H} + \text{GRAIN0}$	7.9×10^{-5}	0.5
$\text{He}^+ + \text{GRAIN}^- \rightarrow \text{He} + \text{GRAIN0}$	3.9×10^{-5}	0.5
$\text{Mg}^+ + \text{GRAIN}^- \rightarrow \text{Mg} + \text{GRAIN0}$	1.6×10^{-5}	0.5
$\text{N}^+ + \text{GRAIN}^- \rightarrow \text{N} + \text{GRAIN0}$	2.1×10^{-5}	0.5
$\text{Na}^+ + \text{GRAIN}^- \rightarrow \text{Na} + \text{GRAIN0}$	1.6×10^{-5}	0.5
$\text{O}^+ + \text{GRAIN}^- \rightarrow \text{O} + \text{GRAIN0}$	2.0×10^{-5}	0.5
$\text{S}^+ + \text{GRAIN}^- \rightarrow \text{S} + \text{GRAIN0}$	1.4×10^{-5}	0.5
$\text{Si}^+ + \text{GRAIN}^- \rightarrow \text{Si} + \text{GRAIN0}$	1.5×10^{-5}	0.5
$\text{H}_3^+ + \text{GRAIN}^- \rightarrow \text{H}_2 + \text{H} + \text{GRAIN0}$	4.6×10^{-5}	0.5
$\text{HCO}^+ + \text{GRAIN}^- \rightarrow \text{H} + \text{CO} + \text{GRAIN0}$	1.5×10^{-5}	0.5
$\text{e}^- + \text{GRAIN0} \rightarrow \text{GRAIN}^-$	3.4×10^{-3}	0.5




Fig. 1.— Abundances as a function of time for selected species computed for dense cloud conditions (see section 3.5) using three different chemical networks: udfa06 (blue dashed lines), osu-01-2009 (red dashed-dotted lines) and kida.uva.2011 (black lines). Dotted curves represent 2σ error bars on the abundances computed with the kida.uva.2011 network using uncertainty propagation in the reaction rate coefficients.