



*Research article*

## Bi-Bazilevič functions of order $\vartheta + i\delta$ associated with $(p, q)$ - Lucas polynomials

Ala Amourah<sup>1,\*</sup>, B. A. Frasin<sup>2</sup>, G. Murugusundaramoorthy<sup>3</sup> and Tariq Al-Hawary<sup>4</sup>

<sup>1</sup> Department of Mathematics, Faculty of Science and Technology, Irbid National University, Irbid, Jordan

<sup>2</sup> Faculty of Science, Department of Mathematics, Al al-Bayt University, Mafraq, Jordan

<sup>3</sup> Department of Mathematics, School of Advanced Sciences, VIT , Vellore 632014, Tamilnadu, India

<sup>4</sup> Department of Applied Science, Ajloun College, Al-Balqa Applied University, Ajloun 26816, Jordan

\* **Correspondence:** Email: alaammour@yahoo.com; Tel: 00962792290138.

**Abstract:** By means of  $(p, q)$ - Lucas polynomials, a class of Bazilevič functions of order  $\vartheta + i\delta$  in the open unit disk  $\mathbb{U}$  of analytic and bi-univalent functions is introduced. Further, we estimate coefficients bounds and Fekete-Szegö inequalities for functions belonging to this class. Several corollaries and consequences of the main results are also obtained.

**Keywords:** Bazilevič functions; Lucas polynomial; analytic functions; univalent functions; bi-univalent functions

**Mathematics Subject Classification:** 30C45

### 1. Introduction and preliminaries

Let  $\mathcal{A}$  indicate an analytic functions family, which is normalized under the condition  $f(0) = f'(0) - 1 = 0$  in  $\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$  and given by the following Taylor-Maclaurin series:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n. \tag{1.1}$$

Further, by  $\mathcal{S}$  we shall denote the class of all functions in  $\mathcal{A}$  which are univalent in  $\mathbb{U}$ .

With a view to recalling the principle of subordination between analytic functions, let the functions  $f$  and  $g$  be analytic in  $\mathbb{U}$ . Then we say that the function  $f$  is subordinate to  $g$  if there exists a Schwarz function  $w(z)$ , analytic in  $\mathbb{U}$  with

$$\omega(0) = 0, |\omega(z)| < 1, (z \in \mathbb{U})$$

such that

$$f(z) = g(\omega(z)).$$

We denote this subordination by

$$f < g \text{ or } f(z) < g(z).$$

In particular, if the function  $g$  is univalent in  $\mathbb{U}$ , the above subordination is equivalent to

$$f(0) = g(0), f(\mathbb{U}) \subset g(\mathbb{U}).$$

The Koebe-One Quarter Theorem [11] asserts that image of  $\mathbb{U}$  under every univalent function  $f \in \mathcal{A}$  contains a disc of radius  $\frac{1}{4}$ . Thus every univalent function  $f$  has an inverse  $f^{-1}$  satisfying  $f^{-1}(f(z)) = z$  and  $f^{-1}(f^{-1}(w)) = w$  ( $|w| < r_0(f^{-1}), r_0(f^{-1}) > \frac{1}{4}$ ), where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots \quad (1.2)$$

A function  $f \in \mathcal{A}$  is said to be bi-univalent functions in  $\mathbb{U}$  if both  $f$  and  $f^{-1}$  are univalent in  $\mathbb{U}$ . A function  $f \in \mathcal{S}$  is said to be bi-univalent in  $\mathbb{U}$  if there exists a function  $g \in \mathcal{S}$  such that  $g(z)$  is an univalent extension of  $f^{-1}$  to  $\mathbb{U}$ . Let  $\Lambda$  denote the class of bi-univalent functions in  $\mathbb{U}$ . The functions  $\frac{z}{1-z}, -\log(1-z), \frac{1}{2} \log\left(\frac{1+z}{1-z}\right)$  are in the class  $\Lambda$  (see details in [20]). However, the familiar Koebe function is not bi-univalent. Lewin [17] investigated the class of *bi-univalent* functions  $\Lambda$  and obtained a bound  $|a_2| \leq 1.51$ . Motivated by the work of Lewin [17], Brannan and Clunie [9] conjectured that  $|a_2| \leq \sqrt{2}$ . The coefficient estimate problem for  $|a_n|$  ( $n \in \mathbb{N}, n \geq 3$ ) is still open ([20]). Brannan and Taha [10] also worked on certain subclasses of the bi-univalent function class  $\Lambda$  and obtained estimates for their initial coefficients. Various classes of bi-univalent functions were introduced and studied in recent times, the study of *bi-univalent* functions gained momentum mainly due to the work of Srivastava et al. [20]. Motivated by this, many researchers [1], [4–8], [13–15], [20], [21], and [27–29], also the references cited there in) recently investigated several interesting subclasses of the class  $\Lambda$  and found non-sharp estimates on the first two Taylor-Maclaurin coefficients. Recently, many researchers have been exploring bi-univalent functions, few to mention Fibonacci polynomials, Lucas polynomials, Chebyshev polynomials, Pell polynomials, Lucas–Lehmer polynomials, orthogonal polynomials and the other special polynomials and their generalizations are of great importance in a variety of branches such as physics, engineering, architecture, nature, art, number theory, combinatorics and numerical analysis. These polynomials have been studied in several papers from a theoretical point of view (see, for example, [23–30] also see references therein).

We recall the following results relevant for our study as stated in [3].

Let  $p(x)$  and  $q(x)$  be polynomials with real coefficients. The  $(p, q)$ -Lucas polynomials  $\mathcal{L}_{p,q,n}(x)$  are defined by the recurrence relation

$$\mathcal{L}_{p,q,n}(x) = p(x)\mathcal{L}_{p,q,n-1}(x) + q(x)\mathcal{L}_{p,q,n-2}(x) \quad (n \geq 2),$$

from which the first few Lucas polynomials can be found as

$$\begin{aligned}\mathcal{L}_{p,q,0}(x) &= 2, \\ \mathcal{L}_{p,q,1}(x) &= p(x), \\ \mathcal{L}_{p,q,2}(x) &= p^2(x) + 2q(x), \\ \mathcal{L}_{p,q,3}(x) &= p^3(x) + 3p(x)q(x), \dots\end{aligned}\tag{1.3}$$

For the special cases of  $p(x)$  and  $q(x)$ , we can get the polynomials given  $\mathcal{L}_{x,1,n}(x) \equiv \mathcal{L}_n(x)$  Lucas polynomials,  $\mathcal{L}_{2x,1,n}(x) \equiv \mathcal{D}_n(x)$  Pell–Lucas polynomials,  $\mathcal{L}_{1,2x,n}(x) \equiv j_n(x)$  Jacobsthal–Lucas polynomials,  $\mathcal{L}_{3x,-2,n}(x) \equiv F_n(x)$  Fermat–Lucas polynomials,  $\mathcal{L}_{2x,-1,n}(x) \equiv T_n(x)$  Chebyshev polynomials first kind.

**Lemma 1.1.** [16] Let  $G\{\mathcal{L}(x)\}(z)$  be the generating function of the  $(p, q)$ –Lucas polynomial sequence  $\mathcal{L}_{p,q,n}(x)$ . Then,

$$G\{\mathcal{L}(x)\}(z) = \sum_{n=0}^{\infty} \mathcal{L}_{p,q,n}(x)z^n = \frac{2 - p(x)z}{1 - p(x)z - q(x)z^2}$$

and

$$\mathcal{G}_{\{\mathcal{L}(x)\}}(z) = G\{\mathcal{L}(x)\}(z) - 1 = 1 + \sum_{n=1}^{\infty} \mathcal{L}_{p,q,n}(x)z^n = \frac{1 + q(x)z^2}{1 - p(x)z - q(x)z^2}.$$

**Definition 1.2.** [22] For  $\vartheta \geq 0$ ,  $\delta \in \mathbb{R}$ ,  $\vartheta + i\delta \neq 0$  and  $f \in \mathcal{A}$ , let  $\mathcal{B}(\vartheta, \delta)$  denote the class of Bazilevič function if and only if

$$\operatorname{Re} \left[ \left( \frac{zf'(z)}{f(z)} \right) \left( \frac{f(z)}{z} \right)^{\vartheta+i\delta} \right] > 0.$$

Several authors have researched different subfamilies of the well-known Bazilevič functions of type  $\vartheta$  from various viewpoints (see [3] and [19]). For Bazilevič functions of order  $\vartheta + i\delta$ , there is no much work associated with Lucas polynomials in the literature. Initiating an exploration of properties of Lucas polynomials associated with Bazilevič functions of order  $\vartheta + i\delta$  is the main goal of this paper. To do so, we take into account the following definitions. In this paper motivated by the very recent work of Altinkaya and Yalcin [3] (also see [18]) we define a new class  $\mathcal{B}(\vartheta, \delta)$ , bi-Bazilevič function of  $\Lambda$  based on  $(p, q)$ –Lucas polynomials as below:

**Definition 1.3.** For  $f \in \Lambda$ ,  $\vartheta \geq 0$ ,  $\delta \in \mathbb{R}$ ,  $\vartheta + i\delta \neq 0$  and let  $\mathcal{B}(\vartheta, \delta)$  denote the class of Bi-Bazilevič functions of order  $\vartheta + i\delta$  if only if

$$\left[ \left( \frac{zf'(z)}{f(z)} \right) \left( \frac{f(z)}{z} \right)^{\vartheta+i\delta} \right] < \mathcal{G}_{\{\mathcal{L}(x)\}}(z) \quad (z \in \mathbb{U})\tag{1.4}$$

and

$$\left[ \left( \frac{zg'(w)}{g(w)} \right) \left( \frac{g(w)}{w} \right)^{\vartheta+i\delta} \right] < \mathcal{G}_{\{\mathcal{L}(x)\}}(w) \quad (w \in \mathbb{U}),\tag{1.5}$$

where  $\mathcal{G}_{\mathcal{L}_{p,q,n}}(z) \in \Phi$  and the function  $g$  is described as  $g(w) = f^{-1}(w)$ .

*Remark 1.4.* We note that for  $\delta = 0$  the class  $R(\vartheta, 0) = R(\vartheta)$  is defined by Altinkaya and Yalcin [2].

The class  $\mathcal{B}(0, 0) = \mathcal{S}_\Lambda^*$  is defined as follows:

**Definition 1.5.** A function  $f \in \Lambda$  is said to be in the class  $\mathcal{S}_\Lambda^*$ , if the following subordinations hold

$$\frac{zf'(z)}{f(z)} < \mathcal{G}_{\{\mathcal{L}(x)\}}(z) \quad (z \in \mathbb{U})$$

and

$$\frac{wg'(w)}{g(w)} < \mathcal{G}_{\{\mathcal{L}(x)\}}(w) \quad (w \in \mathbb{U})$$

where  $g(w) = f^{-1}(w)$ .

We begin this section by finding the estimates of the coefficients  $|a_2|$  and  $|a_3|$  for functions in the class  $\mathcal{B}(\vartheta, \delta)$ .

## 2. Coefficient bounds for the function class $\mathcal{B}(\vartheta, \delta)$

**Theorem 2.1.** Let the function  $f(z)$  given by 1.1 be in the class  $\mathcal{B}(\vartheta, \delta)$ . Then

$$|a_2| \leq \frac{p(x) \sqrt{2p(x)}}{\sqrt{|((\vartheta + i\delta)^2 + 3(\vartheta + i\delta) + 2) - 2(\vartheta + i\delta + 1)^2| p^2(x) - 4q(x)(\vartheta + i\delta + 1)^2|}}.$$

and

$$|a_3| \leq \frac{p^2(x)}{(\vartheta + 1)^2 + \delta^2} + \frac{p(x)}{\sqrt{(\vartheta + 2)^2 + \delta^2}}.$$

*Proof.* Let  $f \in \mathcal{B}(\vartheta, \delta)$  there exist two analytic functions  $u, v : \mathbb{U} \rightarrow \mathbb{U}$  with  $u(0) = 0 = v(0)$ , such that  $|u(z)| < 1$ ,  $|v(w)| < 1$ , we can write from (1.4) and (1.5), we have

$$\left[ \left( \frac{zf'(z)}{f(z)} \right) \left( \frac{f(z)}{z} \right)^{\vartheta+i\delta} \right] = \mathcal{G}_{\{\mathcal{L}(x)\}}(z) \quad (z \in \mathbb{U}), \quad (2.1)$$

and

$$\left[ \left( \frac{zg'(w)}{g(w)} \right) \left( \frac{g(w)}{w} \right)^{\vartheta+i\delta} \right] = \mathcal{G}_{\{\mathcal{L}(x)\}}(w) \quad (w \in \mathbb{U}), \quad (2.2)$$

It is fairly well known that if

$$|u(z)| = |u_1z + u_2z^2 + \dots| < 1,$$

and

$$|v(w)| = |v_1w + v_2w^2 + \dots| < 1,$$

then

$$|u_k| \leq 1 \quad \text{and} \quad |v_k| \leq 1 \quad (k \in \mathbb{N}),$$

so we have,

$$\begin{aligned}\mathcal{G}_{\{\mathcal{L}(x)\}}(u(z)) &= 1 + \mathcal{L}_{p,q,1}(x)u(z) + \mathcal{L}_{p,q,2}(x)u^2(z) + \dots \\ &= 1 + \mathcal{L}_{p,q,1}(x)u_1z + [\mathcal{L}_{p,q,1}(x)u_2 + \mathcal{L}_{p,q,2}(x)u_1^2]z^2 + \dots\end{aligned}\quad (2.3)$$

and

$$\begin{aligned}\mathcal{G}_{\{\mathcal{L}(x)\}}(v(w)) &= 1 + \mathcal{L}_{p,q,1}(x)v(w) + \mathcal{L}_{p,q,2}(x)v^2(w) + \dots \\ &= 1 + \mathcal{L}_{p,q,1}(x)v_1w + [\mathcal{L}_{p,q,1}(x)v_2 + \mathcal{L}_{p,q,2}(x)v_1^2]w^2 + \dots\end{aligned}\quad (2.4)$$

From the equalities (2.1) and (2.2), we obtain that

$$\left[ \left( \frac{zf'(z)}{f(z)} \right) \left( \frac{f(z)}{z} \right)^{\vartheta+i\delta} \right] = 1 + \mathcal{L}_{p,q,1}(x)u_1z + [\mathcal{L}_{p,q,1}(x)u_2 + \mathcal{L}_{p,q,2}(x)u_1^2]z^2 + \dots, \quad (2.5)$$

and

$$\left[ \left( \frac{zg'(w)}{g(w)} \right) \left( \frac{g(w)}{w} \right)^{\vartheta+i\delta} \right] = 1 + \mathcal{L}_{p,q,1}(x)v_1w + [\mathcal{L}_{p,q,1}(x)v_2 + \mathcal{L}_{p,q,2}(x)v_1^2]w^2 + \dots, \quad (2.6)$$

It follows from (2.5) and (2.6) that

$$(\vartheta + i\delta + 1) a_2 = \mathcal{L}_{p,q,1}(x)u_1, \quad (2.7)$$

$$\frac{(\vartheta + i\delta - 1)(\vartheta + i\delta + 2)}{2} a_2^2 - (\vartheta + i\delta + 2) a_3 = \mathcal{L}_{p,q,1}(x)u_2 + \mathcal{L}_{p,q,2}(x)u_1^2, \quad (2.8)$$

and

$$-(\vartheta + i\delta + 1) a_2 = \mathcal{L}_{p,q,1}(x)v_1, \quad (2.9)$$

$$\frac{(\vartheta + i\delta + 2)(\vartheta + i\delta + 3)}{2} a_2^2 + (\vartheta + i\delta + 2) a_3 = \mathcal{L}_{p,q,1}(x)v_2 + \mathcal{L}_{p,q,2}(x)v_1^2. \quad (2.10)$$

From (2.7) and (2.9)

$$u_1 = -v_1 \quad (2.11)$$

and

$$2(\vartheta + i\delta + 1)^2 a_2^2 = \mathcal{L}_{p,q,1}^2(x)(u_1^2 + v_1^2), \quad (2.12)$$

by adding (2.8) to (2.10), we get

$$\left( (\vartheta + i\delta)^2 + 3(\vartheta + i\delta) + 2 \right) a_2^2 = \mathcal{L}_{p,q,1}(x)(u_2 + v_2) + \mathcal{L}_{p,q,2}(x)(u_1^2 + v_1^2), \quad (2.13)$$

by using (2.12) in equality (2.13), we have

$$\left[ \left( (\vartheta + i\delta)^2 + 3(\vartheta + i\delta) + 2 \right) - \frac{2\mathcal{L}_{p,q,2}(x)(\vartheta + i\delta + 1)^2}{\mathcal{L}_{p,q,1}^2(x)} \right] a_2^2 = \mathcal{L}_{p,q,1}(x)(u_2 + v_2),$$

$$a_2^2 = \frac{\mathcal{L}_{p,q,1}^3(x)(u_2 + v_2)}{\left[ \left( (\vartheta + i\delta)^2 + 3(\vartheta + i\delta) + 2 \right) \mathcal{L}_{p,q,1}^2(x) - 2\mathcal{L}_{p,q,2}(x)(\vartheta + i\delta + 1)^2 \right]}. \quad (2.14)$$

Thus, from (1.3) and (2.14) we get

$$|a_2| \leq \frac{p(x) \sqrt{2p(x)}}{\sqrt{|((\vartheta + i\delta)^2 + 3(\vartheta + i\delta) + 2) - 2(\vartheta + i\delta + 1)^2| p^2(x) - 4q(x)(\vartheta + i\delta + 1)^2}}.$$

Next, in order to find the bound on  $|a_3|$ , by subtracting (2.10) from (2.8), we obtain

$$\begin{aligned} 2(\vartheta + i\delta + 2)a_3 - 2(\vartheta + i\delta + 2)a_2^2 &= \mathcal{L}_{p,q,1}(x)(u_2 - v_2) + \mathcal{L}_{p,q,2}(x)(u_1^2 - v_1^2) \\ 2(\vartheta + i\delta + 2)a_3 &= \mathcal{L}_{p,q,1}(x)(u_2 - v_2) + 2(\vartheta + i\delta + 2)a_2^2 \\ a_3 &= \frac{\mathcal{L}_{p,q,1}(x)(u_2 - v_2)}{2(\vartheta + i\delta + 2)} + a_2^2 \end{aligned} \quad (2.15)$$

Then, in view of (2.11) and (2.12), we have from (2.15)

$$a_3 = \frac{\mathcal{L}_{p,q,1}^2(x)}{2(\vartheta + i\delta + 2)^2}(u_1^2 + v_1^2) + \frac{\mathcal{L}_{p,q,1}(x)}{2(\vartheta + i\delta + 2)}(u_2 - v_2).$$

$$\begin{aligned} |a_3| &\leq \frac{p^2(x)}{|\vartheta + i\delta + 1|^2} + \frac{p(x)}{|\vartheta + i\delta + 2|} \\ &= \frac{p^2(x)}{(\vartheta + 1)^2 + \delta^2} + \frac{p(x)}{\sqrt{(\vartheta + 2)^2 + \delta^2}} \end{aligned}$$

This completes the proof.  $\square$

Taking  $\delta = 0$ , in Theorem 2.1, we get the following corollary.

**Corollary 2.2.** *Let the function  $f(z)$  given by (1.1) be in the class  $\mathcal{B}(\vartheta)$ . Then*

$$|a_2| \leq \frac{p(x) \sqrt{2p(x)}}{\sqrt{|(\vartheta^2 + 3\vartheta + 2) - 2(\vartheta + 1)^2| p^2(x) - 4q(x)(\vartheta + 1)^2}}$$

and

$$|a_3| \leq \frac{p^2(x)}{(\vartheta + 2)^2} + \frac{p(x)}{\vartheta + 2}$$

Also, taking  $\vartheta = 0$  and  $\delta = 0$ , in Theorem 2.1, we get the results given in [18].

### 3. Fekete-Szegő inequality for the class $\mathcal{B}(\vartheta, \delta)$

Fekete-Szegő inequality is one of the famous problems related to coefficients of univalent analytic functions. It was first given by [12], the classical Fekete-Szegő inequality for the coefficients of  $f \in \mathcal{S}$  is

$$|a_3 - \mu a_2^2| \leq 1 + 2 \exp(-2\mu/(1 - \mu)) \text{ for } \mu \in [0, 1).$$

As  $\mu \rightarrow 1^-$ , we have the elementary inequality  $|a_3 - \mu a_2^2| \leq 1$ . Moreover, the coefficient functional

$$\zeta_\mu(f) = a_3 - \mu a_2^2$$

on the normalized analytic functions  $f$  in the unit disk  $\mathbb{U}$  plays an important role in function theory. The problem of maximizing the absolute value of the functional  $\zeta_\mu(f)$  is called the Fekete-Szegő problem.

In this section, we are ready to find the sharp bounds of Fekete-Szegő functional  $\zeta_\mu(f)$  defined for  $f \in \mathcal{B}(\vartheta, \delta)$  given by (1.1).

**Theorem 3.1.** *Let  $f$  given by (1.1) be in the class  $\mathcal{B}(\vartheta, \delta)$  and  $\mu \in \mathbb{R}$ . Then*

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{p(x)}{\sqrt{(\vartheta+2)^2 + \delta^2}}, & 0 \leq |h(\mu)| \leq \frac{1}{2\sqrt{(\vartheta+2)^2 + \delta^2}} \\ 2p(x)|h(\mu)|, & |h(\mu)| \geq \frac{1}{2\sqrt{(\vartheta+2)^2 + \delta^2}} \end{cases}$$

where

$$h(\mu) = \frac{\mathcal{L}_{p,q,1}^2(x)(1-\mu)}{\left((\vartheta+i\delta)^2 + 3(\vartheta+i\delta) + 2\right) \mathcal{L}_{p,q,1}^2(x) - 2\mathcal{L}_{p,q,2}(x)(\vartheta+i\delta+1)^2}.$$

*Proof.* From (2.14) and (2.15), we conclude that

$$\begin{aligned} a_3 - \mu a_2^2 &= (1-\mu) \frac{\mathcal{L}_{p,q,1}^3(x)(u_2 + v_2)}{\left[\left((\vartheta+i\delta)^2 + 3(\vartheta+i\delta) + 2\right) \mathcal{L}_{p,q,1}^2(x) - 2\mathcal{L}_{p,q,2}(x)(\vartheta+i\delta+1)^2\right]} \\ &\quad + \frac{\mathcal{L}_{p,q,1}(x)}{2(\vartheta+i\delta+2)}(u_2 - v_2) \\ &= \mathcal{L}_{p,q,1}(x) \left[ \left( h(\mu) + \frac{1}{2(\vartheta+i\delta+2)} \right) u_2 + \left( h(\mu) - \frac{1}{2(\vartheta+i\delta+2)} \right) v_2 \right] \end{aligned}$$

where

$$h(\mu) = \frac{\mathcal{L}_{p,q,1}^2(x)(1-\mu)}{\left((\vartheta+i\delta)^2 + 3(\vartheta+i\delta) + 2\right) \mathcal{L}_{p,q,1}^2(x) - 2\mathcal{L}_{p,q,2}(x)(\vartheta+i\delta+1)^2}.$$

Then, in view of (1.3), we obtain

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{p(x)}{\sqrt{(\vartheta+2)^2 + \delta^2}}, & 0 \leq |h(\mu)| \leq \frac{1}{2\sqrt{(\vartheta+2)^2 + \delta^2}} \\ 2p(x)|h(\mu)|, & |h(\mu)| \geq \frac{1}{2\sqrt{(\vartheta+2)^2 + \delta^2}} \end{cases}$$

□

We end this section with some corollaries.

Taking  $\mu = 1$  in Theorem 3.1, we get the following corollary.

**Corollary 3.2.** *If  $f \in \mathcal{B}(\vartheta, \delta)$ , then*

$$|a_3 - a_2^2| \leq \frac{p(x)}{\sqrt{(\vartheta + 2)^2 + \delta^2}}.$$

Taking  $\delta = 0$  in Theorem 3.1, we get the following corollary.

**Corollary 3.3.** *Let  $f$  given by (1.1) be in the class  $\mathcal{B}(\vartheta, 0)$ . Then*

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{p(x)}{\vartheta+2}, & 0 \leq |h(\mu)| \leq \frac{1}{2(\vartheta+2)} \\ 2p(x)|h(\mu)|, & |h(\mu)| \geq \frac{1}{2(\vartheta+2)} \end{cases}$$

Also, taking  $\vartheta = 0$ ,  $\delta = 0$  and  $\mu = 1$  in Theorem 3.1, we get the following corollary.

**Corollary 3.4.** *Let  $f$  given by (1.1) be in the class  $\mathcal{B}$ . Then*

$$|a_3 - a_2^2| \leq \frac{p(x)}{2}.$$

### Conflict of interest

All authors declare no conflicts of interest in this paper.

### References

1. I. Aldawish, T. Al-Hawary, B. A. Frasin, Subclasses of bi-univalent functions defined by Frasin differential operator, *Mathematics*, **8** (2020), 783.
2. Ş. Altinkaya, S. Yalçın, On the Chebyshev polynomial coefficient problem of bi-Bazilevic function, *TWMS J. App. Eng. Math.*, **10** (2020), 251–258.
3. Ş. Altinkaya, S. Yalçın, On the  $(p, q)$ -Lucas polynomial coefficient bounds of the bi-univalent function class  $\sigma$ , *Boletín de la Sociedad Matemática Mexicana*, **25** (2019), 567–575.
4. Ş. Altinkaya, S. Yalçın, Coefficient estimates for two new subclasses of bi-univalent functions with respect to symmetric points, *J. Funct. Spaces*, **2015** (2015), 1–5.
5. A. Amourah, *Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to  $(p, q)$ -Lucas polynomials*, TWMS Journal of Applied and Engineering (in press).
6. A. Amourah, , Faber polynomial coefficient estimates for a class of analytic bi-univalent functions, *AIP Conference Proceedings*, **2096** (2019), 020024. AIP Publishing LLC.
7. A. Amourah, *Initial bounds for analytic and bi-univalent functions by means of  $(p, q)$ - Chebyshev polynomials defined by differential operator*, *General Letters in Mathematics*, **7** (2019), 45–51.
8. A. Amourah, M. Illafe, A Comprehensive subclass of analytic and Bi-univalent functions associated with subordination, *Palestine Journal of Mathematics*, **9** (2020), 187–193.
9. D. Brannan, J. Clunie, *Aspects of contemporary complex analysis*, Academic Press, New York, 1980.



10. D. Brannan, T. S. Taha, On some classes of bi-univalent functions, *Mathematical Analysis and Its Applications*, (1988), 53–60.
11. P. L. Duren, *Univalent Functions, Grundlehren der Mathematischen Wissenschaften*, Springer, New York, USA, 1983.
12. M. Fekete, G. Szegő, Eine Bemerkung ber Ungerade Schlichte Funktionen, *J. London Math. Soc.*, **1** (1933), 85–89.
13. B. A. Frasin, M. K. Aouf, New subclasses of bi-univalent functions, *Appl. Math. Lett.*, **24** (2011), 1569–1573.
14. B. A. Frasin, Coefficient bounds for certain classes of bi-univalent functions, *Hacet. J. Math. Stat.*, **43** (2014), 383–389.
15. B. A. Frasin, T. Al-Hawary, Initial Maclaurin coefficients bounds for new subclasses of bi-univalent functions, *Theory Appl. Math. Comp. Sci.*, **5** (2015), 186–193.
16. G. Y. Lee, M. Aşçı, Some properties of the  $(p, q)$ -Fibonacci and  $(p, q)$ -Lucas polynomials, *J. Appl. Math.*, **2012** (2012), 1–18.
17. M. Lewin, On a coefficient problem for bi-univalent functions, *Proc. Amer. Math. Soc.*, **18** (1967), 63–68.
18. G. Murugusundaramoorthy, S. Yalçın, On the  $\lambda$ -Pseudo-bi-starlike functions related to  $(p, q)$ -Lucas polynomial, *Libertas Mathematica (newseries)*, **39** (2019), 79–88.
19. H. M. Srivastava, G. Murugusundaramoorthy, K. Vijaya, Coefficient estimates for some families of bi-Bazilevič functions of the Ma-Minda type involving the Hohlov operator, *J. Class. Anal.*, **2** (2013), 167–181.
20. H. M. Srivastava, A. K. Mishra, P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.*, **23** (2010), 1188–1192.
21. H. M. Srivastava, S. Bulut, CM. ağlar, N. Yağmur, Coefficient estimates for a general subclass of analytic and bi-univalent functions, *Filomat*, **27** (2013), 831–842.
22. T. Sheil-Small, On Bazilevič functions, *The Quarterly Journal of Mathematics*, **23** (1972), 135–142.
23. P. Vellucci, A. M. Bersani, The class of Lucas–Lehmer polynomials, *Rend. Mat. Appl. serie VII*, **37** (2016), 43–62.
24. P. Vellucci, A. M. Bersani, Orthogonal polynomials and Riesz bases applied to the solution of Love’s equation, *Math. Mech. Complex Syst.*, **4** (2016), 55–66.
25. P. Vellucci, A. M. Bersani, Ordering of nested square roots of according to the Gray code, *Ramanujan J.*, **45** (2018), 197–210.
26. T. Wang, W. Zhang, Some identities involving Fibonacci, Lucas polynomials and their applications, *Bull. Math. Soc. Sci. Math. Roum.*, **55** (2012), 95–103.
27. F. Yousef, T. Al-Hawary, G. Murugusundaramoorthy, Fekete-Szegő functional problems for some subclasses of bi-univalent functions defined by Frasin differential operator, *Afr. Mat.*, **30** (2019), 495–503.

28. F. Yousef, B. Frasin, T. Al-Hawary, Fekete-Szegő inequality for analytic and bi-univalent functions subordinate to Chebyshev polynomials, *Filomat*, **32** (2018), 3229–3236.
29. F. Yousef, S. Alroud, M. Illafe, A comprehensive subclass of bi-univalent functions associated with Chebyshev polynomials of the second kind, *Boletín de la Sociedad Matemática Mexicana*, (2019), 1–11.
30. A. Zireh, E. Analouei Adegani, S. Bulut, Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions defined by subordination, *Bull. Belg. Math. Soc. Simon Stevin*, **23** (2016), 487–504.



AIMS Press

©2021 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)