

# Candidate OP Phyla: Importance, Ecology and Cultivation Prospects

M. Rohini Kumar · V. S. Saravanan

Received: 21 September 2010 / Accepted: 22 December 2010 / Published online: 25 January 2011  
© Association of Microbiologists of India 2011

**Abstract** OP phyla were created in the domain bacteria, based on the group of 16S rRNA gene sequences recovered from the Obsidian Pool. However, due to the lack of cultured representative it is referred to as candidate phyla. Wider ecological occurrence was predicted for the OP phyla, especially OP3, OP10 and OP11. Recently, members of phylum OP5 and OP10 were cultured, providing clues to their cultivation prospects. At last the bio-prospecting potentials of the OP members are discussed herein.

**Keywords** Candidate OP group phyla · Uncultivated · Obsidian pool

## Introduction

Nearly a decade ago, some novel 16S rRNA gene sequences were recovered from the Obsidian pool, a hot spring (75–95°C) located in Yellow Stone National Park (Fig. 1), rich in iron, sulfide, carbon dioxide and hydrogen, supporting a unique ecosystem dominated by lithotrophic communities [1, 2]. Those sequences could not be assigned to any other phylum known at that time, thus, novel OP group phyla were created [1]. Based on the variability of the sequences recovered, 12 OP phyla were initially created [1]. However,

currently only seven phyla are validated (OP 1, 3, 5, 8, 9, 10 and 11). In this letter, the importance of Candidate phyla, ecological occurrence of OP phyla, recent cultivation efforts to underpin the OP10 and OP5 phylotypes and possible biotechnological applications are emphasized.


## Radiance of Bacterial Phyla

The enormous diversity of bacteria in nature was revealed through the community sampling of ribosomal RNA genes (16S rRNA). It is believed that at least 100 phyla of bacteria are probably in existence. Among these, bacteria belonging to at least 40 phyla are well identified [3], among these validly identified phyla, only about 25 have cultured representatives (Fig. 2). The remaining phyla are referred to as Candidate, denoting phyla without cultured representatives, assigned by directly obtained gene sequences (16S rRNA sequences), at present they number 39 [4]. Thus candidate status indicates their unconfirmed position as a valid phylum due to the absence of cultured representatives. However, their members are abundant and widely distributed in soil ecosystems and ecologically they may play key roles in soil processes [5]. Candidatus nature is observed even among bacterial groups occurring in defined phyla including the dominant soil dwelling Acidobacteria [6] and Planctomycetes accommodating the anaerobic ammonium oxidizers (ANAMMOX) an important player in the nitrogen cycle [7]. The promise of being able to study the genome without cultivation led to the emergence of metagenomics as a popular, cultivation independent approach to bacterial community studies. Paradoxically, completed genomes posed a concern due to genomic micro heterogeneity and high species diversity within samples. Ultimately, a situation has arisen where 40% or more of the proteins in any

M. Rohini Kumar  
Hex-226 Lab, School of Bio Sciences and Technology,  
Vellore Institute of Technology, University,  
Vellore, Tamilnadu 632 014, India

V. S. Saravanan (✉)  
Department of Microbiology, Indira Gandhi College  
of Arts and Science, Puducherry 605 007, India  
e-mail: saraphd2003@yahoo.com





**Fig. 3** Phase-contrast (a) and transmission electron (b–h) images of *C. exile*. Negatively stained cells are shown in b and c. Ultra-thin sections of the cells are shown in d–h. The cellular membrane and outer envelope are indicated by *open and solid arrows*, respectively (d) [19]

Recently, a few strains belonging to the members of OP5 and OP10 have been cultured [9, 14, 18]. Interestingly, their culturing conditions give us clues to the successful isolation of other members belonging to OP series. Members of OP phyla are ovoid rod (OP10) to filamentous (OP5) in shape, litho or chemo organotrophs and represent either aerobic or anaerobic form of metabolism.

A member of OP5 phylum has been isolated from hot springs of Japan. It is a Gram negative, filamentous (Fig. 3) chemoheterotrophic and anaerobic bacteria. It has been named as *Caldisericum exile* and its taxonomic position was validly published [9, 19].

In another study, members belonging to OP10 phylum were successfully isolated from the geothermally heated soils of Taupo volcanic zone, New Zealand. These isolates (P488 and T49) are thermophilic in nature, and grow only in the mineral medium. They are aerobes and able to utilize complex polysaccharides like gellan, xanthan and carboxymethylcellulose [18]. The success behind their cultivation was mainly attributed to maintenance of low pH medium and the use of gellan as gelling agent as an alternate to agar. Recently, another member of this phylum was cultured from the rhizoplanes of the aquatic plant, *Phragmites australis* (reed), commonly grown in the mesophilic fresh water lakes of Japan using low nutrient medium [14]. Both New Zealand and Japanese OP10 strains are chemoheterotrophic and aerobic however, the mesophilic strain isolated from Japan can be easily differentiated from those recovered from geothermal soils of

New Zealand based on the temperature, pH range of growth, complex polysaccharide utilization and differences in certain biochemical features. Polyphasic taxonomic analysis placed the Japanese strain into a novel genera and it was validly published as *Armatimonas rosea*. Besides, the candidate OP10 phylum was renamed as phylum *Armatimonadetes* [14].

### Bioprospecting Potentials

Based on the habitat diversity of OP phylum series, it can be hypothesized that it is a highly widespread and ecologically diverse phyla. Further, cultivation studies could pave the way to understanding the genetic, metabolic and species diversity in these phyla. The isolates obtained till date may throw more light on the cultivation aspects of OP phyla in the future, further phylum level identity could be better resolved for various OP phyla. The available evidence envisages their putative role in biogeochemical cycling, which is unexplored at present [12]. Interestingly, sequences of OP phyla have been recovered from high or low temperature environments including hydrocarbon and chlorinated solvent contaminated aquifers [11]. Therefore, bioprospecting potentials in the areas of extremophilic enzymes development and intrinsic bioremediation approaches could be explored on cultivation of those OP members or identifying the functional genes associated with their survival in those environments.

## References

- Hugenholtz P, Pitulle C, Hershberger KL, Pace NR (1998) Novel division level bacterial diversity in a Yellowstone hot spring. *J Bacteriol* 180:366–376
- Hugenholtz P, Goebel BM, Pace NR (1998) Impact of culture-independent studies on the emerging phylogenetic view of bacterial diversity. *J Bacteriol* 180:4765–4774
- Ley RE, Harris JK, Wilcox J, Spear JR, Miller SR, Bebout BM, Maresca JA, Bryant DA, Sogin ML, Pace NR (2006) Unexpected diversity and complexity of the Guerrero Negro hypersaline microbial mat. *Appl Environ Microbiol* 72:3685–3695
- Glöckner J, Kube M, Shrestha PM, Weber M, Glöckner FO, Reinhardt R, Liesack W (2010) Phylogenetic diversity and metagenomics of candidate division OP3. *Environ Microbiol* 12:1218–1229
- Van Elsas JD, Torsvik V, Hartmann A, Øvreås L, Jansson JK (2007) The bacteria and archaea in soil. In: Van Elsas JD, Jansson JK, Trevors JT (eds) *Modern soil microbiology*, 2nd edn. Taylor and Francis, London, pp 84–102
- Dedysh SN, Pankratov TA, Belova SE, Kulichevskaya IS, Liesack W (2006) Phylogenetic analysis and in situ identification of bacteria community composition in an acidic *Sphagnum* peat bog. *Appl Environ Microbiol* 72:2110–2117
- Kuenen GJ (2008) Anammox bacteria: from discovery to application. *Nat Rev Microbiol* 6:320–326
- Nichols D (2007) Cultivation gives context to the microbial ecologist. *FEMS Microbiol Ecol* 60:351–357
- Mori K, Sunamura M, Yanagawa K, Ishibashi J, Miyoshi Y, Iino T, Suzuki K, Urabe T (2008) First cultivation and ecological investigation of a bacterium affiliated with the candidate phylum OP5 from hot springs. *Appl Environ Microbiol* 74:6223–6229
- Rastogi G, Osman S, Kukkadapu R, Engelhard M, Vaishampayan PA, Andersen GL, Sani RK (2010) Microbial and mineralogical characterizations of soils collected from the deep biosphere of the former Homestake gold mine, South Dakota. *Microb Ecol* 60:539–550
- Dojka MA, Hugenholtz P, Haack SK, Pace NR (1998) Microbial diversity in a hydrocarbon- and chlorinated-solvent contaminated aquifer undergoing intrinsic bioremediation. *Appl Environ Microbiol* 64:3869–3877
- Harris JK, Kelley ST, Pace NR (2004) New perspective on uncultured bacterial phylogenetic division OP11. *Appl Environ Microbiol* 70:845–849
- Stott MB, Saito JA, Crowe MA, Dunfield PF, Hou S, Nakasone E, Daughney CJ, Smirnova AV, Mountain BW, Takai K, Alam M (2008) Culture-independent characterization of a novel microbial community at a hydrothermal vent at Brothers volcano, Kermadec arc, New Zealand. *J Geophys Res* 113:B08S06
- Tamaki H, Tanaka Y, Matsuzawa H, Muramatsu M, Meng XY, Hanada S, Mori K, Kamagata Y (2011) *Armatimonas rosea* gen. nov., sp. nov., a Gram-negative, aerobic, chemoheterotrophic bacterium of a novel bacterial phylum, *Armatimonadetes* phyl. nov., formally called the candidate phylum OP10. *Int J Syst Evol Microbiol*. doi:10.1099/ijs.0.025643-0
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35:7188–7196
- Portillo MC, Gonzalez JM (2009) Members of the Candidate Division OP10 are spread in a variety of environments. *World J Microbiol Biotechnol* 25:347–353
- Schloss PD, Handelsman J (2004) Status of the microbial census. *Microbiol Mol Biol Rev* 68:686–691
- Stott MB, Crowe MA, Mountain BW, Smirnova AV, Hou S, Alam M, Dunfield PF (2008) Isolation of novel bacteria, including a candidate division, from geothermal soils in New Zealand. *Environ Microbiol* 10:2030–2041
- Mori K, Yamaguchi K, Sakiyama Y, Urabe T, Suzuki K (2009) *Caldisericum exile* gen. nov., sp. nov., an anaerobic, thermophilic, filamentous bacterium of a novel bacterial phylum, *Caldiserica* phyl. nov., originally called the candidate phylum OP5, and description of *Caldiseriaceae* fam. nov., *Caldisericales* ord. nov. and *Caldisericia* classis nov. *Int J Syst Evol Microbiol* 59:2894–2898