

C. SELVARAJ, O. S. BABU and G. MURUGUSUNDARAMOORTHY

Coefficient bounds for some subclasses of p -valently starlike functions

ABSTRACT. For functions of the form $f(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} z^{p+n}$ we obtain sharp bounds for some coefficients functionals in certain subclasses of starlike functions. Certain applications of our main results are also given. In particular, Fekete–Szegö-like inequality for classes of functions defined through extended fractional differintegrals are obtained.

1. Introduction. Let \mathcal{A}_p denote the class of all functions of the form

$$(1.1) \quad f(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} z^{p+n} \quad (p \in \mathbb{N} = \{1, 2, 3, \dots\})$$

which are analytic in the open disk $\Delta = \{z \in \mathbb{C} : |z| < 1\}$ and let $\mathcal{A} = \mathcal{A}_1$. For $f(z)$ given by (1.1) and $g(z)$ given by $g(z) = z^p + \sum_{n=1}^{\infty} b_{p+n} z^{p+n}$, their *convolution* (or Hadamard product), denoted by $f * g$, is defined as

$$(f * g)(z) = z^p + \sum_{n=1}^{\infty} a_{p+n} b_{p+n} z^{p+n}.$$

Given two functions f and g , which are analytic in Δ , the function f is said to be *subordinate* to g in Δ , written $f \prec g$ or $f(z) \prec g(z)$, if there exists a Schwarz function $w(z)$, analytic in Δ with $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = g(w(z))$, $z \in \Delta$. In particular, if the function g is univalent in Δ , the above subordination is equivalent to $f(0) = g(0)$ and $f(\Delta) \subset g(\Delta)$.

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic functions, starlike functions, convex functions, p -valent functions, subordination, convolution, Fekete–Szegö inequality.

Definition 1.1. Let $\phi(z)$ be an analytic function with positive real part in the unit disk Δ with $\phi(0) = 1$ and $\phi'(0) > 0$ that maps Δ onto a region starlike with respect to 1 and symmetric with respect to the real axis. The class $\mathcal{M}_p(\lambda; \phi)$ is the subclass of \mathcal{A}_p consisting of functions $f(z)$ satisfying

$$(1.2) \quad \frac{\frac{1}{p} z f'(z)}{(1 - \lambda) z^p + \lambda f(z)} \prec \phi(z) \quad (z \in \Delta, 0 < \lambda \leq 1).$$

As special cases, let

$$\mathcal{M}_p(1; \phi) = S_p^*(\phi) = \left\{ f(z) \in \mathcal{A}_p : \frac{1}{p} \frac{z f'(z)}{f(z)} \prec \phi(z) \right\},$$

$$\mathcal{M}_1(1; \phi) = S^*(\phi) = \left\{ f(z) \in \mathcal{A} : \frac{z f'(z)}{f(z)} \prec \phi(z) \right\}.$$

When $\phi(z) = \frac{1+Az}{1+Bz}$, $-1 \leq B < A \leq 1$, we denote the subclass $\mathcal{M}_p(\lambda; \phi)$ by $\mathcal{M}_p(\lambda; A, B)$. The class $\mathcal{M}_1(1; A, B) = S^*[A, B]$ was studied by Janowski [2]. For $0 \leq \alpha < 1$, let $\mathcal{M}_p(\lambda; \alpha) = \mathcal{M}_p(\lambda; 1 - 2\alpha, -1)$.

For a fixed analytic function $g \in \mathcal{A}_p$ with positive coefficients, define the class $\mathcal{M}_{p,g}(\lambda; \phi)$ as the class of all functions $f \in \mathcal{A}_p$ satisfying $f * g \in \mathcal{M}_p(\lambda; \phi)$. This class includes as special cases several other classes studied in the literature. For example, when $g(z) = z^p + \sum_{n=1}^{\infty} \frac{p+n}{p} z^{p+n}$, the class $\mathcal{M}_{p,g}(1; \phi)$ reduces to the class $C_p(1 : \phi) = C_p(\phi)$ consisting of functions $f \in \mathcal{A}_p$ satisfying

$$(1.3) \quad \frac{1}{p} \left(1 + \frac{z f''(z)}{f'(z)} \right) \prec \phi(z), \quad z \in \Delta.$$

The classes $S^*(\phi)$ and $C(\phi) = C_1(\phi)$ were introduced and studied by Ma and Minda [4]. They have obtained the Fekete–Szegö inequality for functions in the class $C(\phi)$. Since $f(z) \in C(\phi)$ if and only if $z f'(z) \in S^*(\phi)$, we get the Fekete–Szegö inequality for functions in the class $S^*(\phi)$. For a brief history of Fekete–Szegö problem for classes of starlike, convex and close-to-convex functions see the recent paper by Srivastava et al. [10].

Let Ω be the class of analytic functions of the form

$$(1.4) \quad w(z) = w_1 z + w_2 z^2 + \dots$$

in the unit disk Δ satisfying the condition $|w(z)| < 1$.

There has been triggering interest in the literature (see [1, 3, 4, 9, 10]) to define certain subclasses of analytic functions and to discuss Fekete–Szegö inequalities. Making use of the techniques, in this paper we defined two new classes $\mathcal{M}_p(\lambda; \phi)$ and $\mathcal{M}_{p,g}(\lambda; \phi)$ to obtain Fekete–Szegö inequalities and to discuss the results on upper bounds for the coefficient a_{p+3} .

Lemma 1.2 ([1]). *If $w \in \Omega$, then*

$$|w_2 - tw_1^2| \leq \begin{cases} -t & \text{if } t \leq -1, \\ 1 & \text{if } -1 \leq t \leq 1, \\ t & \text{if } t \geq 1. \end{cases}$$

When $t < -1$ or $t > 1$, equality holds if and only if $w(z) = z$ or one of its rotations. If $-1 < t < 1$, then equality holds if and only if $w(z) = z^2$ or one of its rotations. Equality holds for $t = -1$ if and only if $w(z) = \frac{z(\lambda+z)}{1+\lambda z}$, ($0 \leq \lambda \leq 1$) or one of its rotations while for $t = 1$, equality holds if and only if $w(z) = -\frac{z(\lambda+z)}{1+\lambda z}$, ($0 \leq \lambda \leq 1$) or one of its rotations.

Also sharp upper bound above can be improved as follows when $-1 < t < 1$:

$$|w_2 - tw_1^2| + (1+t)|w_1|^2 \leq 1 \quad (-1 < t \leq 0)$$

and

$$|w_2 - tw_1^2| + (1-t)|w_1|^2 \leq 1 \quad (0 < t < 1).$$

Lemma 1.2 is a reformulation of Lemma of Ma and Minda [4].

Lemma 1.3 ([3]). *If $w \in \Omega$, then for any complex number t*

$$|w_2 - tw_1^2| \leq \max\{1, |t|\}.$$

The result is sharp for the functions $w(z) = z$ or $w(z) = z^2$.

Lemma 1.4 ([8]). *If $w \in \Omega$, then for any real numbers q_1 and q_2 the following sharp estimate holds:*

$$|w_3 + q_1 w_1 w_2 + q_2 w_1^3| \leq H(q_1, q_2)$$

where

$$H(q_1, q_2) = \begin{cases} 1 & \text{for } (q_1, q_2) \in D_1 \cup D_2 \\ |q_2| & \text{for } (q_1, q_2) \in \bigcup_{k=3}^7 D_k \\ \frac{2}{3}(|q_1| + 1) \left(\frac{|q_1| + 1}{3(|q_1| + 1 + q_2)} \right)^{\frac{1}{2}} & \text{for } (q_1, q_2) \in D_8 \cup D_9 \\ \frac{q_2}{3} \left(\frac{q_1^2 - 4}{q_1^2 - 4q_2} \right) \left(\frac{q_1^2 - 4}{3(q_2 - 1)} \right)^{\frac{1}{2}} & \text{for } (q_1, q_2) \in D_{10} \cup D_{11} \setminus \{\pm 2, 1\} \\ \frac{2}{3}(|q_1 - 1|) \left(\frac{|q_1| - 1}{3(|q_1| - 1 - q_2)} \right)^{\frac{1}{2}} & \text{for } (q_1, q_2) \in D_{12}. \end{cases}$$

The extremal functions, up to rotations, are of the form

$$w(z) = z^3, \quad w(z) = z, \quad w(z) = w_0(z) = \frac{(z[(1-\lambda)\varepsilon_2 + \lambda\varepsilon_1] - \varepsilon_1\varepsilon_2 z)}{1 - [(1-\lambda)\varepsilon_1 + \lambda\varepsilon_2]z},$$

$$w(z) = w_1(z) = \frac{z(t_1 - z)}{1 - t_1 z}, \quad w(z) = w_2(z) = \frac{z(t_2 + z)}{1 + t_2 z},$$

$$|\varepsilon_1| = |\varepsilon_2| = 1, \quad \varepsilon_1 = t_0 - e^{-\frac{i\theta_0}{2}}(a \mp b), \quad \varepsilon_2 = -e^{-\frac{i\theta_0}{2}}(ia \pm b),$$

$$a = t_0 \cos \frac{\theta_0}{2}, \quad b = \sqrt{1 - t_0^2 \sin^2 \frac{\theta_0}{2}}, \quad \lambda = \frac{b \pm a}{2b},$$

$$t_0 = \left[\frac{2q_2(q_1^2 + 2) - 3q_1^2}{3(q_2 - 1)(q_1^2 - 4q_2)} \right]^{\frac{1}{2}}, \quad t_1 = \left(\frac{|q_1| + 1}{3(|q_1| + 1 + q_2)} \right)^{\frac{1}{2}},$$

$$t_2 = \left(\frac{|q_1| - 1}{3(|q_1| - 1 - q_2)} \right)^{\frac{1}{2}}, \quad \cos \frac{\theta_0}{2} = \frac{q_1}{2} \left[\frac{q_2(q_1^2 + 8) - 2(q_1^2 + 2)}{2q_2(q_1^2 + 2) - 3q_1^2} \right].$$

The sets D_k , $k = 1, 2, \dots, 12$ are defined as follows:

$$\begin{aligned} D_1 &= \left\{ (q_1, q_2) : |q_1| \leq \frac{1}{2}, |q_2| \leq 1 \right\}, \\ D_2 &= \left\{ (q_1, q_2) : \frac{1}{2} \leq |q_1| \leq 2, \frac{4}{27}(|q_1| + 1)^3 - (|q_1| + 1) \leq q_2 \leq 1 \right\}, \\ D_3 &= \left\{ (q_1, q_2) : |q_1| \leq \frac{1}{2}, q_2 \leq -1 \right\}, \\ D_4 &= \left\{ (q_1, q_2) : |q_1| \geq \frac{1}{2}, q_2 \leq -\frac{2}{3}(|q_1| + 1) \right\}, \\ D_5 &= \{(q_1, q_2) : |q_1| \leq 2, q_2 \geq 1\}, \\ D_6 &= \left\{ (q_1, q_2) : 2 \leq |q_1| \leq 4, q_2 \geq \frac{1}{12}(q_1^2 + 8) \right\}, \\ D_7 &= \left\{ (q_1, q_2) : |q_1| \geq 4, q_2 \geq \frac{2}{3}(|q_1| - 1) \right\}, \\ D_8 &= \left\{ (q_1, q_2) : \frac{1}{2} \leq |q_1| \leq 2, \right. \\ &\quad \left. -\frac{2}{3}(|q_1| + 1) \leq q_2 \leq \frac{4}{27}(|q_1| + 1)^3 - (|q_1| + 1) \right\}, \\ D_9 &= \left\{ (q_1, q_2) : |q_1| \geq 2, -\frac{2}{3}(|q_1| + 1) \leq q_2 \leq \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \right\}, \\ D_{10} &= \left\{ (q_1, q_2) : 2 \leq |q_1| \leq 4, \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \leq q_2 \leq \frac{1}{12}(q_1^2 + 8) \right\}, \\ D_{11} &= \left\{ (q_1, q_2) : |q_1| \geq 4, \frac{2|q_1|(|q_1| + 1)}{q_1^2 + 2|q_1| + 4} \leq q_2 \leq \frac{2|q_1|(|q_1| - 1)}{q_1^2 - 2|q_1| + 4} \right\}, \\ D_{12} &= \left\{ (q_1, q_2) : |q_1| \geq 4, \frac{2|q_1|(|q_1| - 1)}{q_1^2 - 2|q_1| + 4} \leq q_2 \leq \frac{2}{3}(|q_1| - 1) \right\}. \end{aligned}$$

2. Coefficient bounds. By making use of Lemmas 1.2–1.4, we prove the following bounds for the class $\mathcal{M}_p(\lambda; \phi)$.

Theorem 2.1. Let $\phi(z) = 1 + B_1 z + B_2 z^2 + \dots$, where B_n 's are real with $B_1 > 0$ and $B_2 \geq 0$, let $0 < \lambda \leq 1$, and

$$\begin{aligned}\sigma_1 &:= \frac{[pB_1^2\lambda + (B_2 - B_1)(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}, \\ \sigma_2 &:= \frac{[pB_1^2\lambda + (B_2 + B_1)(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}, \\ \sigma_3 &:= \frac{[pB_1^2\lambda + B_2(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}.\end{aligned}$$

If $f(z)$ given by (1.1) belongs to $\mathcal{M}_p(\lambda; \phi)$, then

$$(2.1) \quad |a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} -\frac{pB_1\Lambda}{p - p\lambda + 2} & \text{if } \mu \leq \sigma_1 \\ \frac{pB_1}{p - p\lambda + 2} & \text{if } \sigma_1 \leq \mu \leq \sigma_2 \\ \frac{pB_1\Lambda}{p - p\lambda + 2} & \text{if } \mu \geq \sigma_2. \end{cases}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$(2.2) \quad |a_{p+2} - \mu a_{p+1}^2| + \frac{(p - p\lambda + 1)^2}{pB_1(p - p\lambda + 2)}(1 + \Lambda)|a_{p+1}|^2 \leq \frac{pB_1}{p - p\lambda + 2}.$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$(2.3) \quad |a_{p+2} - \mu a_{p+1}^2| + \frac{(p - p\lambda + 1)^2}{pB_1(p - p\lambda + 2)}(1 - \Lambda)|a_{p+1}|^2 \leq \frac{pB_1}{p - p\lambda + 2},$$

where

$$\Lambda = \frac{\mu(p - p\lambda + 2)pB_1^2 - \lambda(p - p\lambda + 1)pB_1^2 - (p - p\lambda + 1)^2B_2}{(p - p\lambda + 1)^2B_1}.$$

For any complex number μ ,

$$(2.4) \quad |a_{p+2} - \mu a_{p+1}^2| \leq \frac{pB_1}{p - p\lambda + 2} \max\{1, |\Lambda|\}$$

Further,

$$(2.5) \quad |a_{p+3}| \leq \frac{pB_1}{p - p\lambda + 3} H(q_1, q_2),$$

where $H(q_1, q_2)$ is as defined in Lemma 1.3,

$$q_1 := \frac{2B_2}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda pB_1}{(p - p\lambda + 1)(p - p\lambda + 2)}$$

and

$$q_2 := \frac{B_3}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_2 - \lambda^2 p^2 B_1^2}{(p - p\lambda + 1)(p - p\lambda + 2)}.$$

These results are sharp.

Proof. If $f(z) \in \mathcal{M}_p(\lambda; \phi)$, then there is an analytic function $w(z)$ given by (1.4) such that

$$(2.6) \quad \frac{\frac{1}{p}zf'(z)}{(1 - \lambda)z^p + \lambda f(z)} = \phi(w(z)).$$

Since

$$\begin{aligned} \frac{\frac{1}{p}zf'(z)}{(1 - \lambda)z^p + \lambda f(z)} &= 1 + \left[\frac{1}{p}(p+1) - \lambda \right] a_{p+1} z + \left[\left(\frac{1}{p}(p+2) - \lambda \right) a_{p+2} \right. \\ &\quad + \left(\lambda^2 - \frac{\lambda}{p}(p+1) \right) a_{p+1}^2 \Big] z^2 + \left[\left(\frac{1}{p}(p+3) - \lambda \right) a_{p+3} \right. \\ &\quad + \left(2\lambda^2 - \frac{\lambda}{p}(p+2) - \frac{\lambda}{p}(p+1) \right) a_{p+1} a_{p+2} \\ &\quad \left. \left. + \left(\frac{1}{p}(p+1)\lambda^2 - \lambda^3 \right) a_{p+1}^3 \right] z^3 + \dots \end{aligned}$$

and

$$\phi(w(z)) = 1 + B_1 w_1 z + (B_1 w_2 + B_2 w_1^2) z^2 + (B_1 w_3 + 2B_2 w_1 w_2 + B_3 w_1^3) z^3 + \dots,$$

we have from (2.6),

$$(2.7) \quad a_{p+1} = \frac{p B_1 w_1}{p - p\lambda + 1},$$

$$(2.8) \quad a_{p+2} = \frac{p(B_1 w_2 + B_2 w_1^2)}{p - p\lambda + 2} + \frac{\lambda p^2 B_1^2 w_1^2}{(p - p\lambda + 1)(p - p\lambda + 2)}$$

and

$$(2.9) \quad \begin{aligned} a_{p+3} &= \frac{p B_1}{p - p\lambda + 3} \left\{ w_3 + \left[\frac{2B_2}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_1}{(p - p\lambda + 1)(p - p\lambda + 2)} \right] w_1 w_2 \right. \\ &\quad \left. + \left[\frac{B_3}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_2 - \lambda^2 p^2 B_1^2}{(p - p\lambda + 1)(p - p\lambda + 2)} \right] w_1^3 \right\}. \end{aligned}$$

Now,

$$(2.10) \quad a_{p+2} - \mu a_{p+1}^2 = \frac{p B_1}{p - p\lambda + 2} \{w_2 - \Lambda w_1^2\}.$$

The results (2.1)–(2.3) are established by an application of Lemma 1.2, inequality (2.4) by Lemma 1.3 and (2.5) follows from Lemma 1.4. To show

that the bounds in (2.1), (2.2) and (2.3) are sharp, we define $K_{\phi n}$, $n = 2, 3, 4, \dots$ by

$$\frac{\frac{1}{p}zK'_{\phi n}(z)}{(1-\lambda)z^p + \lambda K_{\phi n}(z)} = \phi(z^{n-1}), \quad K_{\phi n}(0) = 0 = K'_{\phi n}(0) - 1$$

and the functions F_λ and G_λ , $0 < \lambda \leq 1$ by

$$\frac{\frac{1}{p}zF'_\lambda(z)}{(1-\lambda)z^p + \lambda F_\lambda(z)} = \phi\left(\frac{z(z+\lambda)}{1+\lambda z}\right), \quad F_\lambda(0) = 0 = F'_\lambda(0) - 1$$

and

$$\frac{\frac{1}{p}zG'_\lambda(z)}{(1-\lambda)z^p + \lambda G_\lambda(z)} = \phi\left(-\frac{z(z+\lambda)}{1+\lambda z}\right), \quad G_\lambda(0) = 0 = G'_\lambda(0) - 1.$$

Clearly the functions $K_{\phi n}$, F_λ , $G_\lambda \in \mathcal{M}_p(\lambda; \phi)$.

If $\mu < \sigma_1$ or $\mu > \sigma_2$, then equality holds if and only if f is $K_{\phi 2}$ or one of its rotations. When $\sigma_1 < \mu < \sigma_2$, equality holds if and only if f is $K_{\phi 3}$ or one of its rotations. If $\mu = \sigma_1$ then equality holds if and only if f is F_λ or one of its rotations. Equality holds for $\mu = \sigma_2$ if and only if f is G_λ or one of its rotations. \square

Remark 2.2. For $\lambda = 1$, results (2.1)–(2.4) coincide with the results obtained for the class $S_p^*(\phi)$ by Ali et al. [1].

Remark 2.3. For $\lambda = 1$, $p = 1$, results (2.1)–(2.4) coincide with the results obtained for the class $S^*(\phi)$ by Ma and Minda [4].

Example 2.4. Let $-1 \leq B < A \leq 1$. If $f(z)$ given by (1.1) belongs to $\mathcal{M}_p(\lambda; A, B)$, then

$$|a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} -\frac{p(A-B)\Lambda}{p-p\lambda+2} & \text{if } \mu \leq \sigma_1 \\ \frac{p(A-B)}{p-p\lambda+2} & \text{if } \sigma_1 \leq \mu \leq \sigma_2 \\ \frac{p(A-B)\Lambda}{p-p\lambda+2} & \text{if } \mu \geq \sigma_2. \end{cases}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{(p-p\lambda+1)^2}{p(A-B)(p-p\lambda+2)}(1+\Lambda)|a_{p+1}|^2 \leq \frac{p(A-B)}{p-p\lambda+2}.$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{(p-p\lambda+1)^2}{p(A-B)(p-p\lambda+2)}(1-\Lambda)|a_{p+1}|^2 \leq \frac{p(A-B)}{p-p\lambda+2},$$

where

$$\begin{aligned}\sigma_1 &:= \frac{[p(A-B)\lambda - (1+B)(p-p\lambda+1)](p-p\lambda+1)}{p(p-p\lambda+2)(A-B)}, \\ \sigma_2 &:= \frac{[p(A-B)\lambda + (1-B)(p-p\lambda+1)](p-p\lambda+1)}{p(p-p\lambda+2)(A-B)}, \\ \sigma_3 &:= \frac{[p(A-B)\lambda - B(p-p\lambda+1)](p-p\lambda+1)}{p(p-p\lambda+2)(A-B)}\end{aligned}$$

and

$$\Lambda_A = \frac{\mu p(p-p\lambda+2)(A-B) - (p-p\lambda+1)[(A-B)\lambda p - (p-p\lambda+1)B]}{(p-p\lambda+1)^2}.$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{p(A-B)}{p-p\lambda+2} \max\{1, |\Lambda_A|\}.$$

In particular, if $f \in \mathcal{M}_p(\lambda; \alpha)$, then

$$|a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} -\frac{2p(1-\alpha)\Lambda}{p-p\lambda+2} & \text{if } \mu \leq \sigma_1 \\ \frac{2p(1-\alpha)}{p-p\lambda+2} & \text{if } \sigma_1 \leq \mu \leq \sigma_2 \\ \frac{2p(1-\alpha)\Lambda}{p-p\lambda+2} & \text{if } \mu \geq \sigma_2. \end{cases}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{(p-p\lambda+1)^2}{2p(1-\alpha)(p-p\lambda+2)}(1+\Lambda)|a_{p+1}|^2 \leq \frac{2p(1-\alpha)}{p-p\lambda+2}.$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$|a_{p+2} - \mu a_{p+1}^2| + \frac{(p-p\lambda+1)^2}{2p(1-\alpha)(p-p\lambda+2)}(1-\Lambda)|a_{p+1}|^2 \leq \frac{2p(1-\alpha)}{p-p\lambda+2},$$

where

$$\begin{aligned}\sigma_1 &:= \frac{\lambda(p-p\lambda+1)}{p-p\lambda+2}, \\ \sigma_2 &:= \frac{(p-p\lambda\alpha+1)(p-p\lambda+1)}{p(1-\alpha)(p-p\lambda+2)}, \\ \sigma_3 &:= \frac{[2p(1-\alpha)\lambda + (p-p\lambda+1)](p-p\lambda+1)}{2p(1-\alpha)(p-p\lambda+2)}\end{aligned}$$

and

$$\Lambda_\alpha = \frac{2\mu p(1-\alpha)(p-p\lambda+2) - (p-p\lambda+1)[2\lambda p(1-\alpha) + (p-p\lambda+1)]}{(p-p\lambda+1)^2}.$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{2p(1-\alpha)}{p-p\lambda+2} \max\{1, |\Lambda_\alpha|\}.$$

The results are sharp.

Corollary 2.5. *Let $\phi(z)$ be as in Theorem 2.1,*

$$g(z) = z^p + \sum_{n=1}^{\infty} g_{p+n} z^{p+n} \quad (g_{p+n} > 0),$$

and let

$$\begin{aligned} \sigma_1 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{[pB_1^2\lambda + (B_2 - B_1)(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}, \\ \sigma_2 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{[pB_1^2\lambda + (B_2 + B_1)(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}, \\ \sigma_3 &:= \frac{g_{p+1}^2}{g_{p+2}} \frac{[pB_1^2\lambda + B_2(p - p\lambda + 1)](p - p\lambda + 1)}{(p - p\lambda + 2)pB_1^2}. \end{aligned}$$

If $f(z)$ given by (1.1) belongs to $\mathcal{M}_{p,g}(\lambda; \phi)$, then

$$(2.11) \quad |a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} -\frac{pB_1\Lambda}{g_{p+2}(p - p\lambda + 2)} & \text{if } \mu \leq \sigma_1 \\ \frac{pB_1}{g_{p+2}(p - p\lambda + 2)} & \text{if } \sigma_1 \leq \mu \leq \sigma_2 \\ \frac{pB_1\Lambda}{g_{p+2}(p - p\lambda + 2)} & \text{if } \mu \geq \sigma_2. \end{cases}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$(2.12) \quad \begin{aligned} |a_{p+2} - \mu a_{p+1}^2| + \frac{g_{p+1}^2(p - p\lambda + 1)^2}{g_{p+2}(p - p\lambda + 2)pB_1} (1 + \Lambda) |a_{p+1}|^2 \\ \leq \frac{pB_1}{g_{p+2}(p - p\lambda + 2)}. \end{aligned}$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$(2.13) \quad \begin{aligned} |a_{p+2} - \mu a_{p+1}^2| + \frac{g_{p+1}^2(p - p\lambda + 1)^2}{g_{p+2}(p - p\lambda + 2)pB_1} (1 - \Lambda) |a_{p+1}|^2 \\ \leq \frac{pB_1}{g_{p+2}(p - p\lambda + 2)}, \end{aligned}$$

where

$$\Lambda_g = \frac{\frac{g_{p+2}}{g_{p+1}^2} \mu(p - p\lambda + 2)pB_1^2 - \lambda(p - p\lambda + 1)pB_1^2 - (p - p\lambda + 1)^2 B_2}{(p - p\lambda + 1)^2 B_1}.$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{pB_1}{g_{p+2}(p-p\lambda+2)} \max\{1, |\Lambda_g|\}.$$

Further,

$$|a_{p+3}| \leq \frac{pB_1}{g_{p+3}(p-p\lambda+3)} H(q_1, q_2),$$

where $H(q_1, q_2)$ is as defined in Lemma 1.3,

$$q_1 := \frac{2B_2}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_1}{(p - p\lambda + 1)(p - p\lambda + 2)}$$

and

$$q_2 := \frac{B_3}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_2 - \lambda^2 p^2 B_1^2}{(p - p\lambda + 1)(p - p\lambda + 2)}.$$

These results are sharp.

3. Applications to functions defined by extended fractional differ-integrals. With a view to define fractional differintegral operator $\Omega_z^{(\delta,p)}$, we recall Gauss hypergeometric function ${}_2F_1$ defined by [6].

$$(3.1) \quad {}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!} \quad (a, b, c \in \mathbb{C}, c \neq 0, -1, -2, \dots),$$

where $(d)_n$ denotes the Pochhammer symbol given in terms of Gamma function Γ by

$$(d)_n = \frac{\Gamma(d+n)}{\Gamma(d)} = \begin{cases} 1 & (n=0; d \in \mathbb{C} \setminus \{0\}) \\ d(d+1)\dots(d+n-1) & (n \in \mathbb{N}; d \in \mathbb{C}). \end{cases}$$

We note that the series defined by (3.1) converges absolutely for $z \in \Delta$ and hence ${}_2F_1$ represents an analytic function in Δ .

Also we recall the definitions of fractional calculus considered by Owa [5] (see also [6, 11, 12]).

Definition 3.1. The fractional integral of order δ ($\delta > 0$) is defined, for a function f , analytic in a simply connected region of the complex plane containing the origin, by

$$(3.2) \quad D_z^{-\delta} f(z) = \frac{1}{\Gamma(\delta)} \int_0^z \frac{f(\zeta)}{(z-\zeta)^{1-\delta}} d\zeta,$$

where the multiplicity of $(z-\zeta)^{\delta-1}$ is removed by requiring $\log(z-\zeta)$ to be real when $z-\zeta > 0$.

Definition 3.2. Under the hypothesis of Definition 3.1, the fractional derivative of f of order δ ($\delta \geq 0$) is defined by

$$(3.3) \quad D_z^\delta f(z) = \begin{cases} \frac{1}{\Gamma(1-\delta)} \frac{d}{dz} \int_0^z \frac{f(\zeta)}{(z-\zeta)^\delta} d\zeta, & (0 \leq \delta < 1) \\ \frac{d^n}{dz^n} D_z^{\delta-n} f(z), & (n \leq \delta < n+1; n \in \mathbb{N}_0) \end{cases}$$

where $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ and the multiplicity of $(z - \zeta)^{-\delta}$ is removed as in Definition 3.1.

Definition 3.3 ([7]). The extended fractional differintegral operator $\Omega_z^{(\delta,p)} : \mathcal{A}_p \rightarrow \mathcal{A}_p$ for a function f of the form (1.1) and for a real number δ ($-\infty < \delta < p+1$) is defined by

$$(3.4) \quad \Omega_z^{(\delta,p)} f(z) = \frac{\Gamma(p+1-\delta)}{\Gamma(p+1)} z^\delta D_z^\delta f(z) \quad (-\infty < \delta < p+1; z \in \Delta),$$

where $D_z^\delta f(z)$ is respectively, the fractional integral of f of order $-\delta$ when $-\infty < \delta < 0$ and the fractional derivative of f of order δ when $0 \leq \delta < p+1$.

We note that

$$\begin{aligned} \Omega_z^{(\delta,p)} f(z) &= z^p + \sum_{n=1}^{\infty} \frac{\Gamma(n+p+1)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(n+p+1-\delta)} a_{n+p} z^{n+p} \\ &= z^p {}_2F_1(1, p+1; p+1-\delta; z) * f(z) \end{aligned}$$

$(-\infty < \delta < p+1; z \in \Delta)$.

Let $\mathcal{M}_{p,\delta}(\lambda; \phi)$ be the class of functions $f \in \mathcal{A}_p$ for which $\Omega_z^{(\delta,p)} f(z) \in \mathcal{M}_p(\lambda; \phi)$. The class $\mathcal{M}_{p,\delta}(\lambda; \phi)$ is the special case of the class $\mathcal{M}_{p,g}(\lambda; \phi)$, when

$$g(z) = z^p + \sum_{n=1}^{\infty} \frac{\Gamma(n+p+1)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(n+p+1-\delta)} z^{n+p}.$$

Since

$$(\Omega_z^{(\delta,p)} f)(z) = z^p + \sum_{n=1}^{\infty} \frac{\Gamma(n+p+1)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(n+p+1-\delta)} a_{n+p} z^{n+p},$$

we have

$$(3.5) \quad g_{p+1} = \frac{\Gamma(p+2)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(p+2-\delta)} = \frac{p+1}{p+1-\delta},$$

$$(3.6) \quad g_{p+2} = \frac{\Gamma(p+3)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(p+3-\delta)} = \frac{(p+1)(p+2)}{(p+1-\delta)(p+2-\delta)},$$

$$(3.7) \quad g_{p+3} = \frac{\Gamma(p+4)\Gamma(p+1-\delta)}{\Gamma(p+1)\Gamma(p+4-\delta)} = \frac{(p+1)(p+2)(p+3)}{(p+1-\delta)(p+2-\delta)(p+3-\delta)}.$$

For g_{p+1}, g_{p+2} and g_{p+3} given by (3.5), (3.6) and (3.7), Corollary 2.5 reduces to the following:

Theorem 3.4. *Let $\phi(z)$ be as in Theorem 2.1, and let*

$$\begin{aligned}\sigma_1 &:= \frac{(p+1)(p+2-\delta)}{(p+2)(p+1-\delta)} \frac{[pB_1^2\lambda + (B_2 - B_1)(p-p\lambda+1)](p-p\lambda+1)}{(p-p\lambda+2)pB_1^2}, \\ \sigma_2 &:= \frac{(p+1)(p+2-\delta)}{(p+2)(p+1-\delta)} \frac{[pB_1^2\lambda + (B_2 + B_1)(p-p\lambda+1)](p-p\lambda+1)}{(p-p\lambda+2)pB_1^2}, \\ \sigma_3 &:= \frac{(p+1)(p+2-\delta)}{(p+2)(p+1-\delta)} \frac{[pB_1^2\lambda + B_2(p-p\lambda+1)](p-p\lambda+1)}{(p-p\lambda+2)pB_1^2}.\end{aligned}$$

If $f(z)$ given by (1.1) belongs to $\mathcal{M}_{p,\delta}(\lambda; \phi)$, then

$$|a_{p+2} - \mu a_{p+1}^2| \leq \begin{cases} -\frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1\Lambda}{(p-p\lambda+2)} & \text{if } \mu \leq \sigma_1, \\ \frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1}{(p-p\lambda+2)} & \text{if } \sigma_1 \leq \mu \leq \sigma_2, \\ \frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1\Lambda}{(p-p\lambda+2)} & \text{if } \mu \geq \sigma_2. \end{cases}$$

Further, if $\sigma_1 \leq \mu \leq \sigma_3$, then

$$\begin{aligned}(3.8) \quad |a_{p+2} - \mu a_{p+1}^2| &+ \frac{(p+1)(p+2-\delta)}{(p+2)(p+1-\delta)} \frac{(p-p\lambda+1)^2}{(p-p\lambda+2)} (1+\Lambda) |a_{p+1}|^2 \\ &\leq \frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1}{(p-p\lambda+2)}.\end{aligned}$$

If $\sigma_3 \leq \mu \leq \sigma_2$, then

$$\begin{aligned}(3.9) \quad |a_{p+2} - \mu a_{p+1}^2| &+ \frac{(p+1)(p+2-\delta)}{(p+2)(p+1-\delta)} \frac{(p-p\lambda+1)^2}{(p-p\lambda+2)} (1+\Lambda) |a_{p+1}|^2 \\ &\leq \frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1}{(p-p\lambda+2)},\end{aligned}$$

where

$$\Lambda_\delta = \frac{\frac{(p+2)(p+1-\delta)}{(p+1)(p+2-\delta)} \mu (p-p\lambda+2)pB_1^2 - \lambda (p-p\lambda+1)pB_1^2 - (p-p\lambda+1)^2 B_2}{(p-p\lambda+1)^2 B_1}.$$

For any complex number μ ,

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{(p+1-\delta)(p+2-\delta)}{(p+1)(p+2)} \frac{pB_1}{(p-p\lambda+2)} \max\{1, |\Lambda_\delta|\}.$$

Further,

$$|a_{p+3}| \leq \frac{(p+1-\delta)(p+2-\delta)(p+3-\delta)}{(p+1)(p+2)(p+3)} \frac{pB_1}{(p-p\lambda+3)} H(q_1, q_2),$$

where

$$q_1 := \frac{2B_2}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_1}{(p - p\lambda + 1)(p - p\lambda + 2)}$$

and

$$q_2 := \frac{B_3}{B_1} - \frac{(2p\lambda - 2p - 3)\lambda p B_2 - \lambda^2 p^2 B_1^2}{(p - p\lambda + 1)(p - p\lambda + 2)}.$$

These results are sharp.

Acknowledgements. The authors thank the referee for his insightful suggestions.

REFERENCES

- [1] Ali, R. M., Ravichandran, V., Seenivasagan, N., *Coefficient bounds for p -valent functions*, Appl. Math. Comput. **187** (2007), 35–46.
- [2] Janowski, W., *Some extremal problems for certain families of analytic functions*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **21** (1973), 17–25.
- [3] Keogh, F. R., Merkes, E. P., *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12.
- [4] Ma, W. C., Minda, D., *A unified treatment of some special classes of univalent functions*, Proceedings of the Conference on Complex Analysis (Tianjin, 1992), Int. Press, Cambridge, MA, 1994, 157–169.
- [5] Owa, S., *On the distortion theorem. I*, Kyungpook Math. J. **18** (1) (1978), 53–59.
- [6] Owa, S., Srivastava, H. M., *Univalent and starlike generalized hypergeometric functions*, Canad. J. Math. **39** (5) (1987), 1057–1077.
- [7] Patel, J., Mishra, A., *On certain subclasses of multivalent functions associated with an extended differintegral operator*, J. Math. Anal. Appl. **332** (2007), 109–122.
- [8] Prokhorov, D. V., Szymal, J., *Inverse coefficients for (α, β) -convex functions*, Ann. Univ. Mariae Curie-Skłodowska Sect. A **35** (1981), 125–143.
- [9] Selvaraj, C., Selvakumaran, K. A., *Fekete–Szegő problem for some subclass of analytic functions*, Far East J. Math. Sci. (FJMS) **29** (3) (2008), 643–652.
- [10] Srivastava, H. M., Mishra, A. K., Das, M. K., *The Fekete–Szegő problem for a subclass of close-to-convex functions*, Complex Variables Theory Appl. **44** (2) (2001), 145–163.
- [11] Srivastava, H. M., Owa, S., *An application of the fractional derivative*, Math. Japon. **29** (3) (1984), 383–389.
- [12] Srivastava, H. M., Owa, S., *Univalent Functions, Fractional Calculus and their Applications*, Halsted Press/John Wiley & Sons, Chichester–New York, 1989.

C. Selvaraj
 Department of Mathematics
 Presidency College (Autonomous)
 Chennai - 600005
 India
 e-mail: pamc9439@yahoo.co.in

O. S. Babu
 Department of Mathematics
 Dr. Ambedkar Govt. Arts College
 Chennai - 600039
 India
 e-mail: osbabu1009@gmail.com

G. Murugusundaramoorthy
School of Advanced Sciences
VIT university
Vellore - 632 014
India
e-mail: gmsmoorthy@yahoo.com

Received June 25, 2012

Copyright of Annales UMCS, Mathematica is the property of De Gruyter Open and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.