

EQUALITY OF GRAPHOIDAL AND ACYCLIC GRAPHOIDAL COVERING NUMBER OF A GRAPH

INDRA RAJASINGH AND P. ROUSHINI LEELY PUSHPAM

Abstract. A *graphoidal cover* of a graph G is a collection ψ of (not necessarily open) paths in G such that every vertex of G is an internal vertex of at most one path in ψ and every edge of G is in exactly one path in ψ . If no member of ψ is a cycle, then ψ is called an *acyclic graphoidal cover* of G . The minimum cardinality of a graphoidal cover is called the *graphoidal covering number* of G and is denoted by η and the minimum cardinality of an acyclic graphoidal cover is called an *acyclic graphoidal covering number* of G and is denoted by η_a . In this paper we characterize the class of graphs for which $\eta = \eta_a$.

1. Introduction

The concept of graphoidal cover was introduced by B.D. Acharya and E. Sampathkumar in 1987 and a study of the graphoidal covering number was initiated by them [1]. Since then this area of research has been explored by several authors [2, 3, 10, 11, 12, 13, 19]. The concept of an acyclic graphoidal cover was introduced by Suresh Suseela and pursued by Arumugam *et al.*, [14, 15].

By a *graph* $G = (V, E)$ we mean a finite, undirected, connected graph without loops or multiple edges. The order and size of G are denoted by p and q respectively. For graph theoretic terminology we refer to Harary [16].

If $P = (v_0, v_1, \dots, v_n)$ is a path or a cycle in G , v_1, \dots, v_{n-1} are called *internal vertices* of P . If $P = (v_0, v_1, \dots, v_n)$ and $Q = (v_n = w_0, w_1, \dots, w_m)$ are two paths in G then the walk obtained by concatenating P and Q at v_n is denoted by $P \circ Q$.

Definition 1.1. A *graphoidal cover* of a graph G is a set ψ of (not necessarily open) paths in G satisfying the following conditions.

- (i) Every path in ψ has at least two vertices.
- (ii) Every vertex of G is an internal vertex of at most one path in ψ .
- (iii) Every edge of G is in exactly one path in ψ .

ψ is called an *acyclic graphoidal cover* of G if no member of ψ is a cycle in G . The minimum cardinality of (an acyclic) a graphoidal cover of G is called the (*acyclic*) *graphoidal covering number* of G and is denoted by (η_a) η .

Received February 13, 2003; revised May 24, 2004.

2000 *Mathematics Subject Classification.* 05C.

Key words and phrases. Graphoidal covering number, acyclic graphoidal covering number.

Definition 1.2. Let ψ be a collection of internally disjoint paths in G . A vertex of G is said to be an *interior vertex* of ψ if it is an internal vertex of some path in ψ . Any vertex which is not an interior vertex of ψ is said to be an *exterior vertex* of ψ .

Theorem 1.3.([18]) For any graphoidal cover ψ of G , let t_ψ denote the number of exterior vertices of ψ . Let $t = \min t_\psi$, where the minimum is taken over all graphoidal covers of G . Then $\eta = q - p + t$ where p and q denote respectively the order and size of G .

Corollary 1.4.([18]) For any graph G , $\eta \geq q - p$. Moreover the following are equivalent.

- (i) $\eta = q - p$.
- (ii) There exists a graphoidal cover without exterior vertices.
- (iii) There exists a set of internally disjoint and edge disjoint paths without exterior vertices.

Theorem 1.5.([17]) For any graph G with $\delta \geq 3$, $\eta = q - p$

Remark 1.6. Results analogous to Theorem 1.3, Corollary 1.4 and Theorem 1.5 are also true for an acyclic graphoidal covering number of a graph.

Theorem 1.7.([15]) For any graph with $\delta \geq 3$, $\eta = \eta_a$.

For graphs with $\delta \leq 2$ it is not necessary that $\eta = \eta_a = q - p$. Hence the problem of characterizing graphs with $\delta \leq 2$ satisfying $\eta = \eta_a$ is challenging. In this paper we solve this problem. For our further discussion we confine ourselves to graphs with $\delta \leq 2$. We need the following definition and theorems.

Definition 1.8.([7]) Let $\mathcal{G}(f)$ denote the collection of all blocks whose edge set can be decomposed into a cycle C and a collection \wp of internally disjoint paths such that each path P in \wp has $f \in V(C)$ as its origin and $|V(P) \cap V(C)| \leq 2$ (The collection \wp may be empty in which case the corresponding member of $\mathcal{G}(f)$ is a cycle). We observe that if $G \in \mathcal{G}(f)$ and G is not a cycle, then $\deg f = |\wp| + 2 = \Delta$ and there is at most one vertex $v \neq f$ with $\deg v = \Delta$.

Let \mathcal{F} and \mathcal{F}_a denote respectively the class of all graphs G with $\eta = q - p$ and $\eta_a = q - p$.

Theorem 1.9.([4]) Let G be a 2-connected graph with $p \geq 3$. Then $G \notin \mathcal{F}$ if and only if G is a cycle or a cycle with exactly one chord or a theta graph.

Theorem 1.10.([4]) Let G be a 2-edge connected graph with $\delta = 2$. Then $G \notin \mathcal{F}$ if and only if every block of G is a cycle or a cycle with exactly one chord or a theta graph and at most one block of G is not a cycle.

Theorem 1.11.([6]) Let G be a connected graph with $\delta = 2$ and edge connectivity one. Then $G \notin \mathcal{F}$ if and only if there exists a cut edge e of G such that at least one component of $G - e$ is a graph, all of whose blocks are cycles.

Lemma 1.12.([6]) *Let G be a 2-edge connected graph such that exactly one block of G is either a cycle with exactly one chord or a theta graph and all other blocks are cycles. Let $v \in V(G)$. Then there exists a minimum graphoidal cover ψ of G such that v is the only vertex exterior to ψ and there exists a path in ψ which contains v as an exterior vertex.*

Theorem 1.13.([7]) *Let G be a 2-connected graph with $p \geq 3$. Then $G \in \mathcal{F}_a$ if and only if $G \notin \mathcal{G}(f)$.*

Theorem 1.14.([7]) *Let G be a graph with $\delta > 1$ and connectivity one. Then $G \notin \mathcal{F}_a$ if and only if at least one end block of G is a member of $\mathcal{G}(f)$ with f as a cut vertex of G .*

Theorem 1.15.([5]) *Let $G \in \mathcal{G}(f)$ and $\Delta(G) \geq 3$. Let v be a vertex of G with $\deg v \neq \Delta$. Then there exists a minimum acyclic graphoidal cover ψ of G such that v is the only vertex exterior to ψ and $\eta_a = |\psi| = \Delta - 1$.*

2. Main Results

Lemma 2.1. *Let G be a graph whose blocks are all cycles. Let $u, v \in V(G)$ with $u \neq v$. Then there exists a graphoidal cover ψ of G such that u and v are the only vertices exterior to ψ and a path P in ψ which contains u and v as end vertices.*

Proof. Let C_1, C_2, \dots, C_k be the blocks of G . By hypothesis each C_i is a cycle, $i = 1, 2, \dots, k$. Let P be a (u, v) -path in G . Let S_1, S_2, \dots, S_s , $s \in \{1, 2, \dots, k\}$ be the segments of P that are part of the cycles C_1, C_2, \dots, C_s . Let $R_i = C_i \setminus S_i$, $1 \leq i \leq s$. Then $\psi = \{P, R_1, R_2, \dots, R_s, C_{s+1}, \dots, C_k\}$ is a graphoidal cover of G with u and v as the only vertices exterior to ψ .

Theorem 2.2. *Let G be a 2-edge connected graph. Then either $\eta = q - p$ or $\eta = q - p + 1$.*

Proof. Suppose $\eta \neq q - p$. By Theorem 1.10, every block of G is a cycle or a cycle with exactly one chord or a theta graph and at most one block of G is not a cycle. We now prove that $\eta = q - p + 1$ by induction on m where m is the number of blocks of G . When $m = 1$, G is either a cycle or a cycle with exactly one chord or a theta graph. Clearly for any minimum graphoidal cover ψ of G , exactly one vertex of G is exterior to ψ and hence by Theorem 1.3, $\eta = q - p + 1$.

We now assume that $\eta = q - p + 1$ for all 2-edge connected graphs with m blocks, $m \geq 1$. Let G be a 2-edge connected graph with $m + 1$ blocks. Let C be an end block which is a cycle and $v \in V(C)$ be a cut vertex of G . Let G' be the subgraph of G obtained by removing all the vertices of $C - v$. Clearly G' has m blocks. By induction

hypothesis $\eta(G') = q' - p' + 1$ when p' and q' are the order and size of G' respectively.

$$\begin{aligned} \text{Now } \eta(G) &= \eta(G') + 1 \\ &= q' - p' + 2 \\ &= q - p + 1. \text{ This completes the induction and the proof.} \end{aligned}$$

Theorem 2.3. *Let G be a graph with $\kappa' = 1$ and $\delta > 1$, where κ' is the edge connectivity of G . Let $S = \{e \mid e \text{ is a cut edge of } G \text{ and the blocks of at least one component of } G-e \text{ are all cycles}\}$. Let $\mathcal{H}_G = \{H_1, H_2, \dots, H_m\}$, be the collection of all such components. Let $v_i \in V(H_i)$, $1 \leq i \leq m$. Then $\eta = q - p + m$ and there exists a minimum graphoidal cover ψ of G such that v_i , $1 \leq i \leq m$ are the only vertices exterior to ψ and there exist paths in ψ which contain v_i as an end vertex, $1 \leq i \leq m$.*

Proof. Since contracting an edge incident with a cut vertex of degree 2 does not affect the value of $q - p$ and η , we may assume without loss of generality that any cut vertex has degree at least 3. We observe that if $m = 0$, then the result follows from Theorem 1.11. Hence we assume that $m > 0$ and prove the result by induction on r , where r is the number of cut edges of G . Since $\kappa' = 1$, $r \geq 1$. Suppose $r = 1$ and let $e = x_1x_2$ be the cut edge of G . Let G_1 and G_2 be the components of $G-e$ with $x_1 \in V(G_1)$ and $x_2 \in V(G_2)$. Clearly G_1 and G_2 are 2-edge connected graphs. If both G_1 and G_2 are in \mathcal{H}_G then by Lemma 2.1 there exists a graphoidal cover ψ_i of G_i , $i = 1, 2$ and a (v_i, x_i) -path P_i in G_i such that v_i and x_i are the only vertices exterior to ψ_i , $i = 1, 2$. Then by Theorem 1.3 $\psi = (\psi_1 \setminus \{P_1\}) \cup (\psi_2 \setminus \{P_2\}) \cup \{P_1 \circ e \circ P_2^{-1}\}$ is a minimum graphoidal cover such that v_1 and v_2 are the only vertices exterior to ψ . Thus $\eta = q - p + 2$. Suppose $G_1 \notin \mathcal{H}_G$. For $G_2 \in \mathcal{H}_G$ we define ψ_2 as before. If $G_1 \in \mathcal{F}$, then there exists a minimum graphoidal cover ψ_1 of G_1 such that all vertices of G_1 are interior to ψ_1 . Then by Theorem 1.3 $\psi = \psi_1 \cup (\psi_2 \setminus \{P_2\}) \cup (e \circ P_2^{-1})$ is a minimum graphoidal cover of G with v_2 as the only vertex exterior to ψ , where $e \circ P_2^{-1}$ is a path in ψ which contains v_2 as an end vertex. If $G_1 \notin \mathcal{F}$, then since $G_1 \notin \mathcal{H}_G$, by Theorem 1.10 exactly one block of G_1 is a cycle with exactly one chord or a theta graph and all other blocks are cycles. Hence by Lemma 1.12 there exists a minimum graphoidal cover ψ_1 of G_1 and a path P_1 in ψ_1 with x_1 as its terminus such that x_1 is the only vertex exterior to ψ_1 . Then $\psi = (\psi_1 \setminus \{P_1\}) \cup (\psi_2 \setminus \{P_2\}) \cup \{P_1 \circ e \circ P_2^{-1}\}$ is a minimum graphoidal cover of G such that v_2 is the only vertex exterior to ψ and $P_1 \circ e \circ P_2^{-1}$ is a path in ψ which contains v_2 as an end vertex. Thus $\eta = q - p + 1$. Hence the result is true for $r = 1$.

We now assume that the result is true for all graphs with at most $r - 1$ cut edges. Let G be a graph with r cut edges satisfying the conditions of the theorem. Let $e = x_1y_1$ be a cut edge of G such that one of the components of $G - e$ is H_1 . Let G_1 be the other component of $G - e$ and let $x_1 \in V(G_1)$ and $y_1 \in V(H_1)$. By Lemma 2.1 there exists a graphoidal cover ψ_1 of H_1 whose only exterior vertices are y_1 and v_1 and there exists a (y_1, v_1) -path Q_1 in ψ_1 . Clearly $\delta(G_1) > 1$.

Case(i). $x_1 \in H$ for some $H \in \mathcal{H}_{G_1}$

By induction hypothesis there exists a minimum graphoidal cover ψ_2 of G_1 whose only exterior vertices are x_1, v_2, \dots, v_m and there exists a path R_1 in ψ_2 with x_1 as its terminus and path in ψ_2 with v_1 as terminus, $2 \leq i \leq m$. Then $\psi = (\psi_1 \setminus \{Q_1\}) \cup (\psi_2 \setminus \{R_1\}) \cup \{P_1\}$ where $P_1 = R_1 \circ e \circ Q_1$ is the required minimum graphoidal cover of G .

Case(ii). $x_1 \notin H$ for all $H \in \mathcal{H}_{G_1}$

By induction hypothesis there exists a minimum graphoidal cover ψ_2 of G_1 whose only exterior vertices are v_2, v_3, \dots, v_m and there exists a path ψ_2 with v_i , $2 \leq i \leq m$ as terminus. Now $\psi = (\psi_1 \setminus \{Q_1\}) \cup \psi_2 \cup \{e \circ Q_1\}$ is the required minimum graphoidal cover of G . This completes the induction and the proof.

Theorem 2.4. *Let G be a connected graph which is not a tree with n pendant vertices, $n \geq 1$ and $|\mathcal{H}_G| = m$, $m \geq 0$. Then $\eta = q - p + m + n$.*

Proof. We prove the result by induction on n , the number of pendant vertices in G . Let $\mathcal{H}_G = \{H_1, H_2, \dots, H_m\}$. Let $v_i \in V(H_i)$, $1 \leq i \leq m$ where $H_i \in \mathcal{H}_G$. Suppose $n = 1$. Let $v \in V(G)$ be such that $\deg v = 1$ and $P = (v, u_1, u_2, \dots, u_k, w)$ be a path in G such that $\deg u_i = 2$, for all i , $1 \leq i \leq k$ and $\deg w > 2$. Such a vertex w exists because G is not a path. Let $G_1 = G \setminus \{v, u_1, u_2, \dots, u_k\}$.

Case(i). $w \in H$ for some $H \in \mathcal{H}_{G_1}$

Then by Theorem 2.3, there exists a minimum graphoidal cover ψ_1 of G_1 such that v_1, v_2, \dots, v_m, w are the only vertices exterior to ψ_1 and there exists paths in ψ_1 which contains v_i , $1 \leq i \leq m$ and w as end vertices. Let P_1 be the path in ψ_1 with w as its terminus. Then $\psi = (\psi_1 \setminus \{P_1\}) \cup \{P_1 \circ P^{-1}\}$ is a minimum graphoidal cover of G such that v_1, v_2, \dots, v_m, v are the only vertices exterior to ψ and $\eta = q - p + m + 1$.

Case(ii). $w \notin H$ for all $H \in \mathcal{H}_{G_1}$

By Theorem 2.3, there exists a minimum graphoidal cover ψ_1 of G_1 such that v_1, v_2, \dots, v_m are the only vertices exterior to ψ_1 . Hence $\psi = \psi_1 \cup \{P\}$ is a minimum graphoidal cover with v_1, v_2, \dots, v_m and v as the only vertices exterior to ψ . Hence $\eta = q - p + m + 1$. Therefore the result is true for $n = 1$.

We assume the result is true for k pendant vertices. Let G be a graph with $k + 1$ pendant vertices and let z_1, z_2, \dots, z_{k+1} be the pendant vertices of G . Let $P = (z_1, u_1, \dots, u_k, w)$ be a path in G such that $\deg u_i = 2$, for all i , $1 \leq i \leq k$ and $\deg w > 2$. Let $G_1 = G \setminus \{z_1, u_1, u_2, \dots, u_k\}$. If $w \in H$ for some $H \in \mathcal{H}_{G_1}$, then by induction hypothesis there exists a minimum graphoidal cover ψ_1 of G_1 such that $v_1, v_2, \dots, v_m, w, z_2, z_3, \dots, z_{k+1}$ are the only vertices exterior to ψ_1 and there exists a path P_1 in ψ_1 with w as its terminus. Hence $\psi = (\psi_1 \setminus \{P_1\}) \cup \{P_1 \circ P^{-1}\}$ is a minimum graphoidal cover of G with $z_1, z_2, \dots, z_{k+1}, v_1, v_2, \dots, v_m$ as the only vertices exterior to ψ . Thus $\eta = q - p + n + m$. If $w \notin H$ for all $H \in \mathcal{H}_G$, then by induction hypothesis there exists a minimum graphoidal cover ψ_1 of G_1 such that $v_1, v_2, \dots, v_m, z_2, \dots, z_{k+1}$ are the only vertices exterior to ψ_1 . Hence $\psi = \psi_1 \cup \{P\}$ is a minimum graphoidal cover

of G with v_1, v_2, \dots, v_m and z_1, z_2, \dots, z_{k+1} as the only vertices exterior to ψ . Thus $\eta = q - p + m + n$.

The next theorem determines the acyclic graphoidal covering number of a graph with $\delta = 2$.

Theorem 2.5. *Let G be a graph with $\delta = 2$. Let B_1, B_2, \dots, B_m , $m \geq 0$ be end blocks of G which are in $\mathcal{G}(f_i)$ with f_i as a cut vertex. Let $v_i \in V(B_i)$, $1 \leq i \leq m$ and v_i is not a cut vertex of G . Then there exists a minimum acyclic graphoidal cover ψ of G whose only exterior vertices are v_1, v_2, \dots, v_m and $\eta_a = q - p + m$.*

Proof. We observe that if $m = 0$, then the result follows from Theorem 1.13 and Theorem 1.14. Hence we assume that $m > 0$ and prove the result by induction on n , where n is the number of blocks of G . If $n = 1$, the result follows from Theorem 1.15. We now assume that the result is true for all graphs with at most $n - 1$ blocks. Let G be a graph with n blocks satisfying the conditions of the theorem. Let ψ_1 be a minimum acyclic graphoidal cover of B_1 whose only exterior vertex is v_1 . Let $H = G \setminus (V(B_1) \setminus \{f_1\})$.

Case(i). $\deg_H f_1 = 1$.

We choose a path $P = (f_1, u_1, \dots, u_k, w)$, such that $\deg u_i = 2$ for each i and $\deg w > 2$. Let $G_1 = H \setminus \{f_1, u_1, u_2, \dots, u_k\}$. Clearly $\delta(G_1) > 1$. Let B be a block of G_1 containing w . If B is an end block of G_1 and $B \in \mathcal{G}(f)$ with f a cut vertex of G , then by induction hypothesis there exists a minimum acyclic graphoidal cover ψ_2 of G_1 whose only exterior vertices are w, v_2, \dots, v_m . Let P_1 be a path in ψ_2 having w as its terminus. Then $\psi = (\psi_2 \setminus \{P_1\}) \cup \{P_1 \circ P^{-1}\} \cup \psi_1$ is a minimum acyclic graphoidal cover of G with v_1, v_2, \dots, v_m as its only exterior vertices.

Otherwise by induction hypothesis there exists a minimum acyclic graphoidal cover ψ_2 of G_1 whose only exterior vertices are v_2, v_3, \dots, v_m and $\psi = \psi_2 \cup \{P\} \cup \psi_1$ is a minimum acyclic graphoidal cover of G whose only exterior vertices are v_1, v_2, \dots, v_m .

Case(ii). $\deg_H f_1 > 1$.

Let B be a block of H containing f_1 . If B is an end block of G and $B \in \mathcal{G}(f)$ with f as a cut vertex then by induction hypothesis there exists a minimum acyclic graphoidal cover ψ_2 of H with v_2, v_3, \dots, v_m and f_1 as its only exterior vertices. Then $\psi = \psi_1 \cup \psi_2$ is the required minimum acyclic graphoidal cover of G .

Otherwise by induction hypothesis there exists a minimum acyclic graphoidal cover ψ_2 of H with v_2, v_3, \dots, v_m as its only exterior vertices. Let P be the path in ψ_2 having f_1 as an internal vertex. Let x, y be the terminal vertices of P . Let P_1 and P_2 be the (x, f_1) and (f_1, y) - sections of P respectively. Then $\psi = \psi_1 \cup (\psi_2 \setminus \{P\}) \cup \{P_1, P_2\}$ is the required minimum acyclic graphoidal cover of G . This completes the induction and the proof.

Theorem 2.6. *Let G be a graph with n pendant vertices, $n \geq 1$ and let B_1, B_2, \dots, B_m , $m \geq 0$ be end blocks of G which are in $\mathcal{G}(f_i)$ with f_i as a cut vertex. Then $\eta_a = q - p + m + n$.*

Proof. Similar to the proof of Theorem 2.3.

We now proceed to the main theorem of characterizing the class of graphs with $\eta = \eta_a$.

Remark 2.7. Since $\eta \leq \eta_a$, $\eta_a = q - p$ implies that $\eta = q - p$. Theorem 1.13 and Theorem 1.14 characterizes the class of all graphs for which $\eta_a = q - p$. Hence for these graphs $\eta = q - p$, in turn $\eta = \eta_a$. Hence we need to consider the case when $\eta_a \neq q - p$.

Theorem 2.8. *Let G be a connected graph with $\eta_a \neq q - p$ and $\delta \leq 2$. Then $\eta = \eta_a$ if and only if one of the following holds.*

- (i) *If G has no cut edge, then G is a graph such that an end block of G is either a theta graph or a cycle with exactly one chord whose vertices of degree 3 are not cut vertices and all other blocks are cycles and the block-cut point tree of G is a path (Refer Figure 1).*



Figure 1.

- (ii) *If G has a cut edge with $|\mathcal{H}_G| = m$ and if l is the number of end blocks in $\mathcal{G}(f)$ with f as a cut vertex, then $m = l$, where \mathcal{H}_G is as defined in Theorem 2.3 (Refer Figure 2).*




Figure 2.

Proof. If G is of type (i) by Lemma 1.12 and Theorem 2.3, $\eta = \eta_a = q - p + 1$. If G is of type (ii), let $|\mathcal{H}_G| = m$ and let n be the number of pendant vertices in G . Then by Theorem 2.4 and Theorem 2.6, $\eta = \eta_a = q - p + m + n$. Hence $\eta = \eta_a$.

Conversely suppose $\eta = \eta_a$. We first prove the theorem when $\delta = 2$. Since $\eta_a \neq q - p$, we have $\eta \neq q - p$. Hence by Theorem 1.9, Theorem 1.10 and Theorem 1.11, G is either

- (a) a block which is either a cycle or a cycle with exactly one chord or a theta graph, or
- (b) a graph in which each block is a cycle or a cycle with exactly one chord or a theta graph and at most one block is not a cycle, or
- (c) a graph which has a cut edge e such that at least one component of $G - e$ is a graph whose blocks are all cycles.

If G is of type (a) or (b) then by Theorem 2.2, $\eta = q - p + 1$. Suppose G is of type (a), a clock. If G is a cycle then clearly $\eta \neq \eta_a$. Hence G is a cycle with exactly one chord or a theta graph and G reduces to a graph of type (i) given in the theorem. Suppose G is of type (b), not a clock. Let s be the number of end blocks of G which are cycles. If each block of G is a cycle then by Theorem 2.5, $\eta_a = q - p + s$. Since $\eta = \eta_a$, $s = 1$ which is a contradiction to the fact that G is not a block. Hence there exists a block B in G which is not a cycle. We claim that $s = 1$. Suppose $s > 1$. Then by Theorem 2.5,

$$\eta_a = \begin{cases} q - p + s + 1, & \text{if } B \text{ is an end block of } G \text{ and a vertex of} \\ & \text{degree 3 is a cut vertex of } G, \\ q - p + s, & \text{otherwise.} \end{cases}$$

Hence $\eta_a - \eta = s$ or $s - 1$ and $s > 1$ which is a contradiction to the fact that $\eta = \eta_a$. Hence $s = 1$ and this proves that the block-cut point tree of G is a path. If a vertex of degree 3 in B is a cut vertex then again $\eta_a = q - p + 2$ which is a contradiction. Hence vertices of degree 3 in B are not cut vertices and G reduces to a graph of type (i) given in the theorem.

If G is of type (c), let l be the number of end blocks in G which are in $\mathcal{G}(f)$ with f as a cut vertex and let $|\mathcal{H}_G| = m$. Then by Theorem 2.3 and Theorem 2.5, $\eta = q - p + m$, $\eta_a = q - p + l$. Since $\eta = \eta_a$, $m = l$. Thus G reduces to a graph of type (ii) given in the theorem.

Now let $\delta = 1$. Let n be the number of pendant vertices of G . We define l, m as before. By Theorem 2.4 and Theorem 2.6, $\eta = q - p + m + n$ and $\eta_a = q - p + l + n$. Since $\eta = \eta_a$, $l = m$. Thus G reduces to a graph of type (ii) given in the theorem.

References

- [1] B. D. Acharya and E. Sampathkumar, *Graphoidal covers and graphoidal covering number of a graph*, Indian J. Pure Appl. Math. **18**(1987), 882-890.
- [2] B. D. Acharya and Purnima Gupta, *Domination in graphoidal covers of a graph*, Discrete Mathematics **206**(1999), 3-33.

- [3] S. Arumugam, B. D. Acharya and E. Sampathkumar, *Graphoidal covers of a graph a certificate review.*, Graph Theory and Its Applications, Proceedings of the National Workshop, Manonmaniam Sundaranar University, Tirunelveli, Eds. S. Arumugam, B. D. Acharya and E. Sampathkumar, Tata-McGraw Hill (1997), 1-28.
- [4] S. Arumugam, Indra Rajasingh, R. Bharathi and P. Roushini Leely Pushpam, *On graph whose graphoidal covering number is one less than its cyclomatic number*, Graph theory and Its Applications, Proceedings of the National Workshop, Manonmaniam Sundaranar University, Thirunelveli, Eds. S. Arumugam, B. D. Acharya and E. Sampathkumar, Tata-McGraw Hill (1997), 29-34.
- [5] S. Arumugam, Indra Rajasingh and P. Roushini Leely Pushpam, *On graphs whose acyclic graphoidal covering number is one less than its maximum degree*, Discrete Mathematics **240**(2001), 231-237.
- [6] S. Arumugam, Indra Rajasingh and P. Roushini Leely Pushpam, *A note on the graphoidal covering number of a graph*, Journal of Discrete Mathematical Sciences and Cryptography **5**(2002), 145-150.
- [7] S. Arumugam, Indra Rajasingh and P. Roushini Leely Pushpam, *On graphs whose acyclic graphoidal covering number is one less than its cyclomatic number - To appear in Ars Combinatoria.*
- [8] S. Arumugam, Indra Rajasingh and P. Roushini Leely Pushpam, *Characterization of graphs with unique minimum graphoidal cover - To appear in Tamkang Journal of Mathematics, Tamkang University Republic of China.*
- [9] S. Arumugam, Indra Rajasingh and P. Roushini Leely Pushpam, *Graphs with unique minimum acyclic graphoidal cover-I - To appear in the Journal of Discrete Mathematical Sciences and Cryptography.*
- [10] S. Arumugam and C. Pakkiam, *The graphoidal covering number of hypergraphs*, International Journal of Management and Systems **10**(1994), 195-200.
- [11] S. Arumugam and C. Pakkiam, *Graphs with unique minimum graphoidal cover*, Indian J. Pure Appl. Math. **25**(1994), 1147-1153.
- [12] S. Arumugam and C. Pakkiam, *Graphoidal bipartite graphs*, Graphs and Combinatorics **10**(1994), 305-310.
- [13] S. Arumugam and C. Pakkiam, *The graphoidal covering number of directed graphs*, J. Math. Phy. Sci. **29**(1995), 33-41.
- [14] S. Arumugam and J. Suresh Suseela, *Acyclic graphoidal covering number of digraphs*, Proceedings of the conference on Graph Connections, Cochin University, Cochin, Eds. R. Balakrishnan *et al.*, Allied Publishers (1998), 72-78.
- [15] S. Arumugam and J. Suresh Suseela, *Acyclic graphoidal covers and path partitions in a graph*, Discrete Mathematics **199**(1998), 67-77.
- [16] F. Harary, Graph Theory, Addison Wesley, Reading Mass, 1969.
- [17] C. Pakkiam and S. Arumugam, *On the graphoidal covering number of a graph*, Indian J. Pure Appl. Math. **20**(1989), 330-333.
- [18] C. Pakkiam and S. Arumugam, *The graphoidal covering number of unicyclic graphs*, Indian J. Pure Appl. Math. **23**(1992), 141-143.
- [19] C. Pakkiam and S. Arumugam, *The graphoidal graphs of a tree*, Indian J. Pure Appl. Math. **21**(1990), 1055-1058.

Department of Mathematics, Loyola College, Chennai - 600 034, India.

Department Mathematics, D. B. Jain College, Chennai - 600 096, India.