

Evaluation of photosynthetic efficacy and CO₂ removal of microalgae grown in an enriched bicarbonate medium

S. Abinandan¹ · S. Shanthakumar¹

Received: 25 September 2015 / Accepted: 14 December 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract Bicarbonate species in the aqueous phase is the primary source for CO₂ for the growth of microalgae. The potential of carbon dioxide (CO₂) fixation by *Chlorella pyrenoidosa* in enriched bicarbonate medium was evaluated. In the present study, effects of parameters such as pH, sodium bicarbonate concentration and inoculum size were assessed for the removal of CO₂ by *C. pyrenoidosa* under mixotrophic condition. Central composite design tool from response surface methodology was used to validate statistical methods in order to study the influence of these parameters. The obtained results reveal that the maximum removal of CO₂ was attained at pH 8 with sodium bicarbonate concentration of 3.33 g/l, and inoculum size of 30 %. The experimental results were statistically significant with R^2 value of 0.9527 and 0.960 for CO₂ removal and accumulation of chlorophyll content, respectively. Among the various interactions, interactive effects between the parameters pH and inoculum size was statistically significant ($P < 0.05$) for CO₂ removal and chlorophyll accumulation. Based on the studies, the application of *C. pyrenoidosa* as a potential source for carbon dioxide removal at alkaline pH from bicarbonate source is highlighted.

Keywords Carbon concentrating mechanism · Mixotrophic condition · *Chlorella pyrenoidosa* · Dissolved inorganic carbon · Response surface methodology

Introduction

Rapid exploitation of fossil fuels such as coal, petroleum, etc., has invariably increased the concentration of CO₂ in the atmosphere to ~400 ppm (Rahaman et al. 2011). Several steps have been taken to control the CO₂ emission rates caused during the pre and post-process of various man-made activities. Some of the process includes application of alkanolamine absorbents (Choi et al. 2012; Kim et al. 2013; Pires et al. 2011), desiccant absorption (Stewart and Hessami 2005), adsorption by activated carbon (Lu et al. 2008), mineral carbonate (Wang et al. 2008), zeolite (Wang et al. 2011), molecular sieve, ocean storage (Metz et al. 2005) and geological storage (Holloway 2007). Biofixation of CO₂ by microalgae attracted researchers due to its affirmative behavior such as carbon neutral and biomass productivity for its various applications. However, the factors such as economic aspects, life cycle analysis, capital investment and other environment aspects have made these technological impacts slower at commercial level. Unlike other gases, CO₂ has some significant limiting factors such as low mass transfer rate, high cost of CO₂ capture, its transportation. Meanwhile, when the CO_{2(g)} is passed onto the aqueous solution, it reduces the pH due to the formation of H₂CO₃ (free CO₂) and escapes into the atmosphere thereby resulting in significant CO₂ loss during algae culture (Chi et al. 2011). Furthermore, pH is the major determinant of the relative concentrations of the carbonaceous system species in water and could affect the availability of carbon for algal photosynthesis in intensive cultures and at certain situations where the supply of adequate CO_{2(g)} is limited, inorganic carbon sources such as bicarbonate salts (NaHCO₃) is best alternative to cultivate microalgae (Hsueh et al. 2009; Lam and Lee 2013; Benemann 1993). Meanwhile, it is also envisaged that

✉ S. Shanthakumar
shanthakumar.s@vit.ac.in

¹ Environmental Engineering Division, School of Mechanical and Building Sciences, VIT University, Vellore 632014, India

mixotrophic mode of cultivating microalgae is best in terms of growth and other applications (Abreu et al. 2012; Bhatnagar et al. 2011; Cheirsilp and Torpee 2012). Based on literature, it can be established that *Chlorella* sp. is found to be most prominent and efficient in bio-fixing HCO_3^- and can grow well at mixotrophic conditions (Jeong et al. 2003; Lam et al. 2012).

Accordingly, this study has attempted to add to the current knowledge for carbon sequestration by optimizing process parameters for maximum CO_2 removal using microalgae in enriched bicarbonate medium. In this context, we aim to discuss the effect of pH, sodium bicarbonate concentration and inoculum size in mixotrophic mode of cultivation for CO_2 removal in microalgae. In addition, we have also monitored chlorophyll as indicator for growth and photosynthetic productivity. Also, in order to optimize the chosen factors and its interactions, central composite design (CCD) using response surface methodology has been employed. The application of response surface methodology is to maximize the effective parameters and minimize the experiments as well as to study the variables individual and interactive effects (Wang et al. 2007; Zhang et al. 2009; Khataee and Dehghan 2011).

Materials and methods

Culturing of microalgae

Chlorella pyrenoidosa (NCIM 2738) was obtained from the National Centre of Industrial Microorganism (NCIM), Pune, India and was maintained in BG-11 medium (UTEX 2009) under axenic conditions. The stock solution was prepared for all the constituents in media with the following composition for 100 ml: K_2HPO_4 0.4 g, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 0.36 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.75 g, NaNO_3 15 g, citric acid 0.06 g, $\text{Na}_2\text{EDTA} \cdot 2\text{H}_2\text{O}$ 0.01 g, sodium carbonate 0.2 g, ammonium ferric citrate 0.06 g, and A_5 trace solution (g/l) components as H_3BO_3 2.86, MnCl_2 1.81, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ 0.222, $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ 0.390, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ 0.079, $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ 0.0494. From the stock solution the media has been prepared for 500 ml with 5 ml from each constituent and 0.5 ml of A_5 trace solution. The culture was subjected to continuous illumination (14 WTL5 tungsten filament lamps; Philips Co.) with 1500 Lux measured using TES light meter (TES CORP) at room temperature 28 ± 1 °C.

Design of experiments

The exponential phase microalgae cells (28×10^5 cells/ml) were taken for experimental studies in synthetic medium (Feng et al. 2011) with the following composition (g/l)

glucose 0.4125, NH_4Cl 0.078, KH_2PO_4 0.018, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.013, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 0.043, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.005, A_5 Trace solution (1 ml/l), respectively. All the experiments in the study were carried in 500 ml conical flasks containing 300 ml of working solution of synthetic medium with variables (pH, inoculum size and sodium bicarbonate). For analysis, 10 ml of sample was collected using autoclaved syringe (20 ml) for every 3 days once and was subjected for centrifugation at 8000 RPM for 10 min. The pellet was taken for estimation of chlorophyll and supernatant for CO_2 analysis. The pH values were chosen to study the microalgae behavior to CO_2 in acidic (pH 4 to mimic more availability of free CO_2), pH 6 (as standard growth medium range) and alkaline range (pH 8). The sodium bicarbonate concentrations were fixed of 1–3 g per 300 ml each which was scaled up in g/l as depicted in Table 1. The inoculum sizes were fixed (10–30 %) on volume per volume basis as to envisage its effects for CO_2 removal and the optical density of 10, 20 and 30 % inoculum sizes used in the study were recorded as 0.594, 0.692 and 0.802, respectively, measured at 660 nm using UV Visible Spectrophotometer (Cyberlab, USA). The pH of the medium was adjusted using 0.1 N NaOH and 0.1 N HCl. All the flasks were manually shaken thrice a day in order to avoid sticking of culture to flasks. The experiments were carried out in duplicates and the average has been taken RSM analysis.

Response surface methodology


In order to study the combined effects of the variables (pH, inoculum size and sodium bicarbonate) on the maximum removal of CO_2 (%), 20 sets of experiments with appropriate combinations of pH, inoculum size and NaHCO_3 concentration were conducted using response surface method (statistical analysis) and the details are presented in Table 2. The CCD under the response surface methodology (RSM) was employed in order to illustrate the nature of the response surface in the experimental design and to elucidate the optimal conditions of the most significant independent variables. A flowchart representation in support of the response surface methodology has been presented in Fig. 1. In this analysis, NaHCO_3 , inoculum size and pH were chosen as independent variables and the carbon

Table 1 Experimental range and levels of independent variables

Independent variables	Design variables	Range and levels		
		−1	0	1
pH	A	4	6	8
NaHCO_3 (g/l)	B	3.33	6.66	10
Inoculum size (%)	C	10	20	30

Table 2 Central composite design matrix with coded and uncoded variables

Run	A	B	C	pH	NaHCO ₃ (g/l)	Inoculum size (%)
1	-1	-1	-1	4.0	3.33	10
2	1	-1	-1	8.0	3.33	10
3	-1	1	-1	4.0	10	10
4	1	1	-1	8.0	10	10
5	-1	-1	1	4.0	3.33	30
6	1	-1	1	8.0	3.33	30
7	-1	1	1	4.0	10	30
8	1	1	1	8.0	10	30
9	-1.68179	0	0	2.6	6.66	20
10	1.68179	0	0	9.4	6.66	20
11	0	-1.68179	0	6.0	1.06	20
12	0	1.68179	0	6.0	12.27	20
13	0	0	-1.68179	6.0	6.66	3.18
14	0	0	1.68179	6.0	6.66	36.81
15	0	0	0	6.0	6.66	20
16	0	0	0	6.0	6.66	20
17	0	0	0	6.0	6.66	20
18	0	0	0	6.0	6.66	20
19	0	0	0	6.0	6.66	20
20	0	0	0	6.0	6.66	20

**Fig. 1** Flow chart representation of response surface methodology

dioxide (CO₂) removal rate (%) and its corresponding chlorophyll content were taken as dependent output response variable.

The three independent variables were varied over two levels with pH between (4 and 8) relative to the centre point (pH 6), the second independent variable (NaHCO₃ in g/l) was varied over two levels (3.3 and 9.9 g/l) relative to the centre point (6.6 g/l) and the third independent variable (inoculum size in %) was varied over two levels (10 and 30 %) relative to the centre point (20 %).

The full factorial CCD matrices of three variables with respect to their uncoded (real) and coded values are presented in Table 2. The response surface method was constructed using MINITAB 16 statistical software. Evaluation of the goodness of fit of the model is done through coefficient determination and analysis of variances. The experimental results were fitted to a second order polynomial Eq. (1):

$$Y = \beta_0 + \beta_1A + \beta_2B + \beta_3C + \beta_{11}A^2 + \beta_{22}B^2 + \beta_{33}C^2 + \beta_{12}AB + \beta_{13}AC + \beta_{23}BC \quad (1)$$

where Y is the dependent variable (CO₂ removal and chlorophyll content); A , B and C are the independent variable; β_0 is the regression coefficient at center point; β_1 , β_2 and β_3 are the linear coefficients; β_{11} , β_{22} and β_{33} are the quadratic coefficients and β_{12} , β_{13} and β_{23} are the second-order interaction coefficients. The developed regression model was evaluated by analyzing the values of regression coefficients, analysis of variance (ANOVA), P and F values. The quality of fit of the polynomial model equation was expressed by the coefficient of determination, R^2 . The statistical software package was used to identify the experimental design as well as to generate a regression model to predict the optimum combinations considering the effects of linear, quadratic and interactive effects on CO₂ removal and corresponding chlorophyll content.

Carbon dioxide removal

Alkalinity based titrimetric method was used to find out the dissolved inorganic carbon species (H_2CO_3^* , HCO_3^- , CO_3^{2-}) by following standard method (APHA 2005). Briefly, all the samples were centrifuged to obtain supernatant which was titrated against standardized H_2SO_4 (0.02 N). In a typical experiment, 10 ml of supernatant was taken in a beaker and titrated against H_2SO_4 using phenolphthalein (pKa 8.6) indicator. After the first end point, titration was continued using methyl orange (pKa 3.8) as the indicator to get the second end point. The readings were noted down and the dissolved inorganic carbon species (H_2CO_3^* , HCO_3^- , CO_3^{2-}) were determined using the formulae in standard method and the sum of the three species are represented as total inorganic carbon species CO_2 as mg CaCO_3/l which is shown in Eq. (2). Furthermore, the ratio of CO_2 to CaCO_3 is 1.4 which is taken for CO_2 estimation from total inorganic carbon species and the same has been shown in the Eq. (3) (Kemmer 1979).

$$\text{CO}_{2(\text{asCaCO}_3)} \text{ mg/l} = \text{H}_2\text{CO}_3^* + \text{HCO}_3^- + \text{CO}_3^{2-} \quad (2)$$

$$\text{CO}_{2(\text{asCaCO}_3)} \text{ mg/l} \div 1.14 = \text{CO}_{2(\text{asCO}_2)} \text{ mg/l} \quad (3)$$

The CO_2 removal (%) was determined by calculating difference between the initial concentration of CO_2 (based on Eq. 3) from each experimental runs and final concentration after growth in stationary phase (until no observation of CO_2 removal was found) and is expressed below

$$\text{CO}_2 \text{ removal (\%)} = \frac{\text{Initial CO}_2 - \text{final CO}_2}{\text{Initial CO}_2} \times 100 \quad (4)$$

Chlorophyll analysis

The chlorophyll content in the medium is determined by spectrometric analysis. Briefly, 5 ml algae culture was centrifuged at 10,000 rpm for 10 min. The supernatant was drained off and the sample was re-suspended in ethanol/diethyl ether and kept boiling for 5 min. After boiling, the sample was made up to 5 ml with ethanol/diethyl ether. The optical density was measured at 660 nm and 642.5 nm with solvent as a blank. The chlorophyll content was determined using the formula (Becker 1994):

$$\text{Chlorophyll (mg/l)} = (9.9 * \text{OD}_{660}) - (0.77 * \text{OD}_{642.5}) \quad (5)$$

Results and discussion

Central composite design: response surface estimation

The growth of *C. pyrenoidosa* under experimental conditions and optimized condition is shown in Fig. 2. It can be




Fig. 2 Growth profile of microalgae under optimized condition [pH 8, NaHCO_3 3.33 (g/l), inoculum size –30 (%)] and experimental condition

noted that, there was a twofold increase in the chlorophyll accumulation in the optimized condition compared to the experimental condition which may be due to the availability of carbon source that helps in synthesizing chlorophyll. The experiments based on design (Table 2) were lasted for 20 days after which the growth was stationary and hence no further improvement was observed on chlorophyll accumulation and CO_2 removal and the same has been given as experimental data in RSM. The results of CO_2 removal and chlorophyll content (both predicted and experimental) for different design variables (pH, inoculum size and NaHCO_3) are presented in Table 3. The second-order polynomial Eq. (1) are fitted with experimental results of maximum CO_2 removal (%) and corresponding chlorophyll content from the estimated regression coefficients. The regression coefficients for CO_2 removal and chlorophyll content are presented as Eqs. (6) and (7), respectively.

$$Y = 67.6120 + 9.8638A - 6.6875B + 8.3419C + 0.0318A^2 - 0.3245B^2 - 1.5856C^2 + 2.9458AB - 5.0963AC - 0.6830BC \quad (6)$$

$$Y = 2.9031 + 0.6274A - 0.2143B + 0.1584C - 0.1050A^2 + 0.3559B^2 + 0.7672C^2 - 0.3266AB - 0.8803AC - 0.7276BC \quad (7)$$

Analysis of variance and residuals

The regression helps to correlate the experimental data with predicted response. R^2 and adjusted R^2 values represent the proportion of variation in the response that is explained by the model where R describes the amount of

Table 3 Central composite design matrix and the output responses for CO₂ removal and chlorophyll

Run	pH	NaHCO ₃ (g/l)	Inoculum size (%)	CO ₂ removal (%)(experimental)	CO ₂ removal (%)(predicted)	Chlorophyll (mg/l)(experimental)	Chlorophyll (mg/l)(predicted)
1	4.0	3.33	10	53.07	51.49	1.016	1.415
2	8.0	3.33	10	80.00	75.52	5.240	5.083
3	4.0	10.00	10	33.95	33.39	3.499	3.095
4	8.0	10.00	10	74.37	69.20	5.064	5.457
5	4.0	3.33	30	78.07	79.73	5.263	4.947
6	8.0	3.33	30	86.32	83.38	4.614	5.095
7	4.0	10.00	30	57.93	58.90	3.484	3.717
8	8.0	10.00	30	76.25	74.33	2.880	2.558
9	2.6	6.66	20	53.07	51.11	1.461	1.550
10	9.4	6.66	20	77.37	84.29	3.858	3.661
11	6.0	1.06	20	75.42	78.07	4.475	4.270
12	6.0	12.27	20	52.98	55.25	3.452	3.549
13	6.0	6.66	3.18	43.75	49.07	4.906	4.806
14	6.0	6.66	36.81	77.50	77.13	5.347	5.339
15	6.0	6.66	20	67.10	66.96	2.895	2.898
16	6.0	6.66	20	67.10	66.96	2.895	2.898
17	6.0	6.66	20	67.10	66.96	2.895	2.898
18	6.0	6.66	20	67.10	66.96	2.895	2.898
19	6.0	6.66	20	67.10	66.96	2.895	2.898
20	6.0	6.66	20	67.10	66.96	2.895	2.898

variation in the observed responses. The value of R^2 is also a measure of fit of the model and the adjusted R^2 value compares models with different independent variables. The ANOVA results are presented in Tables 4 and 5 and it can be noted that, the R^2 values for CO₂ removal and corresponding chlorophyll content are 0.9527 and 0.962 which

Table 4 ANOVA for fit of CO₂ removal (%) from central composite design

Sources of variation	Sum of squares	Degree of freedom	Mean square	<i>F</i> value	<i>P</i>
Regression	3266.42	9	358.49	22.37	0.0000
Residuals	159.50	10	16.02		
Total	3386.65				

$$R^2_{(\text{pred.})} = 95.27\%; R^2_{(\text{adj.})} = 91.01\%$$

Table 5 ANOVA for fit of chlorophyll estimation from central composite design

Sources of variation	Sum of squares	Degrees of freedom	Mean square	<i>F</i> value	<i>P</i>
Regression	27.8792	9	3.09769	28.24	0.000
Residuals	1.0999	10	0.10999		
Total	28.9791				

$$R^2_{(\text{pred.})} = 96.20\%; R^2_{(\text{adj.})} = 92.79\%$$

indicates high degree of correlation between experimental and predicted values.

The estimated regression coefficients for removal of CO₂ and chlorophyll content are presented in Tables 6 and 7, respectively, along with their corresponding *P* value and *T* values. It can be observed from Table 6 for CO₂ removal (%) that, the coefficient for single effect of pH (β_1), sodium bicarbonate (β_2) and inoculum size (β_3) ($P < 0.050$) are highly significant whereas the square effects i.e., linear coefficient β_{11} , β_{22} and β_{33} and the interactive terms β_{23} are not significant except β_{13} which is significant. From Table 7 (chlorophyll content) the coefficients for single effect except inoculum size (β_3) are significant whereas the interactive and square effects except β_{11} are significant.

Response surface plots for CO₂ removal

The main objective of the response surface method is to find out the optimum condition for maximum CO₂ removal with respect to the chosen variables. The interaction effects between the variables (pH and sodium bicarbonate) for carbon dioxide removal is presented in Fig. 3a, as the sodium bicarbonate concentration increases concomitantly with increase in pH, the removal of CO₂ increases. This is due to the fact that the microalgae uptake the HCO₃ inside the cell and by the action of carbonic anhydrase, it converts HCO₃ into CO₂ inside the cell. The observations are

Table 6 Estimated regression coefficients for CO₂ removal (%)


Term	Coefficient	Standard error	<i>T</i>	<i>P</i>
β_0	67.6120	1.6323	41.415	0.000
β_1	9.8638	1.083	9.107	0.000
β_2	-6.7875	1.083	-6.266	0.000
β_3	8.3419	1.083	7.701	0.000
β_{11}	0.0318	1.054	0.030	0.977
β_{22}	-0.3245	1.054	-0.308	0.765
β_{33}	-1.5856	1.054	-1.504	0.164
β_{12}	2.9458	1.415	2.081	0.064
β_{13}	-5.0963	1.415	-3.601	0.005
β_{23}	-0.6830	1.415	-0.483	0.640

Table 7 Estimated regression coefficients for chlorophyll

Term	Coefficient	Standard error	<i>T</i>	<i>P</i>
β_0	2.9031	0.13526	21.463	0.000
β_1	0.6274	0.08974	6.991	0.000
β_2	-0.2143	0.08974	-2.388	0.038
β_3	0.1584	0.08974	1.765	0.108
β_{11}	-0.1050	0.08736	-1.202	0.257
β_{22}	0.3559	0.08736	4.074	0.002
β_{33}	0.7672	0.08736	8.781	0.000
β_{12}	-0.3266	0.11726	-2.786	0.019
β_{13}	-0.8803	0.11726	-7.513	0.000
β_{23}	-0.7276	0.11726	-6.210	0.000

consistent with the results reported in the literature (Devogswami et al. 2011).

Similarly, the interaction effects between the variables (inoculum size and pH) for carbon dioxide removal is presented in Fig. 3b, it can be observed at low pH, even if the inoculum size is high; there is a decrease in the CO₂ removal. This could be due to the fact that at low pH all the carbon dioxide will exist in the form of free CO₂ (H₂CO₃) as the algae cannot utilize for metabolic activity (Van Den Hende et al. 2012). Figure 3c depicts the interaction effect between the variables (inoculum size and sodium bicarbonate) for CO₂ removal. It can be noted from the figure that, increase in inoculum size with concomitant decrease in sodium bicarbonate concentration, helps to increase the uptake of CO₂ (i.e. CO₂ removal). This could be due to the regulation of H⁺ ions into the cell takes place due to which the hydroxide is formed which in turn affects the CO₂ removal process (Yeh et al. 2010). In addition, it is also reported that maximum CO₂ removal varies with different environment conditions. It is reported that at pH 4 and high bicarbonate concentration (0.3 g/l), maximum removal of CO₂ (82.5 %) was achieved (Lam and Lee 2013). Similarly, in another study (Yeh and Chang 2012), it is reported that a maximum removal of CO₂

**Fig. 3** Surface plot for CO₂ removal (%). **a** Conc. of NaHCO₃ (g/l), pH. **b** Inoculum size (%), pH. **c** Inoculum size (%), Conc. of NaHCO₃ (g/l)

in the range of 90–100 % was achieved with increasing bicarbonate concentration (till 1600 mg/l).

In the present study, the maximum absorption of CO₂ was achieved at pH 8 with initial NaHCO₃ concentration of 3.33 g/l and higher inoculum size 30 %. However, with increase in concentration of bicarbonate, there is no significant change in CO₂ removal which is due to the fact that additional carbon mechanism results in decreased growth rate causing them in reduction of fixing CO₂ rates (Lam and Lee 2013).





Fig. 4 Surface plot for chlorophyll (mg/l). **a** Conc. of NaHCO₃ (g/l), pH. **b** Inoculum size (%), pH. **c** Inoculum size (%), Conc. of NaHCO₃ (g/l)

Response surface plots for chlorophyll content

Chlorophyll *a* is the photosynthetic pigment that is widely present in all eukaryotic microalgae and is synthesized during photosynthetic process as mentioned in Eq. (8). Initially the inorganic carbon is accumulated into microalgal cell by capture of light energy by pigments such as chlorophyll *a* and *b* that are presented in chloroplastida of cell (Wang et al. 2012).



The initial concentrations of chlorophyll are 0.5, 1, 1.34 mg/l for inoculum sizes 10, 20, 30 %, respectively. The surface

plots for the interaction effects between the variables (pH and sodium bicarbonate) for the chlorophyll content are presented in Fig. 4a. It can be observed from the figure that, at low sodium bicarbonate concentration, as the pH increases the chlorophyll content increases drastically but at higher sodium bicarbonate concentration only minor variations of chlorophyll content was observed in all pH. This could be due to the excess HCO₃⁻ (Ci) ions which are not utilized by microalgae due to insufficient light, as synthesis of chlorophyll will mainly depend upon the light energy during photosynthesis thereby hindrance the process of synthesizing chlorophyll (Amoroso et al. 1998).

Figure 4b depicts the interaction effects between the variables (inoculum size and pH) for chlorophyll and it can be noticed that at low inoculum size, the chlorophyll content increases with increase in pH. However, the chlorophyll content decreases when the inoculum size increases along with increase in pH. It may be due to the fact that increase in inoculum size causes reduction of H⁺ ions form the medium (Liu et al. 2014; Chi et al. 2011).

The interaction effects between the variables (inoculum size and sodium bicarbonate) are presented in Fig. 4c and it can be observed from the figure, that the chlorophyll content decreases when the sodium bicarbonate increases along with the increase in inoculum size. This is due to the fact that the increase in sodium bicarbonate decreases the photosynthetic efficiency. The obtained results are in consistent with the results reported in the literature (White et al. 2013).

At higher concentration of bicarbonate and pH, there is no significant change in chlorophyll content which could be due to: (1) increasing rates of photorespiration (2) availability of C:N ratio in the medium as the nitrogen is also one of the factor for chlorophyll assimilation.

Conclusion

The interactive effects between the variables (pH, inoculum size and sodium bicarbonate concentration) for CO₂ removal using *C. pyrenoidosa* in synthetic medium under mixotrophic condition was investigated by employing response surface method with CCD. The maximum removal of CO₂ (86 %) was achieved at pH 8 with sodium bicarbonate concentration of 3.33 g/l, and inoculum size of 30 %. The regression value of 0.9527 and 0.962 was observed for CO₂ removal and chlorophyll content which implies that the experimental results are statistically significant. The results reveal that *C. pyrenoidosa* can be used effectively for bio-fixation of CO₂ in the form of bicarbonate at alkaline conditions in higher inoculum size under mixotrophic condition. Further investigations on adapting

microalgae at alkaline conditions will be a useful for large-scale applications to develop bioenergy feedstock.

Acknowledgments The author's gratefully acknowledge VIT University for providing necessary funds and facilities to carry out the research work. Also, the author's expresses their sincere gratitude to V. Sowmya, Research Scholar, School of Mechanical and Building Sciences, VIT University for assisting in writing and proof reading the article.

Compliance with ethical standards

Conflict of interest The authors declare that there is no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Abreu AP, Fernandes B, Vicente AA, Teixeira J, Dragon G (2012) Mixotrophic cultivation of *Chlorella vulgaris* using industrial dairy waste as organic carbon source. *Bioresour Technol* 118:61–66
- Amoroso G, Sultemeyer D, Thyssen C, Fock H (1998) Uptake of HCO_3^- and CO_2 in cells and chloroplasts from the microalgae *Chlamydomonas reinhardtii* and *Dunaliella tertiolecta*. *Plant Physiol* 116:193–201
- American public health association (APHA) (2005) Standard methods for the examination of water and wastewater, 21st edn. AWWA-WPCF, Washington
- Becker EW (1994) Microalgae: biotechnology and microbiology. Cambridge University Press, New York
- Benemann JR (1993) Utilization of carbon dioxide from fossil fuel-burning power plants with biological systems. *Energy Convers Manag* 34:999–1004
- Bhatnagar A, Chinnasamy S, Singh M, Das KC (2011) Renewable biomass production by mixotrophic algae in the presence of various carbon sources and wastewaters. *Appl Energy* 88:3425–3431
- Chairsilp B, Torpee S (2012) Enhanced growth and lipid production of microalgae under mixotrophic culture condition: effect of light intensity, glucose concentration and fed-batch cultivation. *Bioresour Technol* 110:510–516
- Chi Z, O'Fallon JV, Chen S (2011) Bicarbonate produced from carbon capture for algae culture. *Trends Biotechnol* 29:537–541
- Choi W, Kim G, Lee K (2012) Influence of the CO_2 absorbent monoethanolamine on growth and carbon fixation by the green alga *Scenedesmus* sp. *Bioresour Technol* 120:295–299
- Devogswami CR, Kalita MC, Talukdar R, Bora J, Sharma P (2011) Studies on the growth behavior of *Chlorella*, *Haematococcus* and *Scenedesmus* sp. in culture media with different concentrations of sodium bicarbonate and carbon dioxide gas. *Afr J Biotechnol* 10:13128–13138
- Feng Y, Li C, Zhang D (2011) Lipid production of *Chlorella vulgaris* cultured in artificial wastewater medium. *Bioresour Technol* 102:101–105
- Holloway S (2007) In Carbon dioxide capture and geological storage. *Philos Trans R Soc* 36:1095–1107
- Hsueh HT, Li WJ, Chen HH, Chu H (2009) Carbon bio-fixation by photosynthesis of *Thermosynechococcus* sp. CL-1 and *Nannochloropsis oculata*. *J Photochem Photobiol B Biol* 95:33–39
- Jeong ML, Gillis JM, Hwang JY (2003) Carbon dioxide mitigation by microalgal photosynthesis. *Bull Korean Chem Soc* 24:1763–1766
- Kemmer FN (1979) The NALCO water handbook, 2nd edn. McGraw-Hill, USA
- Khataee AR, Dehghan G (2011) Optimization of biological treatment of a dye solution by macroalgae *Cladophora* sp. using response surface methodology. *J Taiwan Inst Chem Eng* 42:26–33
- Kim G, Choi W, Lee CH, Lee K (2013) Enhancement of dissolved inorganic carbon and carbon fixation by green alga *Scenedesmus* sp. in the presence of alkanolamine CO_2 absorbents. *Biochem Eng J* 78:18–23
- Lam MK, Lee KT (2013) Effect of carbon source towards the growth of *Chlorella vulgaris* for CO_2 bio-mitigation and biodiesel production. *Int J Greenh Gas Control* 14:169–176
- Lam MK, Lee KT, Mohamed AR (2012) Current status and challenges on microalgae-based carbon capture. *Int J Greenh Gas Control* 10:456–469
- Liu G, Qiao L, Zhang H, Zhao D, Su X (2014) The effects of illumination factors on the growth and HCO_3^- fixation of microalgae in an experiment culture system. *Energy* 78:40–47
- Lu C, Bai H, Wu B, Su F, Hwang JF (2008) Comparative study of CO_2 capture by carbon nanotubes, activated carbons and zeolites. *Energy Fuels* 22:3050–3056
- Metz B, Davidson O, De Coninck H, Meyer MLL (2005) IPCC report on carbon dioxide capture and storage, 1st edn. Cambridge University Press, New York
- Pires JCM, Martins FG, Alvim-Ferraz MCM, Simoes M (2011) Recent developments on carbon capture and storage: an overview. *Chem Eng Res Des* 89:1446–1460
- Rahaman MSA, Cheng L, Xu X, Zhang L, Chen H (2011) A review of carbon dioxide capture and utilization by membrane integrated microalgal cultivation processes. *Renew Sust Energy Rev* 15:4002–4012
- Stewart C, Hessami MA (2005) A study of methods of carbon dioxide capture and sequestration—the sustainability of a photosynthetic bioreactor approach. *Energy Convers Manag* 46:403–420
- UTEX (2009) BG11 Medium Composition. University of Texas Austin, USA. <http://web.biosci.utexas.edu/utex/mediaDetail.aspx?mediaID=26>. Accessed 22 Nov 2012
- Van Den Hende S, Vervaeren H, Boon N (2012) Flue gas compounds and microalgae: (bio-) chemical interactions leading to biotechnological opportunities. *Biotechnol Adv* 30:1405–1424
- Wang JP, Chen YZ, Ge XW, Yu HQ (2007) Optimization of coagulation–flocculation process for a paper-recycling wastewater treatment using response surface methodology. *Colloid surface A* 302:204–210
- Wang B, Li Y, Wu N, Lan CQ (2008) CO_2 bio-mitigation using microalgae. *Appl Microbiol Biotechnol* 79:707–718
- Wang Q, Luo J, Zhong Z, Borgna A (2011) CO_2 capture by solid adsorbents and their applications: current status and new trends. *Energy Environ Sci* 4:42–55
- Wang H, Fu R, Pei G (2012) A study on lipid production of the mixotrophic microalgae *Phaeodactylum tricornutum* on various carbon sources. *Afr J Microbiol Res* 6:1041–1047
- White DA, Pagarette A, Rooks P, Ali ST (2013) The effect of sodium bicarbonate supplementation on growth and biochemical composition of marine microalgae cultures. *J Appl Phycol* 25:153–165
- Yeh K, Chang JS (2012) Effects of cultivation conditions and media composition on cell growth and lipid productivity of indigenous microalga *Chlorella vulgaris* ESP-31. *Bioresour Technol* 105:120–127

- Yeh KL, Chang JS, Chen W (2010) Effect of light supply and carbon source on cell growth and cellular composition of a newly isolated microalga *Chlorella vulgaris* ESP-31. Eng Life Sci 10:201–208
- Zhang Z, Peng J, Qu W, Zhang L, Zhang Z, Li W, Wan R (2009) Regeneration of high-performance activated carbon from spent catalyst: optimization using response surface methodology. J Taiwan Inst Chem Eng 40:541–548