

RESEARCH

Open Access

# Existence results for impulsive neutral stochastic functional integro-differential inclusions with infinite delays

Jong Yeoul Park<sup>1</sup> and Jae Ug Jeong<sup>2\*</sup>

\*Correspondence: jujeong@deu.ac.kr  
<sup>2</sup>Department of Mathematics, Donggeui University, Busan, 614-714, South Korea  
Full list of author information is available at the end of the article

## Abstract

In this paper, we prove the existence of mild solutions for a class of impulsive neutral stochastic functional integro-differential inclusions with infinite delays in Hilbert spaces. The results are obtained by using the fixed-point theorem for multi-valued operators due to Dhage. An example is provided to illustrate the theory.

**MSC:** 93B05; 93E03

**Keywords:** impulsive equation; stochastic functional inclusion; mild solution; infinite delay

## 1 Introduction

In this paper, we shall consider the existence of mild solutions for impulsive neutral stochastic functional integro-differential inclusions with infinite delay of the following form:

$$d \left[ x(t) - g \left( t, x_t, \int_0^t a(t, s, x_s) ds \right) \right] dt \in [Ax(t) + f(t, x_t)] dt + F(t, x_t) dw(t), \quad t \in J = [0, b], t \neq t_k, \quad (1.1)$$

$$\Delta x(t_k) = x(t_k^+) - x(t_k^-) = I_k(x(t_k^-)), \quad k = 1, 2, \dots, m, \quad (1.2)$$

$$x(t) = \phi(t) \in L^2(\Omega, \mathcal{B}_H) \quad \text{for a.e. } t \in J_0 = (-\infty, 0], \quad (1.3)$$

where the state  $x(\cdot)$  takes values in a separable real Hilbert space  $H$  with inner product  $(\cdot, \cdot)$  and norm  $\|\cdot\|$ ,  $A$  is the infinitesimal generator of a compact analytic resolvent operator  $S(t)$ ,  $t \geq 0$ , in the Hilbert space  $H$ . Suppose that  $\{w(t) : t \geq 0\}$  is a given  $K$ -valued Brownian motion or Wiener process with a finite trace nuclear covariance operator  $Q \geq 0$  and  $L(K, H)$  denotes the space of all bounded linear operators from  $K$  into  $H$ . Further  $a : D \times \mathcal{B}_H \rightarrow H$ ,  $g : J \times \mathcal{B}_H \times H \rightarrow H$ ,  $f : J \times \mathcal{B}_H \rightarrow H$  and  $F : J \times \mathcal{B}_H \rightarrow \mathcal{P}(L_Q(K, H))$  are given functions, where  $D = \{(t, s) \in J \times J : s \leq t\}$ ,  $\mathcal{P}(L_Q(K, H))$  is the family of all nonempty subsets of  $L_Q(K, H)$  and  $L_Q(K, H)$  denotes the space of all  $Q$ -Hilbert-Schmidt operators from  $K$  into  $H$ , which will be defined in Section 2. Here,  $I_k \in C(H, H)$  ( $k = 1, 2, \dots, m$ ) are bounded functions. Furthermore, the fixed times  $t_k$  satisfies  $0 = t_0 < t_1 < t_2 < \dots < t_m < b$ ,  $x(t_k^+)$  and  $x(t_k^-)$  denote the right and left limits of  $x(t)$  at  $t = t_k$ .  $\Delta x(t_k) = x(t_k^+) - x(t_k^-) = I_k(x(t_k^-))$  represents the jump in the state  $x$  at time  $t_k$ , where  $I_k$  determines the size of jump. The his-

tories  $x_t : \Omega \rightarrow \mathcal{B}_H, t \geq 0$ , which are defined by setting  $x_t = \{x(t + s) : s \in (-\infty, 0]\}$ , belong to the abstract phase space  $\mathcal{B}_H$ , which will be defined in Section 2. The initial data  $\phi = \{\phi(t) : -\infty < t \leq 0\}$  is an  $\mathcal{F}_0$ -measurable,  $\mathcal{B}_H$ -valued random variables independent of  $\{w(t) : t \geq 0\}$  with finite second moment.

The theory of impulsive integro-differential inclusions has become an active area of investigation due to their applications in the fields such as mechanics, electrical engineering, medicine biology, ecology, and so on (see [1, 2] and references therein).

The existence of impulsive neutral stochastic functional integro-differential equations or inclusions with infinite delays have attracted great interest of researchers. For example, Lin and Hu [3] consider the existence results for impulsive neutral stochastic functional integro-differential inclusions with nonlocal initial conditions. Hu and Ren [4] studied the existence results for impulsive neutral stochastic functional integro-differential equations with infinite delays.

Motivated by the previous mentioned papers, we prove the existence of solutions for impulsive neutral stochastic functional integro-differential inclusions with infinite delays.

## 2 Preliminaries

Throughout this paper,  $(H, | \cdot |)$  and  $(K, | \cdot |_K)$  denote two real separable Hilbert spaces. Let  $(\Omega, \mathcal{F}, P; F)$  ( $F = \{\mathcal{F}_t\}_{t \geq 0}$ ) be a complete filtered probability space satisfying the requirement that  $\mathcal{F}_0$  contains all  $P$ -null sets of  $\mathcal{F}$ . An  $H$ -valued random variable is an  $\mathcal{F}$ -measurable function  $x(t) : \Omega \rightarrow H$  and the collection of random variables  $S = \{x(t), w : \Omega \rightarrow H | t \in J\}$  is called a stochastic process. Suppose that  $\{w(t) : t \geq 0\}$  is a cylindrical  $K$ -valued Wiener process with a finite trace nuclear covariance operator  $Q \geq 0$ , denote  $T_r Q = \sum_{i=1}^{\infty} \lambda_i = \lambda < \infty$ , which satisfies  $Qe_i = \lambda_i e_i$ . So, actually,  $w(t) = \sum_{i=1}^{\infty} \sqrt{\lambda_i} w_i(t) e_i$ , where  $\{w_i(t)\}_{i=1}^{\infty}$  are mutually independent one-dimensional standard Wiener process. We assume that  $\mathcal{F}_t = \sigma\{w(s) : 0 \leq s \leq t\}$  is the  $\sigma$ -algebra generated by  $w$  and  $\mathcal{F}_T = \mathcal{F}$ . Let  $\psi \in L(K, H)$  and define

$$|\psi|_Q^2 = T_r(\psi Q \psi^*) = \sum_{n=1}^{\infty} |\sqrt{\lambda_n} \psi e_n|^2.$$

If  $|\psi|_Q < \infty$ , then  $\psi$  is called a  $Q$ -Hilbert-Schmidt operator. Let  $L_Q(K, H)$  denote the space of all  $Q$ -Hilbert-Schmidt operator  $\psi : K \rightarrow H$ . The completion  $L_Q(K, H)$  of  $L(K, H)$  with respect to the topology induced by the norm  $|\cdot|_Q$ , where  $|\psi|_Q^2 = \langle \psi, \psi \rangle$  is a Hilbert space with the above norm topology.

Let  $A : D(A) \rightarrow H$  be the infinitesimal generator of a compact, analytic resolvent operator  $S(t), t \geq 0$ . Let  $0 \in \rho(A)$ . Then it is possible to define the fractional power  $(-A)^\alpha$  for  $0 < \alpha \leq 1$  as a closed linear operator with its domain  $D((-A)^\alpha)$  being dense in  $H$ . We denote by  $H_\alpha$  the Banach space  $D((-A)^\alpha)$  endowed with the norm  $\|x\|_\alpha = \|(-A)^\alpha x\|$ , which is equivalent to the graph norm of  $(-A)^\alpha$ .

**Lemma 2.1** ([5]) *The following properties hold:*

- (i) *If  $0 < \beta < \alpha \leq 1$ , the  $H_\alpha \subset H_\beta$  and the embedding is continuous and compact whenever the resolvent operator of  $A$  is compact.*
- (ii) *For every  $0 < \alpha < 1$ , there exists a positive constant  $c_\alpha$  such that*

$$\|(-A)^\alpha S(t)\| \leq \frac{C_\alpha}{t^\alpha}, \quad t > 0.$$

Now, we define the abstract phase space  $\mathcal{B}_h$ . Assume that  $h : (-\infty, 0] \rightarrow (0, \infty)$  is a continuous function with  $l = \int_{-\infty}^0 h(t) dt < \infty$ . For any  $a > 0$  we define

$$\mathcal{B}_h = \left\{ \psi : (-\infty, 0] \rightarrow H : (E|\psi(\theta)|^2)^{\frac{1}{2}} \text{ is a bounded and measurable function on } [-a, 0] \text{ and } \int_{-\infty}^0 h(s) \sup_{s \leq \theta \leq 0} (E|\psi(\theta)|^2)^{\frac{1}{2}} ds < \infty \right\}.$$

If  $\mathcal{B}_h$  is endowed with the norm

$$\|\psi\|_{\mathcal{B}_h} = \int_{-\infty}^0 h(s) \sup_{s \leq \theta \leq 0} (E|\psi(\theta)|^2)^{\frac{1}{2}} ds \quad \text{for all } \psi \in \mathcal{B}_h,$$

then  $(\mathcal{B}_h, \|\cdot\|_{\mathcal{B}_h})$  is a Banach space [6]. Now, we consider the space

$$\mathcal{B}_b = \left\{ x : (-\infty, b] \rightarrow H \text{ such that } x_k \in C(J_k, H) \text{ and there exist } x(t_k^+) \text{ and } x(t_k^-) \text{ with } x(t_k) = x(t_k^-), x_0 = \phi \in L^2(\Omega, \mathcal{B}_h) \text{ on } (-\infty, 0], k = 1, 2, \dots, m \right\},$$

where  $x_k$  is the restriction of  $x$  to  $J_k = (t_k, t_{k+1}]$ ,  $k = 0, 1, \dots, m$ . Let  $\|\cdot\|_b$  be a seminorm in  $\mathcal{B}_b$  defined by

$$\|x\|_b = \|x_0\|_{\mathcal{B}_h} + \sup_{0 \leq s \leq b} (E|x(s)|^2)^{\frac{1}{2}}, \quad x \in \mathcal{B}_b.$$

**Lemma 2.2** ([7]) *Assume that  $x \in \mathcal{B}_b$ , then for  $t \in J$ ,  $x_t \in \mathcal{B}_h$ . Moreover*

$$l(E|x(t)|^2)^{\frac{1}{2}} \leq \|x_t\|_{\mathcal{B}_h} \leq \|x_0\|_{\mathcal{B}_h} + l \sup_{0 \leq s \leq t} (E|x(s)|^2)^{\frac{1}{2}},$$

where  $l = \int_{-\infty}^0 h(s) ds < \infty$ .

We use the notation  $\mathcal{P}(H)$  for the family of all subsets  $H$  and denote

$$\begin{aligned} \mathcal{P}_{cl}(H) &= \{Y \in \mathcal{P}(H) : Y \text{ is closed}\}, \\ \mathcal{P}_{bd}(H) &= \{Y \in \mathcal{P}(H) : Y \text{ is bounded}\}, \\ \mathcal{P}_{cv}(H) &= \{Y \in \mathcal{P}(H) : Y \text{ is convex}\}, \\ \mathcal{P}_{cp}(H) &= \{Y \in \mathcal{P}(H) : Y \text{ is compact}\}. \end{aligned}$$

A multi-valued mapping  $\Gamma : H \rightarrow \mathcal{P}(H)$  is called upper semicontinuous (u.s.c) if for any  $x \in H$ , the set  $\Gamma(x)$  is a nonempty closed subset of  $H$  and if for each open set  $G$  of  $H$  containing  $\Gamma(x)$ , there exists an open neighborhood  $N$  of  $x$  such that  $\Gamma(N) \subseteq G$ .  $\Gamma$  is said to be completely continuous if  $\Gamma(B)$  is relatively compact for every bounded subset of  $B \subseteq H$ . If the multi-valued mapping  $\Gamma$  is completely continuous with nonempty compact values, then  $\Gamma$  is u.s.c. if and only if  $\Gamma$  has a closed graph, i.e.,  $x_n \rightarrow x, y_n \rightarrow y, y_n \in \Gamma(x_n)$  imply  $y \in \Gamma(x)$ .

**Definition 2.1** The multi-valued mapping  $F : J \times \mathcal{B}_h \rightarrow \mathcal{P}(H)$  is said to be  $L^2$ -Carathéodory if

- (i)  $t \mapsto F(t, v)$  is measurable for each  $v \in \mathcal{B}_h$ ,
- (ii)  $v \mapsto F(t, v)$  is u.s.c. for almost all  $t \in J$  and  $v \in \mathcal{B}_h$ ,
- (iii) for each  $q > 0$ , there exists  $h_q \in L^1(J, \mathbb{R}^+)$  such that

$$\|F(t, v)\|^2 = \sup_{f \in F(t, v)} E(|f|^2) \leq h_q(t),$$

for all  $\|v\|_{\mathcal{B}_h}^2 \leq q$  and for a.e.  $t \in J$ .

The following lemma is crucial in the proof of our main result.

**Lemma 2.3** ([8]) *Let  $I$  be a compact interval and  $H$  be a Hilbert space. Let  $F$  be an  $L^2$ -Carathéodory multi-valued mapping with  $N_{F,x} \neq \emptyset$  and let  $\Gamma$  be a linear continuous mapping from  $L^2(I, H)$  to  $C(I, H)$ . Then the operator*

$$\Gamma \circ N_F : C(I, H) \rightarrow \mathcal{P}_{cp,cv}(H), \quad x \mapsto (\Gamma \circ N_F)(x) = \Gamma(N_{F,x})$$

is a closed graph operator in  $C(I, H) \times C(I, H)$ , where  $N_{F,x}$  is known as the selectors set from  $F$ ; it is given by

$$\sigma \in N_{F,x} = \{ \sigma \in L^2(L(K, H)) : \sigma(t) \in F(t, x) \text{ for a.e. } t \in J \}.$$

**Theorem 2.1** ([9]) *Let  $X$  be a Banach space,  $\Phi_1 : X \rightarrow \mathcal{P}_{cl,cv,bd}(X)$  and  $\Phi_2 : X \rightarrow \mathcal{P}_{cp,cv}(X)$  be two multi-valued operators satisfying:*

- (a)  $\Phi_1$  is a contraction,
- (b)  $\Phi_2$  is u.s.c. and completely continuous.

Then either

- (i) the operator inclusion  $\lambda x \in \Phi_1 x + \Phi_2 x$  has a solution for  $\lambda = 1$ , or
- (ii) the set  $G = \{x \in X : \lambda x \in \Phi_1 x + \Phi_2 x, \lambda > 1\}$  is unbounded.

**Lemma 2.4** ([10]) *Let  $v, w : [0, b] \rightarrow [0, \infty)$  be continuous functions. If  $w$  is nondecreasing and there are constants  $\theta > 0, 0 < \alpha < 1$  such that*

$$v(t) \leq w(t) + \theta \int_0^t \frac{v(s)}{(t-s)^{1-\alpha}} ds, \quad t \in J,$$

then

$$v(t) \leq e^{\frac{\theta^n \Gamma(\alpha)^n t^{n\alpha}}{\Gamma(n\alpha)}} \sum_{j=0}^{n-1} \left(\frac{\theta b^\alpha}{\alpha}\right)^j w(t)$$

for every  $t \in J$  and every  $n \in \mathbb{N}$  such that  $n\alpha > 1$  and  $\Gamma(\cdot)$  is the Gamma function.

### 3 Main result

Let  $J_1 = (-\infty, b]$ . First, we present the definition of the mild solution of problem (1.1)-(1.3).

**Definition 3.1** A stochastic process  $x : J_1 \times \Omega \rightarrow H$  is called a mild solution of problem (1.1)-(1.3) if

- (i)  $x(t)$  is measurable and  $\mathcal{F}_t$ -adapted for each  $t \geq 0$ ,
- (ii)  $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$ ,  $k = 1, 2, \dots, m$ ,
- (iii)  $x(t) \in H$  has càdlàg paths on  $t \in J$  a.e. and there exists a function  $\sigma \in N_{F,x}$  such that

$$\begin{aligned}
 x(t) = & S(t)[\phi(0) - g(0, \phi, 0)] + g\left(t, x_t, \int_0^t a(s, x_s) ds\right) \\
 & + \int_0^t AS(t-s)g\left(s, x_s, \int_0^s a(s, \tau, x_\tau) d\tau\right) ds + \int_0^t AS(t-s)f(s, x_s) ds \\
 & + \int_0^t S(t-s)\sigma(s) dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(x(t_k^-)), \quad t \in J,
 \end{aligned}$$

- (iv)  $x_0(\cdot) = \phi \in L^2(\Omega, \mathcal{B}_h)$  on  $J_0 = (-\infty, 0]$  satisfies  $\|\phi\|_{\mathcal{B}_h}^2 < \infty$ .

Now, we assume the following hypotheses:

- (H1)  $A$  is the infinitesimal generator of a compact analytic resolvent operator  $S(t)$ ,  $t \geq 0$ , in the Hilbert space  $H$  and there exist positive constants  $M$  and  $M_1$  such that

$$\|S(t)\|^2 \leq M, \quad \|A^{-\beta}\| \leq M_1, \quad t \in J.$$

- (H2)  $a : D \times \mathcal{B}_h \rightarrow H$ ,  $D = \{(t, s) \in J \times J : t \geq s\}$  is a continuous function and there exists a constant  $M_a$  such that

$$E \left| \int_0^t [a(t, s, x) - a(t, s, y)] ds \right|^2 \leq M_a \|x - y\|_{\mathcal{B}_h}^2 \quad \text{for all } t \in J, x, y \in \mathcal{B}_h.$$

- (H3) There exist constants  $0 < \beta < 1$  and  $M_g$  such that  $g$  is  $H_\beta$ -valued,  $(-A)^\beta g$  is continuous and

$$E|(-A)^\beta g(t, x_1, y_1) - (-A)^\beta g(t, x_2, y_2)|^2 \leq M_g [\|x_1 - x_2\|_{\mathcal{B}_h}^2 + E|y_1 - y_2|^2].$$

- (H4) The function  $f : J \times \mathcal{B}_h \rightarrow H$  satisfies the following conditions:

- (i)  $t \mapsto f(t, s)$  is measurable for each  $x \in \mathcal{B}_h$ ;
- (ii)  $x \mapsto f(t, x)$  is continuous for almost all  $t \in J$ ;
- (iii) There exists a constant  $M_f$  such that

$$E|(-A)^\beta f(t, x) - (-A)^\beta f(t, y)|^2 \leq M_f \|x - y\|_{\mathcal{B}_h}^2$$

for all  $x, y \in \mathcal{B}_h$ ,  $t \in J$  and

$$E|f(t, x)|^2 \leq p(t)\psi(\|x\|_{\mathcal{B}_h}^2)$$

for almost all  $t \in J$ , where  $p \in L^1(J, \mathbb{R})$ ,  $\psi : \mathbb{R}_+ \rightarrow (0, \infty)$  is continuous and increasing with

$$\begin{aligned}
 \int_0^b \frac{1}{\mu(s)} ds & \leq \int_{B_0 k_1}^\infty \frac{1}{\psi(s)} ds, \\
 \bar{\mu}(t) & = B_0 k_3 p(t),
 \end{aligned}$$

$$\begin{aligned}
 k_1 &= \frac{4\|\phi\|_{\mathcal{B}_h}^2 + l^2 F}{1 - 96l^2 \|(-A)^{-\beta}\|^2 M_g(1 + 2M_a)}, \\
 k_2 &= \frac{96bl^2 M_g(1 + 2M_a)c_{1-\beta}^2}{1 - 96l^2 \|(-A)^{-\beta}\|^2 M_g(1 + 2M_a)}, \\
 k_3 &= \frac{48Mb l^2}{1 - 96l^2 \|(-A)^{-\beta}\|^2 M_g(1 + 2M_a)}, \\
 L_0 &= 3l^2 \left[ M_g(1 + M_a) \left( \|(-A)^{-\beta}\|^2 + \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right) + M_f \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right] < 1, \\
 B_0 &= e^{k_2^\beta \Gamma(\beta)^n b^{n\beta} / \Gamma(n\beta)} \sum_{j=1}^{n-1} \left( \frac{k_2 b^\beta}{\beta} \right)^j, \\
 c_1 &= b^2 \sup_{(t,s) \in D} a^2(t, s, 0), \quad c_2 = \|(-A)^\beta\|^2 \sup_{t \in J} \|g(t, 0, 0)\|^2
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{F} &= 4M|\phi(0)|^2 + 96(M + \|(-A)^{-\beta}\|^2)c_2 + 192\|(-A)^{-\beta}\|^2 M_g c_1 \\
 &\quad + \frac{192b^{2\beta} C_{1-\beta}^2}{2\beta - 1} (c_2 + 2M_g c_1) + 48M\|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} b^2 \text{Tr}(Q) \\
 &\quad + 48Mm^2 \sum_{k=1}^m d_k + 96M\|(-A)^{-\beta}\|^2 M_g \|\phi\|_{\mathcal{B}_h}^2.
 \end{aligned}$$

(H5) The multi-valued mapping  $F : J \times \mathcal{B}_h \rightarrow \mathcal{P}_{bd,cl,cv}(L(K, H))$  is an  $L^2$ -Carathéodory function that satisfies the following conditions:

- (i) For each  $t \in J$ , the function  $F(t, \cdot) : \mathcal{B}_h \rightarrow \mathcal{P}_{bd,cl,cv}(L(K, H))$  is u.s.c. and for each fixed  $x \in \mathcal{B}_h$ , the function  $F(\cdot, x)$  is measurable. For each  $x \in \mathcal{B}_h$ , the set

$$N_{F,x} = \{ \sigma \in L^2(K, H) : \sigma(t) \in F(t, x) \text{ for a.e. } t \in J \}$$

is nonempty.

- (ii) There exists a positive function  $\mu \in L^1_{\text{loc}}(J, \mathbb{R}^+)$  such that

$$\|F(t, x)\|^2 = \sup_{\sigma \in F(t, x)} E|\sigma|^2 \leq \mu(t).$$

(H6)  $I_k \in C(H_\alpha, H_\alpha)$  and there exist positive constants  $d_k$  such that for each  $x \in H_\alpha$ ,

$$|I_k(x)|^2 \leq d_k, \quad k = 1, 2, \dots, m.$$

We consider the mapping  $\Phi : \mathcal{B}_h \rightarrow \mathcal{P}(\mathcal{B}_h)$  defined by

$$\Phi x(t) = \begin{cases} \phi(t), & t \in (-\infty, 0], \\ S(t)[\phi(0) - g(0, \phi, 0)] + g(t, x_t, \int_0^t a(t, s, x_s) ds) \\ \quad + \int_0^t AS(t-s)g(s, x_s, \int_0^s a(s, \tau, x_\tau) d\tau) ds \\ \quad + \int_0^t AS(t-s)f(s, x_s) ds + \int_0^t S(t-s)\sigma(s) dw(s) \\ \quad + \sum_{0 < t_k < t} S(t-t_k)I_k(x(t_k^-)), & t \in J, \end{cases}$$

where  $\sigma \in N_{F,x}$ . For each  $\phi \in \mathcal{B}_h$ , we define

$$\tilde{\phi}(t) = \begin{cases} \phi(t), & t \in (-\infty, 0], \\ S(t)\phi(0), & t \in J, \end{cases}$$

and then  $\tilde{\phi} \in \mathcal{B}_h$ . Let  $x(t) = y(t) + \tilde{\phi}(t)$ ,  $t \in (-\infty, b]$ . Then it is easy to see that  $x$  satisfies (1.1)-(1.3) if and only if  $y$  satisfies  $y_0 = 0$  and

$$\begin{aligned} y(t) = & -S(t)g(0, \phi, 0) + g\left(t, y_t + \tilde{\phi}_t, \int_0^t a(t, s, y_s + \tilde{\phi}_s) ds\right) \\ & + \int_0^t AS(t-s)g\left(s, y_s + \tilde{\phi}_s, \int_0^s a(s, \tau, y_\tau + \tilde{\phi}_\tau) d\tau\right) ds \\ & + \int_0^t AS(t-s)f(s, y_s + \tilde{\phi}_s) ds + \int_0^t S(t-s)\sigma(s) dw(s) \\ & + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)), \quad t \in J, \end{aligned}$$

where  $\sigma \in N_{F,y}$ . Let  $\mathcal{B}'_h = \{y \in \mathcal{B}_h : y_0 = 0 \in \mathcal{B}_h\}$ . For any  $y \in \mathcal{B}'_h$ ,

$$\begin{aligned} \|y\|_b &= \|y_0\|_{\mathcal{B}_h} + \sup_{0 \leq s \leq b} (E|y(s)|^2)^{\frac{1}{2}} \\ &= \sup_{0 \leq s \leq b} (E|y(s)|^2)^{\frac{1}{2}} \end{aligned}$$

and thus  $(\mathcal{B}'_h, \|\cdot\|_b)$  is a Banach space. Set  $\mathcal{B}_q = \{y \in \mathcal{B}'_h : \|y\|_b^2 \leq q\}$  for some  $q \geq 0$ . Then  $\mathcal{B}_q \subseteq \mathcal{B}'_h$  is uniformly bounded and for any  $y \in \mathcal{B}_q$ , from Lemma 2.2, we see that

$$\begin{aligned} \|y_t + \tilde{\phi}_t\|_{\mathcal{B}_h}^2 &\leq 2\|y_t\|_{\mathcal{B}_h}^2 + 2\|\tilde{\phi}_t\|_{\mathcal{B}_h}^2 \\ &\leq 4l^2 \sup_{0 \leq s \leq t} E|y(s)|^2 + 4\|y_0\|_{\mathcal{B}_h}^2 \\ &\quad + 4l^2 \sup_{0 \leq s \leq t} \|\tilde{\phi}(s)\|^2 + 4\|\tilde{\phi}_0\|_{\mathcal{B}_h}^2 \\ &\leq 4l^2 (q + M|\phi(0)|^2) + 4\|\tilde{\phi}\|_{\mathcal{B}_h}^2 \\ &:= q'. \end{aligned}$$

Define the operator  $\tilde{\Phi} : \mathcal{B}'_h \rightarrow \mathcal{P}(\mathcal{B}'_h)$  by

$$\tilde{\Phi}y(t) = \begin{cases} 0, & t \in (-\infty, 0], \\ -S(t)g(0, \phi, 0) + g\left(t, y_t + \tilde{\phi}_t, \int_0^t a(t, s, y_s + \tilde{\phi}_s) ds\right) \\ \quad + \int_0^t AS(t-s)g\left(s, y_s + \tilde{\phi}_s, \int_0^s a(s, \tau, y_\tau + \tilde{\phi}_\tau) d\tau\right) ds \\ \quad + \int_0^t AS(t-s)f(s, y_s + \tilde{\phi}_s) ds + \int_0^t S(t-s)\sigma(s) dw(s) \\ \quad + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)), & t \in J, \end{cases}$$

where  $\sigma \in N_{F,y}$ . Obviously, the operator  $\Phi$  has a fixed point is equivalent to proving that  $\tilde{\Phi}$  has a fixed point. Now, we decompose  $\tilde{\Phi}$  as  $\tilde{\Phi}_1 + \tilde{\Phi}_2$ , where

$$\begin{aligned} \tilde{\Phi}_1 y(t) = & -S(t)g(0, \phi, 0) + g\left(t, y_t + \tilde{\phi}_t, \int_0^t a(t, s, y_s + \tilde{\phi}_s) ds\right) \\ & + \int_0^t AS(t-s)g\left(s, y_s + \tilde{\phi}_s, \int_0^s a(s, \tau, y_\tau + \tilde{\phi}_\tau) d\tau\right) ds \\ & + \int_0^t AS(t-s)f(s, y_s + \tilde{\phi}_s) ds \end{aligned}$$

and

$$\tilde{\Phi}_2 y(t) = \int_0^s S(t-s)\sigma(s) dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)), \quad t \in J,$$

where  $\sigma \in N_{F,y}$ . In what follows, we show that the operators  $\tilde{\Phi}_1$  and  $\tilde{\Phi}_2$  satisfy all the conditions of Theorem 2.1.

**Lemma 3.1** *Assume that the assumptions (H1)-(H6) hold. Then  $\tilde{\Phi}_1$  is a contraction and  $\tilde{\Phi}_2$  is u.s.c. and completely continuous.*

*Proof* We give the proof in several steps:

Step 1.  $\tilde{\Phi}_1$  is a contraction.

Let  $u, v \in \mathcal{B}'_h$ . Then we have

$$\begin{aligned} & E|\tilde{\phi}_1 u(t) - \tilde{\phi}_1 v(t)|^2 \\ & \leq 3E\left|g\left(t, u_t + \tilde{\phi}_t, \int_0^t a(t, s, u_s + \tilde{\phi}_s) ds\right) - g\left(t, v_t + \tilde{\phi}_t, \int_0^t a(t, s, v_s + \tilde{\phi}_s) ds\right)\right|^2 \\ & \quad + 3bE\left(\int_0^t |AS(t-s)\left[g\left(s, u_s + \tilde{\phi}_s, \int_0^s a(s, \tau, u_\tau + \tilde{\phi}_\tau) d\tau\right) - g\left(s, v_s + \tilde{\phi}_s, \int_0^s a(s, \tau, v_\tau + \tilde{\phi}_\tau) d\tau\right)\right]|^2 ds\right) \\ & \quad + 3bE\left(\int_0^t |AS(t-s)[f(s, u_s + \tilde{\phi}_s) - f(s, v_s + \tilde{\phi}_s)]|^2 ds\right) \\ & \leq 3\|(-A)^{-\beta}\|^2 M_g (\|u_t - v_t\|_{\mathcal{B}'_h}^2 + M_a \|u_t - v_t\|_{\mathcal{B}'_h}^2) \\ & \quad + 3b \int_0^t \frac{C_{1-\beta}^2}{(t-s)^{2(1-\beta)}} M_g (\|u_s - v_s\|_{\mathcal{B}'_h}^2 + M_a \|u_s - v_s\|_{\mathcal{B}'_h}^2) ds \\ & \quad + 3b \int_0^t \frac{C_{1-\beta}^2}{(t-s)^{2(1-\beta)}} M_f \|u_s - v_s\|_{\mathcal{B}'_h}^2 ds \\ & \leq 3\|(-A)^{-\beta}\|^2 M_g (1 + M_a) \|u_t - v_t\|_{\mathcal{B}'_h}^2 \\ & \quad + 3M_g (1 + M_a) \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \|u_t - v_t\|_{\mathcal{B}'_h}^2 \\ & \quad + 3M_f \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \|u_t - v_t\|_{\mathcal{B}'_h}^2 \end{aligned}$$



$$\begin{aligned} &\leq 3 \left[ M_g(1 + M_a) \left( \|(-A)^{-\beta}\|^2 + \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right) + M_f \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right] \\ &\quad \times \left[ l^2 \sup_{s \in [0, b]} E|u(s) - v(s)|^2 + \|u_0\|_{\mathcal{B}_h}^2 + \|v_0\|_{\mathcal{B}_h}^2 \right] \\ &= 3l^2 \left[ M_g(1 + M_a) \left( \|(-A)^{-\beta}\|^2 + \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right) + M_f \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1} \right] \sup_{s \in [0, b]} E|u(s) - v(s)|^2 \\ &= L_0 \sup_{s \in [0, b]} E|u(s) - v(s)|^2, \end{aligned}$$

where  $L_0 = 3l^2 [M_g(1 + M_a)(\|(-A)^{-\beta}\|^2 + \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1}) + M_f \frac{(C_{1-\beta} b^\beta)^2}{2\beta - 1}] < 1$  and we have used the fact that  $\|u_0\|_{\mathcal{B}_h}^2 = 0$  and  $\|v_0\|_{\mathcal{B}_h}^2 = 0$ . Taking the supremum over  $t$ , we obtain

$$\|\tilde{\Phi}_1 u - \tilde{\Phi}_1 v\|_b^2 \leq L_0 \|u - v\|_b^2$$

and so  $\tilde{\Phi}_1$  is a contraction.

Now, we show that the operator  $\tilde{\Phi}_2$  is completely continuous.

Step 2.  $\tilde{\Phi}_2 y$  is convex for each  $y \in \mathcal{B}'_h$ .

In fact, if  $u_1, u_2 \in \tilde{\Phi}_2(y)$ , then there exist  $\sigma_1, \sigma_2 \in N_{F,y}$  such that

$$u_i(t) = \int_0^t S(t-s)\sigma_i(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-))$$

for  $i = 1, 2$  and  $t \in J$ . Let  $\lambda \in [0, 1]$ . Then for each  $t \in J$ , we have

$$\begin{aligned} \lambda u_1(t) + (1 - \lambda)u_2(t) &= \int_0^t S(t-s)[\lambda\sigma_1(s) + (1 - \lambda)\sigma_2(s)]dw(s) \\ &\quad + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)). \end{aligned}$$

Since  $N_{F,y}$  is convex (because  $F$  has convex values), we obtain

$$\lambda u_1(t) + (1 - \lambda)u_2(t) \in \tilde{\Phi}_2(y).$$

Step 3.  $\tilde{\Phi}_2$  maps bounded sets into bounded sets in  $\mathcal{B}'_h$ .

It is enough to show that there exists a positive constant  $\Lambda$  such that for each  $u \in \tilde{\Phi}_2 y$ ,  $y \in \mathcal{B}_q = \{y \in \mathcal{B}'_h : \|y\|_b \leq q\}$  one has  $\|u\|_b \leq \Lambda$ . If  $u \in \tilde{\Phi}_2(y)$ , there exists  $\sigma \in N_{F,y}$  such that for each  $t \in J$

$$u(t) = \int_0^t S(t-s)\sigma(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-))$$

and so

$$\begin{aligned} E|u(t)|^2 &= E \left| \int_0^t S(t-s)\sigma(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right|^2 \\ &\leq 2E \left| \int_0^t S(t-s)\sigma(s)dw(s) \right|^2 + 2E \left| \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right|^2 \end{aligned}$$

$$\begin{aligned} &\leq 2 \operatorname{Tr}(Q)Mb \int_0^b \mu(s) ds + 2Mm^2 \sum_{k=1}^m d_k \\ &\leq 2 \operatorname{Tr}(Q)Mb^2 \|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} + 2Mm^2 \sum_{k=1}^m d_k \\ &:= \Lambda. \end{aligned}$$

Thus, for each  $y \in \mathcal{B}'_h$ , we get  $\|u\|_b^2 \leq \Lambda$ .

Step 4.  $\tilde{\Phi}_2$  maps bounded sets into equicontinuous sets of  $\mathcal{B}'_h$ .

Let  $0 < \tau_1 < \tau_2 \leq b$ . For each  $y \in \mathcal{B}_q = \{y \in \mathcal{B}'_h : \|y\|_b \leq q\}$  and  $u \in \tilde{\Phi}_2(y)$ . Let  $\tau_1, \tau_2 \in J \setminus \{t_1, t_2, \dots, t_m\}$ . Then there exists  $\sigma \in N_{F,y}$  such that for each  $t \in J$ ,

$$u(t) = \int_0^t S(t-s)\sigma(s) dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)).$$

Thus we have

$$\begin{aligned} &E|u(\tau_2) - u(\tau_1)|^2 \\ &= E \left| \int_0^{\tau_2} S(\tau_2-s)\sigma(s) dw(s) + \sum_{0 < t_k < \tau_2} S(\tau_2-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right. \\ &\quad \left. - \int_0^{\tau_1} S(\tau_1-s)\sigma(s) dw(s) - \sum_{0 < t_k < \tau_1} S(\tau_1-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right|^2 \\ &\leq 2E \left| \int_0^{\tau_1-\varepsilon} (S(\tau_2-s)\sigma(s) - S(\tau_1-s)\sigma(s)) dw(s) \right. \\ &\quad \left. + \int_{\tau_1-\varepsilon}^{\tau_1} (S(\tau_2-s)\sigma(s) - S(\tau_1-s)\sigma(s)) dw(s) + \int_{\tau_1}^{\tau_2} S(\tau_2-s)\sigma(s) dw(s) \right|^2 \\ &\quad + 2E \left| \sum_{0 < t_k < \tau_1} [S(\tau_2-t_k) - S(\tau_1-t_k)]I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right. \\ &\quad \left. + \sum_{\tau_1 < t_k < \tau_2} S(\tau_2-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right|^2 \\ &\leq 6\varepsilon \operatorname{Tr}(Q) \int_0^{\tau_1-\varepsilon} \mu(s) \|S(\tau_2-s) - S(\tau_1-s)\|^2 ds \\ &\quad + 6\varepsilon \operatorname{Tr}(Q) \int_{\tau_1-\varepsilon}^{\tau_1} \mu(s) \|S(\tau_2-s) - S(\tau_1-s)\|^2 ds \\ &\quad + 6(\tau_2 - \tau_1) \operatorname{Tr}(Q) \int_{\tau_1}^{\tau_2} \mu(s) \|S(\tau_2-s)\|^2 ds \\ &\quad + 4m^2 \sum_{0 < t_k < \tau_1} \|S(\tau_2-s) - S(\tau_1-s)\|^2 d_k \\ &\quad + 4m^2 M \sum_{\tau_1 < t_k < \tau_2} d_k. \end{aligned}$$

The right-hand side of the above inequality tends to zero as  $\tau_1 \rightarrow \tau_2$  with  $\varepsilon$  sufficiently small, since  $S(t)$  is strongly continuous and the compactness of  $S(t)$  for  $t > 0$  implies the

continuity in the uniform operator topology. Thus, the set  $\{\tilde{\Phi}_2 y : y \in \mathcal{B}_q\}$  is equicontinuous. Here we consider the case  $0 < \tau_1 < \tau_2 \leq b$ , since the case  $\tau_1 < \tau_2 \leq 0$  or  $\tau_1 \leq 0 \leq \tau_2 \leq b$  is simple.

Step 5.  $\tilde{\Phi}_2$  maps  $\mathcal{B}_q$  into a precompact set in  $H$ .

Let  $0 < t \leq b$  and  $0 < \varepsilon < t$ . For  $y \in \mathcal{B}_q$  and  $u \in \tilde{\Phi}_2(y)$ , there exists  $\sigma \in N_{F,y}$  such that

$$u(t) = \int_0^{t-\varepsilon} S(t-s)\sigma(s)dw(s) + \int_{t-\varepsilon}^t S(t-s)\sigma(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)).$$

Define

$$u_\varepsilon(t) = S(\varepsilon) \int_0^{t-\varepsilon} S(t-\varepsilon-s)\sigma(s)dw(s) + \sum_{0 < t_k < t-\varepsilon} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)).$$

Since  $S(t)$  is a compact operator, the set  $V_\varepsilon(t) = \{u_\varepsilon(t) : u \in \tilde{\Phi}_2(\mathcal{B}_q)\}$  is relatively compact in  $H$  for each  $\varepsilon$ ,  $0 < \varepsilon < t$ . Moreover,

$$\begin{aligned} & E|u(t) - u_\varepsilon(t)|^2 \\ &= E \left| \int_0^{t-\varepsilon} S(t-s)\sigma(s)dw(s) + \int_{t-\varepsilon}^t S(t-s)\sigma(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) - S(\varepsilon) \int_0^{t-\varepsilon} S(t-\varepsilon-s)\sigma(s)dw(s) - \sum_{0 < t_k < t-\varepsilon} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \right|^2 \\ &\leq 4Mb \operatorname{Tr}(Q)\varepsilon \|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} + 4m^2M \sum_{t-\varepsilon < t_k < t} d_k. \end{aligned}$$

Therefore letting  $\varepsilon \rightarrow 0$ , we can see that there are relative compact sets arbitrarily close to the set  $\{u(t) : u \in \tilde{\Phi}_2(\mathcal{B}_q)\}$ . Thus, the set  $\{u(t) : u \in \tilde{\Phi}_2(\mathcal{B}_q)\}$  is relatively compact in  $H$ . Hence, the Arzelá-Ascoli theorem shows that  $\tilde{\Phi}_2$  is a compact multi-valued mapping.

Step 6.  $\tilde{\Phi}_2$  has a closed graph.

Let  $y_n \rightarrow y_*$ ,  $u_n \in \tilde{\Phi}_2(y_n)$  and  $u_n \rightarrow u_*$ . We prove that  $u_* \in \tilde{\Phi}_2(y_*)$ .

Indeed,  $u_n \in \tilde{\Phi}_2(y_n)$  means that there exists  $\sigma_n \in N_{F,y_n}$  such that

$$u_n(t) = \int_0^t S(t-s)\sigma_n(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y_n(t_k^-) + \tilde{\phi}(t_k^-)), \quad t \in J.$$

Thus we must prove that there exists  $\sigma_* \in N_{F,y_*}$  such that

$$u_*(t) = \int_0^t S(t-s)\sigma_*(s)dw(s) + \sum_{0 < t_k < t} S(t-t_k)I_k(y_*(t_k^-) + \tilde{\phi}(t_k^-)), \quad t \in J.$$

Since  $I_k, k = 1, 2, \dots, m$ , are continuous, we see that

$$\left\| \sum_{0 < t_k < t} S(t - t_k) I_k(y_n(t_k^-) + \tilde{\phi}(t_k^-)) - \sum_{0 < t_k < t} S(t - t_k) I_k(y_*(t_k^-) + \tilde{\phi}(t_k^-)) \right\|_b^2 \rightarrow 0$$

as  $n \rightarrow \infty$ . Consider the linear continuous operator  $\Gamma : L^2(J, H) \rightarrow C(J, H)$  with  $\Gamma(\sigma)(t) = \int_0^t S(t - s)\sigma(s) dw(s)$ , where  $\sigma \in N_{F,y}$ . From Lemma 2.3, it follows that  $\Gamma \circ N_F$  is a closed graph operator. Moreover, we have

$$u_n(t) - \sum_{0 < t_k < t} S(t - t_k) I_k(y_n(t_k^-) + \tilde{\phi}(t_k^-)) \in \Gamma(N_{F,y_n}).$$

Since  $y_n \rightarrow y_*$ , from Lemma 2.3, we obtain

$$u_*(t) - \sum_{0 < t_k < t} S(t - t_k) I_k(y_*(t_k^-) + \tilde{\phi}(t_k^-)) \in \Gamma(N_{F,y_*}).$$

That is, there exists a  $\sigma_* \in N_{F,y_*}$  such that

$$\begin{aligned} u_*(t) - \sum_{0 < t_k < t} S(t - t_k) I_k(y_*(t_k^-) + \tilde{\phi}(t_k^-)) &= \Gamma(\sigma_*(t)) \\ &= \int_0^t S(t - s)\sigma_*(s) dw(s). \end{aligned}$$

Therefore  $\tilde{\Phi}_2$  has a closed graph and  $\tilde{\Phi}_2$  is u.s.c. This completes the proof. □

**Lemma 3.2** *Assume that the assumptions (H1)-(H2) hold. Then there exists a constant  $K > 0$  such that  $\|y_t + \tilde{\phi}_t\|_{\mathcal{B}_t}^2 \leq K$  for all  $t \in J$ , where  $K$  is depends only on  $b$  and the functions  $\psi$  and  $\bar{\mu}$ .*

*Proof* Let  $y$  be a possible solution of  $y \in \lambda \tilde{\Phi}(y)$  for some  $0 < \lambda < 1$ . Then there exists  $\sigma \in N_{F,y}$  such that for  $t \in J$  we have

$$\begin{aligned} y(t) &= -\lambda S(t)g(0, \phi, 0) + \lambda g\left(t, y_t + \tilde{\phi}_t, \int_0^t a(t, s, y_s + \tilde{\phi}_s) ds\right) \\ &\quad + \lambda \int_0^t AS(t - s)g\left(s, y_s + \tilde{\phi}_s, \int_0^s a(s, \tau, y_\tau + \tilde{\phi}_\tau) d\tau\right) ds \\ &\quad + \lambda \int_0^t S(t - s)f(s, y_s + \tilde{\phi}_s) ds + \int_0^t S(t - s)\sigma(s) dw(s) \\ &\quad + \lambda \sum_{0 < t_k < t} S(t - t_k) I_k(y(t_k^-) + \tilde{\phi}(t_k^-)). \end{aligned}$$

Then, by the assumptions, we deduce that

$$\begin{aligned} E|y(t)|^2 &\leq E\left| -S(t)g(0, \phi, 0) + g\left(t, y_t + \tilde{\phi}_t, \int_0^t a(t, s, y_s + \tilde{\phi}_s) ds\right) \right. \\ &\quad \left. + \int_0^t AS(t - s)g\left(s, y_s + \tilde{\phi}_s, \int_0^s a(s, \tau, y_\tau + \tilde{\phi}_\tau) d\tau\right) ds \right. \end{aligned}$$

$$\begin{aligned}
 & + \int_0^t S(t-s)f(s, y_s + \tilde{\phi}_s) ds + \int_0^t S(t-s)\sigma(s) dw(s) \\
 & + \sum_{0 < t_k < t} S(t-t_k)I_k(y(t_k^-) + \tilde{\phi}(t_k^-)) \Big|^2 \\
 \leq & 12 \left\{ 2M(\|(-A)^{-\beta}\|^2 M_g \|\phi\|_{\mathcal{B}_h}^2 + c_2) \right. \\
 & + 2\|(-A)^{-\beta}\|^2 [M_g(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 + 2M_a \|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 + 2c_1) + c_2] \\
 & + 2b \int_0^t \frac{c_{1-\beta}^2}{(t-s)^{2(1-\beta)}} [M_g(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 + 2M_a \|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 + 2c_1) + c_2] ds \\
 & \left. + Mb \int_0^t p(s)\psi(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2) ds + M\|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} b^2 \text{Tr}(Q) + Mm^2 \sum_{k=1}^m d_k \right\} \\
 = & 24(M + \|(-A)^{-\beta}\|^2)c_2 + 48\|(-A)^{-\beta}\|^2 M_g c_1 + \frac{48b^{2\beta} c_{1-\beta}^2}{2\beta - 1}(c_2 + 2M_g c_1) \\
 & + 12M\|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} b^2 \text{Tr}(Q) + 12Mm^2 \sum_{k=1}^m d_k + 24M\|(-A)^{-\beta}\|^2 M_g \|\phi\|_{\mathcal{B}_h}^2 \\
 & + 24\|(-A)^{-\beta}\|^2 M_g(1 + 2M_a)\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 \\
 & + 24bM_g(1 + 2M_a)c_{1-\beta}^2 \int_0^t \frac{\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2}{(t-s)^{2(1-\beta)}} ds \\
 & + 12Mb \int_0^t p(s)\psi(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2) ds.
 \end{aligned}$$

From Lemma 2.2 we see that

$$\|y_t + \tilde{\phi}_t\|_{\mathcal{B}_h}^2 \leq 4l^2 \sup_{0 \leq s \leq t} E|y(s)|^2 + 4l^2 M|\tilde{\phi}(0)|^2 + 4\|\tilde{\phi}\|_{\mathcal{B}_h}^2.$$

Thus, for any  $t \in J$ , we have

$$\begin{aligned}
 & \|y_t + \tilde{\phi}_t\|_{\mathcal{B}_h}^2 \\
 \leq & 4l^2 M|\tilde{\phi}(0)|^2 + 4\|\tilde{\phi}\|_{\mathcal{B}_h}^2 + 96l^2(M + \|(-A)^{-\beta}\|^2)c_2 \\
 & + 192l^2\|(-A)^{-\beta}\|^2 M_g c_1 + \frac{192l^{2\beta} b^{2\beta} C_{1-\beta}^2}{2\beta - 1}(c_2 + 2M_g c_1) \\
 & + 48M\|\mu\|_{L^1_{\text{loc}}(J, \mathbb{R}^+)} b^2 l^2 \text{Tr}(Q) + 48Ml^2 m^2 \sum_{k=1}^m d_k \\
 & + 96Ml^2\|(-A)^{-\beta}\|^2 M_g \|\phi\|_{\mathcal{B}_h}^2 \\
 & + 96l^2\|(-A)^{-\beta}\|^2 M_g(1 + 2M_a)\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 \\
 & + 96bl^2 M_g(1 + 2M_a)C_{1-\beta}^2 \int_0^t \frac{\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2}{(t-s)^{2(1-\beta)}} ds \\
 & + 48Mbl^2 \int_0^t p(s)\psi(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2) ds
 \end{aligned}$$

$$\begin{aligned}
 &= 4\|\phi\|_{\mathcal{B}_h}^2 + l^2\mathcal{F} + 96l^2\|(-A)^{-\beta}\|^2M_g(1+2M_a)\sup_{0\leq s\leq t}\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2 \\
 &\quad + 96bl^2M_g(1+2M_a)C_{1-\beta}^2\int_0^t\frac{\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2}{(t-s)^{2(1-\beta)}}ds \\
 &\quad + 48Mbl^2\int_0^t p(s)\psi(\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2)ds.
 \end{aligned}$$

Let  $v(t) = \sup_{0\leq s\leq t}\|y_s + \tilde{\phi}_s\|_{\mathcal{B}_h}^2$ . Then the function  $v(t)$  is nondecreasing in  $J$ . Thus, we obtain

$$\begin{aligned}
 v(t) &\leq 4\|\phi\|_{\mathcal{B}_h}^2 + l^2\mathcal{F} + 96l^2\|(-A)^{-\beta}\|^2M_g(1+2M_a)v(t) \\
 &\quad + 96bl^2M_g(1+2M_a)C_{1-\beta}^2\int_0^t\frac{v(s)}{(t-s)^{2(1-\beta)}}ds \\
 &\quad + 48Mbl^2\int_0^t p(s)\psi(v(s))ds.
 \end{aligned}$$

From this we derive that

$$\begin{aligned}
 v(t) &\leq \frac{4\|\phi\|_{\mathcal{B}_h}^2 + l^2\mathcal{F}}{1 - 96l^2\|(-A)^{-\beta}\|^2M_g(1+2M_a)} \\
 &\quad + \frac{96bl^2M_g(1+2M_a)C_{1-\beta}^2}{1 - 96l^2\|(-A)^{-\beta}\|^2M_g(1+2M_a)}\int_0^t\frac{v(s)}{(t-s)^{1-\beta}}ds \\
 &\quad + \frac{48Mbl^2}{1 - 96l^2\|(-A)^{-\beta}\|^2M_g(1+2M_a)}\int_0^t p(s)\psi(v(s))ds \\
 &\leq k_1 + k_2\int_0^t\frac{v(s)}{(t-s)^{1-\beta}}ds + k_3\int_0^t p(s)\psi(v(s))ds.
 \end{aligned}$$

By Lemma 2.4, we get

$$v(t) \leq B_0\left(k_1 + k_3\int_0^t p(s)\psi(v(s))ds\right),$$

where

$$B_0 = e^{k_2^{\frac{n}{\beta}}\Gamma(\beta)^n b^{n\beta}/\Gamma(n\beta)}\sum_{j=1}^{n-1}\left(\frac{k_2 b^\beta}{\beta}\right)^j.$$

Let us take the right-hand side of the above inequality as  $\mu(t)$ . Then  $\mu(0) = B_0k_1$ ,  $v(t) \leq \mu(t)$ ,  $t \in J$  and

$$\mu'(t) \leq B_0k_3p(t)\psi(v(t)).$$

Since  $\psi$  is nondecreasing, we have

$$\begin{aligned}
 \mu'(t) &\leq B_0k_3p(t)\psi(\mu(t)) \\
 &= \bar{\mu}(t)\psi(\mu(t)).
 \end{aligned}$$

It follows that

$$\begin{aligned} \int_{\mu(0)}^{\mu(t)} \frac{1}{\psi(s)} ds &\leq \int_0^b \overline{\mu(s)} ds \\ &\leq \int_{B_0 K_1}^{\infty} \frac{1}{\psi(s)} ds, \end{aligned}$$

which indicates that  $\mu(t) < \infty$ . Thus, there exists a constant  $K$  such that  $\mu(t) \leq K, t \in J$ . Furthermore, we see that  $\|y_t + \tilde{\phi}_t\|_{\mathcal{B}_h}^2 \leq \nu(t) \leq \mu(t) \leq K, t \in J$ .  $\square$

**Theorem 3.1** *Assume that the assumptions (H1)-(H6) hold. The problem (1.1)-(1.3) has at least one mild solution on  $J$ .*

*Proof* Let us take the set

$$G(\Phi) = \{x \in \mathcal{B}_h : x \in \lambda \Phi(x) \text{ for some } \lambda \in (0, 1)\}.$$

Then for any  $x \in G(\Phi)$ , we have

$$\|x_t\|_{\mathcal{B}_h}^2 = \|y_t + \tilde{\phi}_t\|_{\mathcal{B}_h}^2 \leq K, \quad t \in J,$$

where  $K > 0$  is a constant in Lemma 3.2. This show that  $G$  is bounded on  $J$ . Hence from Theorem 2.1 there exists a fixed point  $x(t)$  for  $\Phi$  on  $\mathcal{B}_h$ , which is a mild solution of (1.1)-(1.3) on  $J$ .  $\square$

#### 4 An example

As an application of Theorem 3.1, we consider the impulsive neutral stochastic functional integro-differential inclusion of the following form:

$$\begin{aligned} &\frac{\partial}{\partial t} \left( z(t, x) + g \left( t, z(t-h, x), \int_0^t a(t, s, z(s-h, x)) ds \right) \right) \\ &\in \frac{\partial^2}{\partial x^2} z(t, x) + (f(t, z(t-h, x)) + [Q_1(t, z(t-h, x)), Q_2(t, z(t-h, x))]) dw(t), \end{aligned} \quad (4.1)$$

$$0 \leq x \leq \pi, t \in J, t \neq t_k,$$

$$\Delta z(t_k, x) = z(t_k^+, x) - z(t_k^-, x) = I_k(z(t_k^-, x)), \quad k = 1, 2, \dots, m, \quad (4.2)$$

$$z(t, 0) = z(t, \pi) = 0, \quad t \in J, \quad (4.3)$$

$$z(t, x) = \rho(t, x), \quad -\infty < t \leq 0, 0 \leq x \leq \pi, \quad (4.4)$$

where  $J = [0, b], k = 1, 2, \dots, m, z(t_k^+, x) = \lim_{h \rightarrow 0^+} z(t_k + h, x), z(t_k^-, x) = \lim_{h \rightarrow 0^-} z(t_k + h, x), Q_1, Q_2 : J \times \mathbb{R} \rightarrow \mathbb{R}$  are two given functions and  $w(t)$  is a one-dimensional standard Wiener process. We assume that for each  $t \in J, Q_1(t, \cdot)$  is lower semicontinuous and  $Q_2(t, \cdot)$  is upper semicontinuous. Let  $J_1 = (-\infty, b]$  and  $H = L^2([0, \pi])$  with norm  $\|\cdot\|$ . Define  $A : H \rightarrow H$  by  $Av = v''$  with domain  $D(A) = \{v \in H : v, v'$  are absolutely continuous,  $v'' \in H, v(0) = v(\pi) = 0\}$ . Then

$$Av = \sum_{n=1}^{\infty} n^2 (v, v_n) v_n, \quad v \in D(A),$$

where  $v_n = \sqrt{\frac{2}{\pi}} \sin(ns)$ ,  $n = 1, 2, \dots$ , is the orthogonal set of eigenvectors in  $A$ . It is well known that  $A$  is the infinitesimal generator of an analytic semigroup  $S(t)$ ,  $t \geq 0$  in  $H$  given by

$$S(t)v = \sum_{n=1}^{\infty} e^{-n^2 t} (v, v_n) v_n, \quad v \in H.$$

For every  $v \in H$ ,  $(-A)^{-\frac{1}{2}} v = \sum_{n=1}^{\infty} \frac{1}{n} (v, v_n) v_n$  and  $\|(-A)^{-\frac{1}{2}}\| = 1$ . The operator  $(-A)^{\frac{1}{2}}$  is given by

$$(-A)^{\frac{1}{2}} v = \sum_{n=1}^{\infty} n (v, v_n) v_n$$

on the space  $D((-A)^{-\frac{1}{2}}) = \{v \in H : \sum_{n=1}^{\infty} n (v, v_n) v_n \in H\}$ . Since the analytic semigroup  $S(t)$  is compact [10], there exists a constant  $M > 0$  such that  $\|S(t)\| \leq M$  and satisfies (H1). Now, we give a special  $\mathcal{B}_h$ -space. Let  $h(s) = e^{2s}$ ,  $s < 0$ . Then  $l = \int_{-\infty}^0 h(s) ds = \frac{1}{2}$  and let

$$\|\varphi\|_{\mathcal{B}_h} = \int_{-\infty}^0 h(s) \sup_{s \leq \theta \leq 0} (E|\varphi(\theta)|^2)^{\frac{1}{2}} ds.$$

It follows from [5] that  $(\mathcal{B}_h, \|\cdot\|_{\mathcal{B}_h})$  is a Banach space. Hence for  $(t, \phi) \in [0, b] \times \mathcal{B}_h$ , let

$$\phi(\theta)x = \phi(\theta, x), \quad (\theta, x) \in (-\infty, 0] \times [0, \pi],$$

$$z(t)(x) = z(t, x)$$

and

$$F(t, \phi)(x) = [Q_1(t, \phi(\theta, x)), Q_2(t, \phi(\theta, x))], \quad -\infty < \theta \leq 0, x \in [0, \pi].$$

Then (4.1)-(4.4) can be rewritten as the abstract form as the system (1.1)-(1.3). If we assume that (H2)-(H6) are satisfied, then the system (4.1)-(4.4) has a mild solution on  $[0, b]$ .

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

**Author details**

<sup>1</sup>Department of Mathematics, Busan National University, Busan, 609-735, South Korea. <sup>2</sup>Department of Mathematics, Donggeui University, Busan, 614-714, South Korea.

Received: 11 September 2013 Accepted: 17 December 2013 Published: 15 Jan 2014

**References**

1. Chang, YK, Anguraj, A, Arjunan, MM: Existence results for impulsive neutral functional differential equations with infinite delay. *Nonlinear Anal. Hybrid Syst.* **2**, 209-218 (2008)
2. Park, JY, Balachandran, K, Annappoorani, N: Existence results for impulsive neutral functional integro-differential equations with infinite delay. *Nonlinear Anal.* **71**, 3152-3162 (2009)
3. Lin, A, Hu, L: Existence results for impulsive neutral stochastic functional integro-differential inclusions with nonlocal initial conditions. *Comput. Math. Appl.* **59**, 64-73 (2010)
4. Hu, L, Ren, Y: Existence results for impulsive neutral stochastic functional integro-differential equations with infinite delays. *Acta Appl. Math.* **111**, 303-317 (2010)



5. Pazy, A: Semigroups of Linear Operators and Applications to Partial Differential Equations. Applied Mathematical Sciences, vol. 44. Springer, New York (1983)
6. Hino, Y, Murakami, S, Naito, T: Functional Differential Equations with Infinite Delay. Lecture Notes in Mathematics, vol. 1473. Springer, Berlin (1991)
7. Chang, YK: Controllability of impulsive functional differential systems with infinite delay in Banach spaces. *Chaos Solitons Fractals* **33**, 1601-1609 (2007)
8. Losa, A, Opial, Z: Application of the Kakutani-Ky-Fan theorem in the theory of ordinary differential equations or noncompact acyclic-valued map. *Bull. Acad. Pol. Sci., Sér. Sci. Math. Astron. Phys.* **13**, 781-786 (1965)
9. Dhage, BC: Multi-valued mappings and fixed points I. *Nonlinear Funct. Anal. Appl.* **10**, 359-378 (2005)
10. Hernandez, E: Existence results for partial neutral functional integro-differential equations with unbounded delay. *J. Math. Anal. Appl.* **292**, 194-210 (2004)

10.1186/1687-1847-2014-17

**Cite this article as:** Park and Jeong: Existence results for impulsive neutral stochastic functional integro-differential inclusions with infinite delays. *Advances in Difference Equations* 2014, 2014:17

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---