

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 6-chloro-2-methyl-4-phenylquinoline-3-carboxylate


 J. Kalyana Sundar,^a S. Natarajan,^a S. Sarveswari,^b
 V. Vijayakumar,^b J. Suresh^c and P. L. Nilantha Lakshman^{d*}
^aDepartment of Physics, Madurai Kamaraj University, Madurai 625 021, India, ^bOrganic Chemistry Division, School of Advanced Sciences, VIT University, Vellore 632 014, India, ^cDepartment of Physics, The Madura College, Madurai 625 011, India, and ^dDepartment of Food Science and Technology, Faculty of Agriculture, University of Ruhuna, Mapalana, Kamburupitiya 81100, Sri Lanka
 Correspondence e-mail: nilanthalakshman@yahoo.co.uk

Received 1 December 2009; accepted 9 December 2009

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.100; data-to-parameter ratio = 13.3.

 In the title compound, $\text{C}_{18}\text{H}_{14}\text{ClNO}_2$, the quinoline ring system is planar (r.m.s. deviation = 0.032 Å) and the phenyl ring is twisted away from it by 57.5 (1)°. The crystal structure is stabilized by weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

 For the anti-tuberculosis activity of quinoline-2-carboxylic acid derivatives, see: Jain *et al.* (2005).


Experimental

Crystal data

 $\text{C}_{18}\text{H}_{14}\text{ClNO}_2$
 $M_r = 311.75$

 Monoclinic, $P2_1/n$
 $a = 10.828$ (5) Å
 $b = 7.535$ (4) Å
 $c = 18.829$ (5) Å
 $\beta = 94.369$ (5)°
 $V = 1531.8$ (12) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.26$ mm⁻¹
 $T = 293$ K
 $0.18 \times 0.16 \times 0.13$ mm

Data collection

 Nonius MACH-3 diffractometer
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.955$, $T_{\max} = 0.967$
 3320 measured reflections

 2682 independent reflections
 2309 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
 2 standard reflections every 2 min
 intensity decay: none

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.100$
 $S = 1.03$
 2682 reflections

 202 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.29$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C9}-\text{H9B}\cdots\text{Cg1}^i$	0.96	2.80	3.744 (3)	166

 Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$. Cg1 is the centroid of the C14-C19 ring.

 Data collection: *CAD-4 EXPRESS* (Enraf-Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

SN thanks the DST for the FIST programme and VV thanks the DST-India for funding through the Young Scientist-Fast Track Proposal.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: C12981).

References

- Enraf-Nonius (1994). *CAD-4 EXPRESS*. Enraf-Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1996). *XCAD4*. University of Marburg, Germany.
- Jain, R., Singh, P. P., Jain, M., Sachdeva, S., Misra, V., Kaul, C. L., Kaur, S., Vaitilingam, B., Nayyar, A. & Bhaskar, P. P. (2005). Indian Patent Appl. IN 2002DE00628.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351-359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148-155.

supporting information

Acta Cryst. (2010). E66, o169 [doi:10.1107/S1600536809052891]

Methyl 6-chloro-2-methyl-4-phenylquinoline-3-carboxylate

J. Kalyana Sundar, S. Natarajan, S. Sarveswari, V. Vijayakumar, J. Suresh and P. L. Nilantha Lakshman

S1. Comment

Quinoline-2-carboxylic acid derivatives are a class of important materials useful as anti-tuberculosis agents (Jain *et al.*, 2005). We report here the crystal structure of the title compound, a quinoline derivative.


The molecular structure of the title compound is shown in Fig.1. The quinoline ring system is planar (r.m.s. deviation is 0.032 Å). Due to phenyl substitution in the pyridine ring, the C4—C5 bond is longer [1.373 (2) Å] and the C3—C4 bond is shorter [1.434 (2) Å] than standard values for C=C (1.334 Å) and C_{sp}²—C_{sp}² (1.455 Å) bond lengths respectively. The phenyl ring is twisted out of the quinoline ring system by 57.5 (1)°. A weak C—H... π interaction involving the C14—C19 ring is observed.

S2. Experimental

A mixture of 2-amino-5-chlorobenzophenone (2.3 g, 0.01 mol) and methyacetate (1.2 g, 0.01 mmol) with 0.15 ml concentrated HCl taken in a beaker was subjected to microwave irradiation for about 6 min. After completion of the reaction (TLC), the reaction mixture was washed with saturated NaHCO₃ solution (10 ml), dried, washed with petroleum ether and recrystallized with chloroform (m.p. 134–135 °C).

S3. Refinement

H atoms were placed at calculated positions and allowed to ride on their carrier atoms with C—H = 0.93–0.96 Å and $U_{iso} = 1.2U_{eq}(C)$ for CH groups, and $1.5U_{eq}$ for CH₃ groups.

**Figure 1**

The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme.

Methyl 6-chloro-2-methyl-4-phenylquinoline-3-carboxylate

Crystal data

$C_{18}H_{14}ClNO_2$

$M_r = 311.75$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 10.828\ (5)\ \text{\AA}$

$b = 7.535\ (4)\ \text{\AA}$

$c = 18.829\ (5)\ \text{\AA}$

$\beta = 94.369\ (5)^\circ$

$V = 1531.8\ (12)\ \text{\AA}^3$

$Z = 4$

$F(000) = 648$

$D_x = 1.352\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71069\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 2\text{--}25^\circ$

$\mu = 0.26\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Block, colourless

$0.18 \times 0.16 \times 0.13\ \text{mm}$

Data collection

Nonius MACH-3

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω - 2θ scans

Absorption correction: ψ scan

(North *et al.*, 1968)

$T_{\min} = 0.955$, $T_{\max} = 0.967$

3320 measured reflections

2682 independent reflections

2309 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.1^\circ$

$h = 0\text{--}12$

$k = -1\text{--}8$

$l = -22\text{--}22$

2 standard reflections every 60 min

intensity decay: none

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.100$
 $S = 1.03$
 2682 reflections
 202 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0502P)^2 + 0.5537P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0132 (14)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C2	0.46859 (14)	0.2554 (2)	-0.02447 (8)	0.0362 (3)
C3	0.53086 (13)	0.25646 (19)	0.04444 (8)	0.0346 (3)
C4	0.46120 (13)	0.30371 (19)	0.10339 (8)	0.0339 (3)
C5	0.33707 (13)	0.33626 (19)	0.08882 (8)	0.0340 (3)
C6	0.28365 (13)	0.3408 (2)	0.01735 (8)	0.0358 (3)
C7	0.15150 (15)	0.3971 (3)	0.00113 (10)	0.0487 (4)
H7A	0.1384	0.4275	-0.0484	0.073*
H7B	0.1343	0.4985	0.0297	0.073*
H7C	0.0973	0.3015	0.0117	0.073*
C8	0.25152 (14)	0.3699 (2)	0.14669 (8)	0.0410 (4)
C9	0.0887 (2)	0.2382 (3)	0.20546 (13)	0.0727 (6)
H9A	0.1254	0.2793	0.2505	0.109*
H9B	0.0525	0.1233	0.2114	0.109*
H9C	0.0257	0.3200	0.1878	0.109*
C10	0.53354 (17)	0.2038 (2)	-0.08359 (9)	0.0485 (4)
H10	0.4928	0.2031	-0.1289	0.058*
C11	0.65448 (17)	0.1555 (2)	-0.07501 (10)	0.0529 (5)
H11	0.6968	0.1224	-0.1141	0.063*
C12	0.71432 (14)	0.1561 (2)	-0.00690 (11)	0.0472 (4)
C13	0.65654 (14)	0.2040 (2)	0.05190 (9)	0.0427 (4)
H13	0.6994	0.2022	0.0966	0.051*
C14	0.52259 (14)	0.3155 (2)	0.17690 (8)	0.0386 (4)
C15	0.62527 (15)	0.4262 (2)	0.19060 (9)	0.0451 (4)

H15	0.6545	0.4927	0.1538	0.054*
C16	0.68357 (17)	0.4376 (3)	0.25823 (10)	0.0548 (5)
H16	0.7522	0.5109	0.2667	0.066*
C17	0.6404 (2)	0.3406 (3)	0.31331 (10)	0.0608 (5)
H17	0.6805	0.3469	0.3587	0.073*
C18	0.5376 (2)	0.2341 (3)	0.30062 (10)	0.0608 (5)
H18	0.5072	0.1710	0.3380	0.073*
C19	0.47923 (17)	0.2203 (2)	0.23292 (9)	0.0489 (4)
H19	0.4106	0.1467	0.2249	0.059*
N1	0.34737 (12)	0.30270 (18)	-0.03732 (7)	0.0394 (3)
O1	0.24231 (13)	0.50480 (19)	0.17883 (8)	0.0677 (4)
O2	0.18315 (11)	0.22595 (16)	0.15517 (7)	0.0536 (3)
Cl1	0.86967 (4)	0.09440 (7)	0.00332 (4)	0.0739 (2)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C2	0.0368 (8)	0.0318 (8)	0.0408 (8)	0.0000 (6)	0.0077 (6)	0.0017 (6)
C3	0.0317 (7)	0.0288 (7)	0.0439 (8)	0.0001 (6)	0.0060 (6)	0.0023 (6)
C4	0.0340 (7)	0.0294 (7)	0.0384 (8)	-0.0006 (6)	0.0036 (6)	0.0018 (6)
C5	0.0322 (7)	0.0302 (7)	0.0398 (8)	0.0004 (6)	0.0048 (6)	-0.0011 (6)
C6	0.0322 (7)	0.0312 (8)	0.0439 (8)	-0.0005 (6)	0.0021 (6)	-0.0014 (6)
C7	0.0339 (8)	0.0538 (11)	0.0572 (10)	0.0064 (7)	-0.0029 (7)	-0.0010 (8)
C8	0.0369 (8)	0.0427 (9)	0.0440 (9)	0.0030 (7)	0.0071 (6)	-0.0028 (7)
C9	0.0644 (13)	0.0759 (15)	0.0835 (15)	0.0002 (11)	0.0421 (11)	0.0110 (12)
C10	0.0536 (10)	0.0489 (10)	0.0445 (9)	0.0026 (8)	0.0140 (7)	-0.0021 (8)
C11	0.0532 (10)	0.0462 (10)	0.0629 (11)	0.0032 (8)	0.0287 (9)	-0.0036 (8)
C12	0.0329 (8)	0.0321 (8)	0.0786 (12)	0.0026 (6)	0.0173 (8)	0.0004 (8)
C13	0.0332 (8)	0.0362 (8)	0.0588 (10)	0.0016 (6)	0.0042 (7)	0.0026 (7)
C14	0.0359 (8)	0.0393 (9)	0.0402 (8)	0.0030 (6)	0.0005 (6)	0.0016 (7)
C15	0.0432 (9)	0.0459 (9)	0.0456 (9)	-0.0035 (7)	0.0002 (7)	0.0039 (7)
C16	0.0505 (10)	0.0550 (11)	0.0568 (10)	-0.0077 (8)	-0.0097 (8)	-0.0015 (9)
C17	0.0713 (13)	0.0643 (12)	0.0440 (10)	-0.0029 (10)	-0.0143 (9)	0.0021 (9)
C18	0.0739 (13)	0.0627 (12)	0.0445 (10)	-0.0086 (10)	-0.0030 (9)	0.0138 (9)
C19	0.0511 (10)	0.0488 (10)	0.0461 (9)	-0.0072 (8)	-0.0004 (7)	0.0076 (8)
N1	0.0374 (7)	0.0410 (7)	0.0396 (7)	0.0020 (6)	0.0013 (5)	-0.0003 (6)
O1	0.0707 (9)	0.0562 (8)	0.0804 (10)	-0.0041 (7)	0.0339 (7)	-0.0240 (7)
O2	0.0524 (7)	0.0487 (7)	0.0629 (8)	-0.0063 (6)	0.0258 (6)	-0.0024 (6)
Cl1	0.0349 (3)	0.0598 (3)	0.1295 (5)	0.0111 (2)	0.0227 (3)	-0.0040 (3)

Geometric parameters (Å, °)

C2—N1	1.364 (2)	C9—H9C	0.96
C2—C10	1.416 (2)	C10—C11	1.357 (3)
C2—C3	1.416 (2)	C10—H10	0.93
C3—C13	1.414 (2)	C11—C12	1.392 (3)
C3—C4	1.434 (2)	C11—H11	0.93
C4—C5	1.373 (2)	C12—C13	1.361 (2)

C4—C14	1.492 (2)	C12—C11	1.7418 (16)
C5—C6	1.424 (2)	C13—H13	0.93
C5—C8	1.505 (2)	C14—C19	1.387 (2)
C6—N1	1.3139 (19)	C14—C15	1.398 (2)
C6—C7	1.501 (2)	C15—C16	1.380 (2)
C7—H7A	0.96	C15—H15	0.93
C7—H7B	0.96	C16—C17	1.379 (3)
C7—H7C	0.96	C16—H16	0.93
C8—O1	1.191 (2)	C17—C18	1.378 (3)
C8—O2	1.330 (2)	C17—H17	0.93
C9—O2	1.448 (2)	C18—C19	1.383 (3)
C9—H9A	0.96	C18—H18	0.93
C9—H9B	0.96	C19—H19	0.93
N1—C2—C10	117.46 (14)	C11—C10—H10	119.6
N1—C2—C3	123.08 (13)	C2—C10—H10	119.6
C10—C2—C3	119.45 (14)	C10—C11—C12	119.08 (15)
C13—C3—C2	118.49 (14)	C10—C11—H11	120.5
C13—C3—C4	123.48 (14)	C12—C11—H11	120.5
C2—C3—C4	117.97 (13)	C13—C12—C11	122.66 (15)
C5—C4—C3	117.02 (13)	C13—C12—C11	118.74 (15)
C5—C4—C14	122.34 (13)	C11—C12—C11	118.60 (13)
C3—C4—C14	120.64 (13)	C12—C13—C3	119.47 (16)
C4—C5—C6	120.93 (13)	C12—C13—H13	120.3
C4—C5—C8	122.19 (13)	C3—C13—H13	120.3
C6—C5—C8	116.88 (13)	C19—C14—C15	118.62 (15)
N1—C6—C5	122.37 (13)	C19—C14—C4	121.47 (14)
N1—C6—C7	116.85 (14)	C15—C14—C4	119.91 (14)
C5—C6—C7	120.75 (14)	C16—C15—C14	120.58 (16)
C6—C7—H7A	109.5	C16—C15—H15	119.7
C6—C7—H7B	109.5	C14—C15—H15	119.7
H7A—C7—H7B	109.5	C17—C16—C15	120.24 (17)
C6—C7—H7C	109.5	C17—C16—H16	119.9
H7A—C7—H7C	109.5	C15—C16—H16	119.9
H7B—C7—H7C	109.5	C18—C17—C16	119.57 (16)
O1—C8—O2	124.49 (15)	C18—C17—H17	120.2
O1—C8—C5	126.29 (15)	C16—C17—H17	120.2
O2—C8—C5	109.17 (13)	C17—C18—C19	120.69 (18)
O2—C9—H9A	109.5	C17—C18—H18	119.7
O2—C9—H9B	109.5	C19—C18—H18	119.7
H9A—C9—H9B	109.5	C18—C19—C14	120.28 (17)
O2—C9—H9C	109.5	C18—C19—H19	119.9
H9A—C9—H9C	109.5	C14—C19—H19	119.9
H9B—C9—H9C	109.5	C6—N1—C2	118.28 (13)
C11—C10—C2	120.85 (17)	C8—O2—C9	116.98 (15)
N1—C2—C3—C13	179.65 (14)	C10—C11—C12—C11	-179.82 (14)
C10—C2—C3—C13	-0.5 (2)	C11—C12—C13—C3	-0.3 (3)

N1—C2—C3—C4	2.2 (2)	C11—C12—C13—C3	179.26 (12)
C10—C2—C3—C4	-177.96 (14)	C2—C3—C13—C12	0.7 (2)
C13—C3—C4—C5	-174.20 (14)	C4—C3—C13—C12	177.98 (14)
C2—C3—C4—C5	3.1 (2)	C5—C4—C14—C19	54.6 (2)
C13—C3—C4—C14	5.6 (2)	C3—C4—C14—C19	-125.18 (17)
C2—C3—C4—C14	-177.17 (14)	C5—C4—C14—C15	-124.62 (17)
C3—C4—C5—C6	-6.4 (2)	C3—C4—C14—C15	55.6 (2)
C14—C4—C5—C6	173.85 (14)	C19—C14—C15—C16	1.2 (3)
C3—C4—C5—C8	174.07 (13)	C4—C14—C15—C16	-179.56 (16)
C14—C4—C5—C8	-5.7 (2)	C14—C15—C16—C17	-0.5 (3)
C4—C5—C6—N1	4.7 (2)	C15—C16—C17—C18	-0.9 (3)
C8—C5—C6—N1	-175.70 (14)	C16—C17—C18—C19	1.6 (3)
C4—C5—C6—C7	-172.83 (15)	C17—C18—C19—C14	-0.9 (3)
C8—C5—C6—C7	6.7 (2)	C15—C14—C19—C18	-0.6 (3)
C4—C5—C8—O1	77.7 (2)	C4—C14—C19—C18	-179.76 (17)
C6—C5—C8—O1	-101.9 (2)	C5—C6—N1—C2	0.7 (2)
C4—C5—C8—O2	-104.54 (17)	C7—C6—N1—C2	178.37 (14)
C6—C5—C8—O2	75.90 (17)	C10—C2—N1—C6	176.04 (14)
N1—C2—C10—C11	179.81 (16)	C3—C2—N1—C6	-4.1 (2)
C3—C2—C10—C11	0.0 (3)	O1—C8—O2—C9	2.1 (3)
C2—C10—C11—C12	0.4 (3)	C5—C8—O2—C9	-175.75 (15)
C10—C11—C12—C13	-0.2 (3)		

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C14—C19 ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C9—H9B...Cg1 ⁱ	0.96	2.80	3.744 (3)	166

Symmetry code: (i) $-x+1/2, y-1/2, -z+1/2$.