

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-[4-(Dimethylamino)benzylidene]-4-methylaniline

 R. K. Balachandar,^a S. Kalainathan,^a Shibu M Eappen^b and Jiban Podder^{c*}

^aCentre for Crystal Growth, School of Advanced Sciences, VIT University, Vellore 632 014, India, ^bSophisticated Test and Instrumentation Centre (STIC), Cochin University PO, Cochin 682 022, Kerala, and ^cDepartment of Physics, Bangladesh University of Engineering and Technology, Dhaka 1000, Bangladesh
Correspondence e-mail: jpodder59@gmail.com


Received 29 April 2013; accepted 7 May 2013

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.061; wR factor = 0.187; data-to-parameter ratio = 16.1.

The molecules of the title compound, $\text{C}_{16}\text{H}_{18}\text{N}_2$, exists in a *trans* conformation with respect to the $\text{C}=\text{N}$ bond [1.270 (3) Å]. The least-squares plane of the dimethylamino group makes a dihedral angle of 1.3 (2)° with the ring to which it is attached. The dihedral angle between the two aromatic rings is 11.70 (2)°. The crystal structure features weak $\text{C}-\text{H}\cdots\pi$ interactions.

Related literature

For the uses and biological importance of diketones, see: Xia *et al.* (2009); Shah *et al.* (1992); Ünver *et al.* (2004). For related structures, see: Fun *et al.* (2011); Khalaji & Simpson (2009).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{18}\text{N}_2$
 $M_r = 238.32$
Orthorhombic, *Pbca*

$a = 10.4814$ (10) Å
 $b = 8.0528$ (8) Å
 $c = 32.571$ (3) Å

$V = 2749.1$ (4) Å³
 $Z = 8$
Mo $K\alpha$ radiation

$\mu = 0.07$ mm⁻¹
 $T = 296$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2004)
 $T_{\min} = 0.980$, $T_{\max} = 0.987$

17121 measured reflections
2692 independent reflections
1737 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.187$
 $S = 1.08$
2692 reflections

167 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C11}-\text{H11}\cdots\text{Cg1}^i$	0.93	2.94	3.670 (2)	137

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SIR92 (Altomare *et al.*, 1993); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

The author acknowledges the STIC, Cochin 682 022, for the single-crystal XRD facility. The authors also thank Mr P. Narayanan and Dr K.Sethusankar, RKM Vivekananda College (Autonomous), Chennai 600 004, and VIT University for providing the excellent research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT6904).

References

- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
Bruker (2004). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
Fun, H.-K., Quah, C. K., Huang, C. & Yu, H. (2011). *Acta Cryst.* **E67**, o1273–o1274.
Khalaji, A. D. & Simpson, J. (2009). *Acta Cryst.* **E65**, o553.
Shah, S., Vyas, R. & Mehta, R. H. (1992). *J. Indian Chem. Soc.* **69**, 590–590.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Ünver, H., Karakas, A. & Elmali, A. (2004). *J. Mol. Struct.* **702**, 49–54.
Xia, D.-G., Ye, Y.-F. & Lei, K.-W. (2009). *Acta Cryst.* **E65**, o3168.

supplementary materials

Acta Cryst. (2013). E69, o905 [doi:10.1107/S160053681301249X]

N-[4-(Dimethylamino)benzylidene]-4-methylaniline

R. K. Balachandar, S. Kalainathan, Shibu M Eappen and Jiban Podder

Comment

Schiff bases are among the most useful ligands in coordination chemistry as they readily form stable complexes with most transition metals (Xia *et al.*, 2009). They are known to exhibit potent anti-bacterial, anti-convulsant, anti-inflammatory and anti-cancer activities (Shah *et al.*, 1992). In addition to that, they show Non-linear optical properties (Ünver *et al.*, 2004). Therefore, successful application of Schiff bases requires a careful study of their characteristics.

The title compound, C₁₆H₁₈N₂, exists in a *trans* configuration with respect to the C=N bond [1.270 (3) Å]. The N1=C8 bond length of 1.270 (3) Å is shorter than the N–C bond [1.413 (3) Å], indicating a typical imine double bond. The C–N–C angle is 120.6 (2)°. X-ray analysis confirms the molecular structure and atom connectivity as illustrated in Fig. 1.

The least square plane of the dimethylamine group has a dihedral angle of 1.31 (2)° with the phenyl ring (C9–C14), which shows that they are almost coplanar to each other. The dimethylamine group attached phenyl ring (C9–C14) forms a dihedral angle of 11.70 (2)° with the methyl group attached phenyl ring (C2–C7).

The crystal packing is stabilized by C11—H11⋯Cg1ⁱ inter-molecular interactions, where Cg1 is the center of gravity of (C8–C14) phenyl ring. The symmetry code is 1/2-*X*, -1/2+*Y*, *Z*.

Experimental


The title compound was synthesized by the reaction of *p*-dimethylaminobenzaldehyde (10 mmol, 1.14919 g) with *p*-toluidine (10 mmol, 1.0717 g) in ethanol (25 ml) under reflux condition for six hours. After filtering, drying the solid product was recrystallized from ethanol/THF (5:1 v/v). After five days yellow colour crystals were obtained which were suitable for X-ray diffraction studies.

Refinement


The positions of hydrogen atoms were localized from the difference electron density maps and their distances were geometrically constrained. The H atoms bound to the C atoms were treated as riding atoms, with d(C–H) = 0.93 Å and $U_{iso}(H) = 1.2U_{eq}(C)$ for aryl atoms; d(C–H) = 0.96 Å and $U_{iso}(H) = 1.5U_{eq}(C)$ for methyl group. The methyl groups were allowed to rotate but not to tip.

Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREF* (Bruker, 2004); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).


Figure 1

The molecular structure of the title compound with the atom numbering scheme, displacement ellipsoids are drawn at 30% probability level. H atoms are present as small spheres of arbitrary radius.


Figure 2

The crystal packing of the title compound, viewed down *a*-axis, showing C11—H11...Cg1ⁱ inter-molecular interactions. The H atoms not involved in the bonding have been excluded for clarity.

N-[4-(Dimethylamino)benzylidene]-4-methylaniline

Crystal data

C₁₆H₁₈N₂

M_r = 238.32

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

a = 10.4814 (10) Å

b = 8.0528 (8) Å

c = 32.571 (3) Å

V = 2749.1 (4) Å³

Z = 8

F(000) = 1024

D_x = 1.152 Mg m⁻³

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 1737 reflections

θ = 2.3–26.0°

μ = 0.07 mm⁻¹

T = 296 K

Block, yellow

0.30 × 0.25 × 0.20 mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and φ scan

Absorption correction: multi-scan

(*SADABS*; Bruker, 2004)

T_{min} = 0.980, *T_{max}* = 0.987

17121 measured reflections

2692 independent reflections

1737 reflections with *I* > 2σ(*I*)

R_{int} = 0.041

θ_{max} = 26.0°, θ_{min} = 2.3°

h = -12→12

k = -9→7

l = -40→40

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.061$

$wR(F^2) = 0.187$

$S = 1.08$

2692 reflections

167 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0814P)^2 + 0.8121P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0130 (19)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C6	0.0030 (3)	0.4888 (3)	0.18005 (8)	0.0768 (8)
H6	-0.0658	0.5293	0.1651	0.092*
C14	0.0373 (2)	0.2889 (2)	0.03147 (7)	0.0523 (6)
H14	-0.0182	0.3671	0.0425	0.063*
C13	0.0307 (2)	0.2534 (3)	-0.00941 (7)	0.0528 (6)
H13	-0.0297	0.3073	-0.0255	0.063*
C15	0.1898 (3)	-0.0172 (4)	-0.08750 (8)	0.0805 (8)
H15A	0.2757	0.0243	-0.0875	0.121*
H15B	0.1629	-0.0378	-0.1152	0.121*
H15C	0.1862	-0.1187	-0.0721	0.121*
C16	0.0139 (3)	0.1841 (4)	-0.09499 (8)	0.0809 (8)
H16A	-0.0705	0.1568	-0.0857	0.121*
H16B	0.0247	0.1468	-0.1228	0.121*
H16C	0.0258	0.3022	-0.0938	0.121*
C10	0.2053 (2)	0.0940 (2)	0.03889 (7)	0.0530 (6)
H10	0.2640	0.0380	0.0552	0.064*
C1	0.1192 (4)	0.5064 (5)	0.29002 (9)	0.1148 (13)
H1A	0.1367	0.6223	0.2939	0.172*
H1B	0.1863	0.4416	0.3021	0.172*
H1C	0.0395	0.4787	0.3029	0.172*
C3	0.2043 (3)	0.3782 (4)	0.22509 (9)	0.0862 (9)
H3	0.2737	0.3400	0.2402	0.103*
C7	0.0115 (3)	0.5248 (4)	0.22110 (9)	0.0831 (9)
H7	-0.0521	0.5886	0.2333	0.100*

C4	0.1978 (3)	0.3420 (4)	0.18370 (8)	0.0788 (8)
H4	0.2632	0.2823	0.1713	0.095*
C2	0.1112 (3)	0.4694 (3)	0.24474 (8)	0.0791 (8)
C9	0.1257 (2)	0.2105 (2)	0.05712 (7)	0.0482 (5)
C12	0.11276 (19)	0.1375 (2)	-0.02776 (7)	0.0477 (5)
C5	0.0941 (2)	0.3941 (3)	0.16042 (7)	0.0607 (6)
N2	0.10664 (19)	0.1037 (2)	-0.06899 (6)	0.0632 (6)
N1	0.0775 (2)	0.3645 (2)	0.11798 (6)	0.0630 (6)
C11	0.2007 (2)	0.0582 (2)	-0.00217 (7)	0.0527 (6)
H11	0.2567	-0.0197	-0.0131	0.063*
C8	0.1356 (2)	0.2458 (3)	0.10037 (7)	0.0541 (6)
H22E	0.1874	0.1778	0.1163	0.065*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C6	0.0832 (19)	0.0809 (18)	0.0664 (17)	0.0181 (15)	0.0077 (14)	0.0081 (14)
C14	0.0464 (12)	0.0402 (11)	0.0703 (15)	0.0053 (9)	0.0058 (11)	0.0021 (10)
C13	0.0470 (13)	0.0445 (11)	0.0669 (14)	0.0029 (9)	-0.0055 (11)	0.0087 (10)
C15	0.0811 (19)	0.0882 (19)	0.0721 (16)	0.0062 (15)	0.0070 (15)	-0.0134 (14)
C16	0.099 (2)	0.0781 (18)	0.0657 (16)	0.0016 (15)	-0.0108 (15)	0.0086 (13)
C10	0.0463 (13)	0.0431 (11)	0.0696 (14)	0.0046 (9)	-0.0046 (11)	0.0095 (10)
C1	0.147 (3)	0.132 (3)	0.0655 (18)	-0.002 (3)	0.001 (2)	-0.0060 (18)
C3	0.094 (2)	0.091 (2)	0.0741 (18)	0.0121 (17)	-0.0165 (16)	-0.0009 (15)
C7	0.092 (2)	0.085 (2)	0.0722 (18)	0.0140 (16)	0.0183 (17)	0.0006 (15)
C4	0.077 (2)	0.0833 (19)	0.0759 (17)	0.0150 (15)	-0.0038 (15)	-0.0099 (14)
C2	0.103 (2)	0.0753 (18)	0.0593 (16)	-0.0054 (16)	0.0072 (16)	0.0052 (13)
C9	0.0459 (12)	0.0381 (11)	0.0607 (13)	-0.0047 (8)	0.0043 (10)	0.0078 (9)
C12	0.0422 (12)	0.0411 (11)	0.0597 (13)	-0.0066 (8)	0.0038 (10)	0.0035 (9)
C5	0.0697 (17)	0.0530 (13)	0.0592 (14)	0.0018 (11)	0.0053 (12)	0.0076 (11)
N2	0.0613 (13)	0.0650 (12)	0.0632 (12)	0.0064 (10)	-0.0013 (10)	0.0009 (10)
N1	0.0684 (14)	0.0608 (12)	0.0599 (12)	0.0050 (10)	0.0046 (10)	0.0042 (9)
C11	0.0443 (13)	0.0429 (11)	0.0710 (14)	0.0051 (9)	0.0024 (11)	0.0004 (10)
C8	0.0524 (13)	0.0428 (12)	0.0670 (14)	-0.0018 (9)	0.0016 (11)	0.0110 (10)

Geometric parameters (\AA , $^\circ$)

C6—C7	1.371 (4)	C1—C2	1.507 (4)
C6—C5	1.378 (3)	C1—H1A	0.9600
C6—H6	0.9300	C1—H1B	0.9600
C14—C13	1.364 (3)	C1—H1C	0.9600
C14—C9	1.398 (3)	C3—C2	1.379 (4)
C14—H14	0.9300	C3—C4	1.381 (4)
C13—C12	1.403 (3)	C3—H3	0.9300
C13—H13	0.9300	C7—C2	1.373 (4)
C15—N2	1.439 (3)	C7—H7	0.9300
C15—H15A	0.9600	C4—C5	1.390 (3)
C15—H15B	0.9600	C4—H4	0.9300
C15—H15C	0.9600	C9—C8	1.441 (3)
C16—N2	1.443 (3)	C12—N2	1.372 (3)

C16—H16A	0.9600	C12—C11	1.397 (3)
C16—H16B	0.9600	C5—N1	1.413 (3)
C16—H16C	0.9600	N1—C8	1.270 (3)
C10—C11	1.369 (3)	C11—H11	0.9300
C10—C9	1.389 (3)	C8—H22E	0.9300
C10—H10	0.9300		
C7—C6—C5	121.6 (3)	C2—C3—H3	119.0
C7—C6—H6	119.2	C4—C3—H3	119.0
C5—C6—H6	119.2	C6—C7—C2	121.8 (3)
C13—C14—C9	121.5 (2)	C6—C7—H7	119.1
C13—C14—H14	119.3	C2—C7—H7	119.1
C9—C14—H14	119.3	C3—C4—C5	120.5 (3)
C14—C13—C12	121.6 (2)	C3—C4—H4	119.8
C14—C13—H13	119.2	C5—C4—H4	119.8
C12—C13—H13	119.2	C7—C2—C3	116.8 (3)
N2—C15—H15A	109.5	C7—C2—C1	121.8 (3)
N2—C15—H15B	109.5	C3—C2—C1	121.4 (3)
H15A—C15—H15B	109.5	C10—C9—C14	116.6 (2)
N2—C15—H15C	109.5	C10—C9—C8	120.6 (2)
H15A—C15—H15C	109.5	C14—C9—C8	122.8 (2)
H15B—C15—H15C	109.5	N2—C12—C11	121.6 (2)
N2—C16—H16A	109.5	N2—C12—C13	121.4 (2)
N2—C16—H16B	109.5	C11—C12—C13	117.1 (2)
H16A—C16—H16B	109.5	C6—C5—C4	117.1 (2)
N2—C16—H16C	109.5	C6—C5—N1	117.5 (2)
H16A—C16—H16C	109.5	C4—C5—N1	125.3 (2)
H16B—C16—H16C	109.5	C12—N2—C15	121.1 (2)
C11—C10—C9	122.6 (2)	C12—N2—C16	121.1 (2)
C11—C10—H10	118.7	C15—N2—C16	117.7 (2)
C9—C10—H10	118.7	C8—N1—C5	120.6 (2)
C2—C1—H1A	109.5	C10—C11—C12	120.6 (2)
C2—C1—H1B	109.5	C10—C11—H11	119.7
H1A—C1—H1B	109.5	C12—C11—H11	119.7
C2—C1—H1C	109.5	N1—C8—C9	123.8 (2)
H1A—C1—H1C	109.5	N1—C8—H22E	118.1
H1B—C1—H1C	109.5	C9—C8—H22E	118.1
C2—C3—C4	122.0 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C11—H11...Cg1 ⁱ	0.93	2.94	3.670 (2)	137

Symmetry code: (i) *x*, $-\gamma+3/2$, *z*-1/2.