

## Ocean Dwelling Actinobacteria as Source of Antitumor Compounds

**Ajitha Gomathi<sup>1</sup>; K M Gothandam<sup>1\*</sup>.**

<sup>1</sup>VIT University, School of Bio Sciences and Technology, India.

### ABSTRACT

*Extending over decades, research has been of great focus and enormous progress on exploring the ocean for natural products from marine actinobacteria. Attraction towards alternative medicine has led to improvements in natural product discovery. With great potential to survive in extreme environments, marine actinobacteria, efficiently produce an array of metabolites with diverse bioactivity by evolving the secondary metabolic pathways. Exploiting the secondary metabolite producing potential of actinobacteria, many compounds with antitumor, antibacterial, antifungal, antimalarial, antiprotozoal, antiparasitic, antiviral, anti-parasitic, anti-inflammatory activities has been discovered. Efforts in bioprospecting alternative sources of natural products have thus led to several explorations and improvements in technologies which has decreased the bottle neck difficulties in the drug discovery process. Emphasizing on the recent advancements in bioactive compound production in actinobacteria, this paper comprises a review of the available literature, compiles the antitumor compounds from marine actinobacteria with brief discussions and the perspectives to develop better antitumor compounds which would stimulate further research.*

**Key words:** Marine actinobacteria, bioactivity, antitumor compounds

---

\*Authors for correspondence: gothandam@gmail.com

## INTRODUCTION

Outstanding biotechnological advances has improved our understanding and led to the achievement of great progress in developing drugs against cancer. Though present in the era of antibiotics, great challenges faced due to development of resistance in microbes and to anticancer drugs by malignant tumor cells has increased medical concern. These alarming needs prompt immediate action towards discovery and development of novel, potent and less toxic drugs. 60% of anticancer agents are derived from natural sources such as plants, marine organisms and micro-organisms. Bioactivities exhibited by the natural products have inspired the researchers to further probe the microbial sources for drug search. Microbes have a long history of use in the treatment of cancer. Microbial metabolites are among the most important of the cancer chemotherapeutic agents. From the 22,500 biologically active compounds that have been obtained so far from microbes, 45% are produced by actinomycetes, 38% by fungi and 17% by unicellular bacteria (Berdy 2005). Actinobacteria being prolific producers of secondary metabolites are novel candidates for synthesis of novel therapeutic drugs, efficient to combat a range of resistant microbes. Past history of extensive actinobacterial research, has proved limited realization of the actinomycete biosynthetic potential (Watve et al. 2001). Thus with considerable scope left over, potential efforts would efficiently increase the novel metabolite discovery.


Actinobacteria, a group of prokaryotic organisms are aerobic, nonmotile, filamentous, Gram-positive saprophytic bacteria with high guanosine–cytosine (GC) content in their DNA (70–80%) (Goodfellow and Williams 1983). Actinobacteria belongs to the phyla actinobacteria which is one of the largest taxonomic unit and the order actinomycetales (Waksman 1959) which currently consists of 10 suborders, 49 families and over 160 genera. Widespread existence in natural ecosystems and greatest potential to produce bioactive compound has made actinobacteria to be greatly exploited for biotechnological applications. With

overexploitation of terrestrial actinobacteria, the focus on marine sources for bioactive compounds gained significant momentum. Ocean, covering more than 70% of the Earth's surface is home to nearly one million multicellular (plants and animals) and one billion unicellular (distributed under 100 different phyla) organisms. (Burgess 2012). Actinobacteria exist with greatest biodiversity in oceans than in terrestrial ecosystems. Actinobacteria adapting to the extreme conditions of temperature, salinity and pressure, produce metabolites for their survival. The secondary metabolites produced by actinobacteria exhibit promising diverse biological activities and also possess varied chemical structures. Figure 1 represents the different biological activities exhibited by the secondary metabolites produced by marine actinobacteria with few listed examples. These metabolites are produced by biosynthetic gene clusters residing in the large genome of the actinobacteria. These clusters contain genes which are responsible for the production of different class of compounds with varied chemical structures. The different classes of compounds reported so far are polyketides, peptides, isoprenoids, indolocarbazoles, aminoglycosides, polyenes, macrolides and many more. These are synthesized by various genes encoding for enzymes such as polyketide synthases, non-ribosomal peptide synthase, dNDP-glucose 4,6 dehydratase, dTDP-glucose 4,6 dehydratase, polyether epoxidase, O-Methyl transferase, P450 Monooxygenase, AHBA synthase, 3-hydroxyl-3-methylglutaryl coenzymeA reductase, oxytryptophan dimertization enzyme, iadomycin cyclase etc. Figure 2 shows the schematic representation of some of the biosynthetic genes present in marine actinobacteria.

This review aims to compile the antitumor compounds from marine actinobacteria, mainly isolated from sea sediments. Table 1-6 comprises of the list of antitumor compounds from marine actinobacteria. The table also represents that a potential number of compounds has been obtained from actinobacteria belonging to *Streptomyceteceae* family.

Biological activity of compounds produced by marine actinobacteria	
Antibacterial activity	<ul style="list-style-type: none"> <li>• Abyssomicins (Riedlinger et al. 2004)</li> <li>• Tirandamycins (Carlson et al. 2009)</li> </ul>
Antifungal activity	<ul style="list-style-type: none"> <li>• Chandrananimycin (Maskey et al. 2003)</li> <li>• N-(2-hydroxyphenyl)-2-phenazinamine (Gao et al. 2012)</li> </ul>
Antitumor activity	<ul style="list-style-type: none"> <li>• Streptokordin (Jeong et al. 2006)</li> <li>• Elaiomycins B and C (Helaly et al. 2011)</li> </ul>
Anti-inflammatory activity	<ul style="list-style-type: none"> <li>• Cyclomarins (Schultz et al. 2008)</li> <li>• Salinamides A and B (Moore et al. 1999)</li> </ul>
Anti-malarial activity	<ul style="list-style-type: none"> <li>• Trioxacarcin (Maskey et al. 2004)</li> </ul>
Anti-viral activity	<ul style="list-style-type: none"> <li>• Benzastatin C (Lee et al. 2007)</li> </ul>
Anti-parasitic activity	<ul style="list-style-type: none"> <li>• Avermectins (Burg et al. 1979)</li> </ul>

**Figure 1**-- Different biological activity of secondary metabolites produced by marine actinobacteria.



**Figure 2** -- Secondary metabolite biosynthetic genes present in marine actinobacteria.

## ANTITUMOR COMPOUNDS FROM MARINE ACTINOBACTERIA

### Polyketides

Table 1 represents the list of compounds from marine actinobacteria with polyketides as chemical structure.

Polyketides are a significant class of compounds exhibiting wide range of pharmacological activities. Three types of polyketide synthases namely type I, II, III, catalyse the condensation reactions of acyl-CoA precursors to produce different polyketides.

**Table 1-** Compounds with polyketide as chemical structure

<b>Compound</b>	<b>Source</b>	<b>Organism</b>	<b>Family</b>	<b>Cancer</b>	<b>Reference</b>
SS-228 Y	Sagami Bay	<i>Streptomyces</i> sp.	<i>Streptomycetaceae</i>	Ehrlich breast adenocarcinoma	Okazaki et al. 1975
Fridamycins(A,B,D) Himalomycin A, B Rabelomycin Aureoverticillactam	Mauritius (Indian ocean) Marine sediment	<i>Streptomyces</i> sp isolate B6921 <i>Streptomyces Aureoverticillatus</i> (NPS001583)	<i>Streptomycetaceae</i> <i>Streptomycetaceae</i>	Colorectal adenocarcinoma, Melanoma, Leukemia	Maskey et al. 2003 Mitchell et al. 2004
(R)-10-methyl-6-undecanolide , (6R, 10S)-10-methyl-6-dodecanolide) Chartreusin	Papua New Guinea Jiaozhou Bay, China	<i>Streptomyces</i> sp. B6007 <i>Streptomyces</i> sp. Strain M518	<i>Streptomycetaceae</i> <i>Streptomycetaceae</i>	Gastric adenocarcinoma, Hepatocarcinoma, Breast cancer Melanoma Leukemia	Stritzke et al. 2004 Wu et al. 2006; McGovren et al. 1977
Actinofuranones A, B	Sediment sample from Guam	<i>Streptomyces</i> Strain CNQ766	<i>Streptomycetaceae</i>		Cho et al. 2006
Nonactin	Sediment sample Ayu Trough	<i>Streptomyces</i> sp. KORDI-3238	<i>Streptomycetaceae</i>	Human erythro leukemia	Jeong et al. 2006; Borrel et al. 2005
Manumycin Chinikomycin A,B	Jiaozhou Bay, China	<i>Streptomyces</i> sp. isolate M045	<i>Streptomycetaceae</i>	Renal cancer Mammary cancer Melanoma	Li et al. 2005
Daryamides A,B	San Diego, California	<i>Streptomyces</i> strain CNQ-085	<i>Streptomycetaceae</i>	Human colon carcinoma	Asolkar et al. 2006
Marinomycins A,B,C,D Saliniketals A and B	La Jolla, California Sediment sample from Guam	<i>Marinispora</i> sp. strain CNQ-140 <i>Salinispora arenicola</i> strain CNR-005	<i>Streptomycetaceae</i> <i>Micromonosporaceae</i>	Human melanoma	Kwon et al. 2006 Williams et al. 2007a; Jensen et al. 2007
Arenicolides A	Sediment sample from Guam	<i>Salinispora arenicola</i> strain CNR-005	<i>Micromonosporaceae</i>	Human colon adenocarcinoma	Williams et al. 2007b
Chalcomycin A	Jiaozhou Bay, China; Pohoiki, Hawaii, Pacific ocean	<i>Streptomyces</i> sp. strain M491; <i>Streptomyces</i> isolate B7064	<i>Streptomycetaceae</i>	Human cervix carcinoma	Wu et al. 2007; Asolkar et al. 2002
Piericidins C7, C8	Iwayama Bay, Palau	<i>Streptomyces</i> sp. YM14-060	<i>Streptomycetaceae</i>		Hayakawa et al. 2007a; Hayakawa et al. 2007b

Resistoflavine, 1-hydroxy-1- norresistomycin	Machilipatnam coast, Bay of Bengal, India	<i>Streptomyces chibaensis</i> (strain AUBN1/7)	<i>Streptomyceteceae</i>	Gastric adenocarcinoma, Hepatic carcinoma	Gorajana et al. 2005; Gorajana et al. 2007
Marmycins A,B	Sea of Cortez, Baja California Sur, México	<i>Streptomyces</i> strain CNH990	<i>Streptomyceteceae</i>	Colon adenocarcinoma , Breast cancer, Prostate cancer, Lung cancer , Leukemia	Martin et al. 2007
Mansouramycin A,B,C,D	North Sea coast, Germany	<i>Streptomyces</i> strain Mei37	<i>Streptomyceteceae</i>	Non-small cell lung cancer, Breast cancer, Melanoma, Prostate cancer	Hawas et al. 2009
Albidopyrone	Atlantic ocean	<i>Streptomyces</i> sp. NTK 227	<i>Streptomyceteceae</i>		Hohmann et al. 2009a
Tartrolon D	East coast of Madagaskar	<i>Streptomyces</i> sp. MDG-04-17-069	<i>Streptomyceteceae</i>	Lung carcinoma, Colorectal carcinoma, Breast adenocarcinoma	Perez et al. 2009
Caerulomycin (F,G,H,I,J,K)	Weihai, China	<i>Actinoalloteichus cyanogriseus</i> WH1-2216-6	<i>Actinomycetaceae</i>	Human promyelocytic leukemia, Human alveolar adenocarcinoma, Human chronic myelogenous leukemia, Human epidermoid carcinoma.	Fu et al. 2011a
Cyanogrisides A,B,C,D	Weihai, China	<i>Actinoalloteichus cyanogriseus</i> WH1-2216-6	<i>Actinomycetaceae</i>	Breast cancer, Human epidermoid carcinoma, Human chronic myelogenous leukemia	Fu et al. 2011b

A species of *Chainia* (now *Streptomyces*) isolated from shallow sea mud in Sagami Bay, produced a compound SS-228 Y which inhibited the growth of Ehrlich carcinoma in mice. SS-228 Y inhibits growth of Ehrlich breast adenocarcinoma in mice. Administering SS-228 Y in mice inoculated with Ehrlich ascites tumor, for 10 days at doses of more than 1.56 µg per day, prolonged the survival period of the mice (Okazaki et al. 1975). Fridamycins (A,B,D), himalomycin A and B, and rabelomycin were reported to be produced by *Streptomyces* sp. isolate B6921 isolated from sandy sediment of a coastal site of Mauritius (Indian Ocean). These

antibiotics possess antibacterial and antitumor activities (Maskey et al. 2003). Aureoverticillactam, a macrocyclic lactam was found to be produced by *Streptomyces aureoverticillatus* (NPS001583) isolated from a marine sediment. The 22 atom macrocyclic lactam has exhibited moderate cytotoxic activity against colorectal adenocarcinoma, melanoma, and leukemia cell lines (Mitchell et al. 2004). A new class of compounds (R)-10-methyl-6-undecanolide and (6R, 10S)-10-methyl-6-dodecanolide belonging to caprolactones are extracted by ethyl acetate from the marine actinomycete *Streptomyces*

sp. B6007. This species was isolated from the mangrove sediment, Papua New Guinea. The compounds (6R, 10S)-10-methyl-6-dodecanolide have shown action against gastric adenocarcinoma (HM02), hepatocarcinoma cell lines (HepG2), and breast cancer cell lines (MCF-7) (Stritzke et al. 2004).

Wu et al. (2006) reported the production of chartreusin from *Streptomyces* sp. M518, derived from a sediment of Jiaozhou Bay in China. Chartreusin causes DNA damage by binding to the DNA and inhibiting RNA and DNA synthesis. Chartreusin was found to be potent inhibitors of the catalytic activity of topoisomerase II (Lorico and Long 1993). Chartreusin exhibited vital chemotherapeutic activity against tumor cell lines such as murine (P388), leukemia (L1210) and melanoma (B16) (McGovren et al. 1977). Cho et al. (2006) reported two new secondary metabolites, namely, actinofuranones A and B produced by marine-derived *Streptomyces* strain designated as CNQ766, which was isolated from sediment sample collected from Guam. *In vitro* cytotoxicity tests revealed weak activity of the polyketides against mouse splenocyte T-cells and macrophages with  $IC_{50}$  values of 20  $\mu$ g/mL.

*Streptomyces* sp. KORDI-3238 isolated from deep-sea sediment sample collected at Ayu Trough, in the western Pacific ocean, produced an ionophore antibiotic macrolide, nonactin (Jeong et al. 2006). Borrel et al. (2005) reported nonactin to possess antitumor activity and to be an effective inhibitor against multidrug-resistant human erythroleukemia cell line K562. Manumycins belong to a class of cytotoxic compounds with side chains consisting of polyketide-derived moieties. The compounds exhibited various other biological activities also. Manumycin A and two novel antitumor antibiotics, chinikomycins A and B were isolated from a marine *Streptomyces* sp. M045, derived from sediment of Jiaozhou Bay in China. These chlorine-containing manumycin derivatives exhibited antitumor activity against different human cancer cell lines namely renal cancer, mammary cancer, melanoma cell line, among which chinikomycin A was potent than chinikomycin B. Chinikomycin A selectively inhibited proliferation in cell lines of mammary cancer MAXF 401NL ( $IC_{50}$  value of 2.41  $\mu$ g/mL), melanoma MEXF 462NL ( $IC_{50}$  value of 4.15  $\mu$ g/mL), and renal cancer RXF 944L ( $IC_{50}$  value of 4.02  $\mu$ g/mL). Chinikomycin B showed selective antitumor activity against MAXF 401NL ( $IC_{50}$  value of 3.04  $\mu$ g/mL) (Li et al. 2005). *Streptomyces*

strain CNQ-085 was isolated from marine sediment collected at a depth of 50 m off San Diego, California. The culture extracts of strain CNQ-085 yielded four new moderately cytotoxic compounds, designated as daryamides A, B, C and (2E,4E)-7-methylocta-2,4-dienoic acid amide. The side chains in the compounds appear to be typical polyketide-derived moieties. Daryamide A exhibited potent cancer cell cytotoxicity ( $IC_{50}$  value of 3.15  $\mu$ g/mL) in the evaluation of cytotoxicity against human colon carcinoma cell line, HCT-116 (Asolkar et al. 2006). *Marinispora* sp. CNQ-140 was isolated from a sediment sample collected at a depth of 56 m offshore of La Jolla, California. The culture extract of the strain led to the isolation of marinomycins A-D which are macrodiolides composed of dimeric 2-hydroxy-6-alkenyl-benzoic acid lactones with conjugated tetraene-pentahydroxy polyketide chains. The compounds inhibited cancer cell proliferation when tested against the NCI's 60 cancer cell line panel with average  $LC_{50}$  values of 0.2-2.7  $\mu$ M. Significant tissue type selectivity was exhibited by marinomycin A against human melanoma cell lines LOX IMVI, M14, SK-MEL-2, SK-MEL-5, UACC-257, and UACC-62, while SK-MEL-5 was the most sensitive strain with an  $LC_{50}$  of 5.0 nM. Marinomycins B and C also showed potent activities with average  $LC_{50}$  values of 0.9 and 0.2  $\mu$ M, respectively (Kwon et al. 2006).

Marine actinomycete *Salinispora arenicola* strain CNR-005, was isolated from a sediment sample collected in Guam at a depth of approximately 30 m. The strain has led to the isolation of two unusual bicyclic polyketides, saliniketals A and B containing a new 1,4-dimethyl-2,8-dioxabicyclo[3.2.1]octan-3-yl ring. Saliniketals A and B were found to inhibit ornithine decarboxylase induction, an important target for the chemoprevention of cancer, with  $IC_{50}$  values of 1.95 and 7.83  $\mu$ g/mL, respectively (Williams et al. 2007a; Jensen et al. 2007). *Salinispora arenicola* CNR-005 also produced three new macrolide polyketides designated arenicolides A-C which are 26-membered polyunsaturated macrolactones containing repeating vicinal hydroxyl methoxyl moieties. Arenicolide A exhibited moderate cytotoxicity toward the HCT-116 cell lines with  $IC_{50}$  value of 30  $\mu$ g/mL (Williams et al. 2007b). *Streptomyces* sp. M491, derived from sediment of Jiaozhou Bay in China, produced macrolide antibiotic chalcomycin A, along with many other compounds (Wu et al. 2007). Another

*Streptomyces* isolate B7064 derived from mangrove sediment near Pohoiki, Hawaii (Pacific ocean), also produced chalcomycin B. Chalcomycin has been found to inhibit protein synthesis in HeLa human cervix carcinoma cell line (Asolkar et al. 2002; Wu et al. 2007). *Streptomyces* sp. YM14-060 was isolated from unidentified greenish ascidians collected at Iwayama Bay, Palau. The isolate produced piericidins C7 and C8, two new members of the piericidin family (Hayakawa et al. 2007a). Rat glial cells transformed with the adenovirus E1A gene (RG-E1A-7), neuro-2a mouse neuroblastoma cells, C6 rat glioma cells and 3Y1 rat normal fibroblasts were used for examining the biological activities of piericidins. Piericidins C7 and C8 exhibited selective cytotoxicity against RG-E1A-7 cells (IC<sub>50</sub> of 1.5 nM and 0.45 nM, respectively), and inhibited the growth of Neuro-2a cells (IC<sub>50</sub> of 0.83 nM and 0.21 nM, respectively) without cytotoxic cell death, while piericidins were not cytotoxic nor cytostatic on C6 rat glioma cells and 3Y1 rat normal fibroblast (Hayakawa et al. 2007b).

*Streptomyces chibaensis* (strain AUBN1/7) was isolated from a marine sediment sample collected at a depth of 30 m at a distance of 8 km off Machilipatnam coast of Bay of Bengal, India. The culture extract of the strain yielded 1-hydroxy-1-norresistomycin and resistoflavine. Both the metabolites showed a potent *in vitro* cytotoxic activity against cell lines namely, HMO2 and HePG2 (Gorajana et al. 2005; Gorajana et al. 2007). *Streptomyces*-related strain CNH990 was obtained from sediment samples collected at the entrance of Sea of Cortez, Baja California Sur, México. The strain yielded two new cytotoxic quinones of the angucycline class, marmycins A and B. Initially the cytotoxic behavior of the compounds were evaluated against the HCT-116 cell lines, in which marmycin A and B showed an IC<sub>50</sub> of 60.5 nM and 1.09 µM respectively. Thus the marmycin A proved to be 18 times more potent in exhibiting cytotoxicity, than marmycin B. Further evaluation of *in vitro* cytotoxicity against 12 human tumor cell lines (breast, prostate, colon, lung, leukemia), revealed marmycin A to be potent, showing a mean IC<sub>50</sub> value of 0.022 µM while marmycin B was significantly less potent with a mean IC<sub>50</sub> value of 3.5 µM (Martin et al. 2007). Mansouramycins are a new class of compounds belonging to isoquinoline quinones. Mansouramycin A was isolated from *Streptomyces* sp. Mei37 from North Sea coast, Germany. Various mansouramycins extracted from

this species are mansouramycin A (7-methylamino-3,4-dimethylisoquinoline-5,8-dione), mansouramycin B (6-chloro-3-methylisoquinoline-5,8-dione), mansouramycin C (3-carbomethoxy-7-methylaminoisoquinoline-5,8-dione), mansouramycin D (3-(1H-indole-3yl)-7-methylaminoisoquinoline-5,8-dione). Different mansouramycins are tested against the 36 cancer cell lines. Mansouramycin C dark red in color, showed highest cytotoxic action compared with other mansouramycins. When tested on 36 different cancer cell lines it exhibited action against 10 cell lines (T-24, SF-268, LXFA629L, MEXF 520L, OVCAR-3, RXF944L and UXF1138L), with inhibition concentrations ranging from 0.008 to 0.02 µM (Hawas et al. 2009). *Streptomyces* sp. NTK 227, a strain isolated from Atlantic ocean sediment, was reported to produce secondary metabolite namely albidopyrone. The pyrone-containing secondary metabolite exhibited cytotoxic activity by moderately inhibiting protein tyrosine phosphatase B (Hohmann et al. 2009a).

Antitumor compound tartrolon D, which belong to a group of macrodiolides, was isolated from the marine actinomycete *Streptomyces* sp. MDG-04-17-069 from the east coast of Madagascar at a depth of 30m. These dimers, with two polyketide chains, had a whitish color and exhibited potent effect on cancer cell lines A549 (lung carcinoma), HT-29 (colorectal carcinoma), and MDA-MB-231 (breast adenocarcinoma) (Pérez et al. 2009). Five bipyridine alkaloids named as caerulomycin F, caerulomycin G, caerulomycin H, caerulomycin I, caerulomycin J, and one phenylpyridine alkaloids named caerulomycin K were isolated from actinomycetes species *Actinoalloteichus cyanogriseus* WH1-2216-6 which showed cytotoxic effect on cancerous cell lines HL-60 (human promyelocytic leukemia), K562, KB (human epidermoid carcinoma of oral cavity) and A549 cell lines. The strain WH1-2216-6 was isolated from marine sediments collected from the seashore of Weihai, China. Analogous of caerulomycins showed antibiotic action against *Escherichia coli*, *Aerobacter aerogenes*, and *Candida albicans* at minimum inhibitory concentrations ranging from 9.7 to 38.6 µM (Fu et al. 2011a). Four novel bipyridine cyclic glycosides namely cyanogrisides A, B, C and D, were isolated from the actinomycete *Actinoalloteichus cyanogriseus* WH1-2216-6 derived from sediment collected from seashore of Weihai, China. Cyanogrisides A exhibited moderate cytotoxicity

against K562, KB, MCF-7 cell lines with IC<sub>50</sub> values of 1.2, 4.7, 9.8 µM while cyanogrisides C exhibited cytotoxicity against K562 and KB cells with IC<sub>50</sub> values of 0.73 and 4.7 µM (Fu et al. 2011b).

## Peptides

Table 2 represents the list of compounds from marine actinobacteria with peptides as chemical structure.

**Table 2-** Compounds with peptides as chemical structure

Compound	Source	Organism	Family	Cancer	Reference
Mechercharmycins	Mecherchar in Palau (North Pacific ocean).	<i>Thermoactinomyces</i> sp. YM3-251	<i>Thermoactinomycetaceae</i>	Human leukemia, Human lung cancer	Kanoh et al. 2005
Lucentamycins	Little San Salvador, Bahamas	<i>Nocardiopsis lucentensis</i> strain CNR-712	<i>Nocardiopsaceae</i>	Human colon carcinoma	Cho et al. 2007
Piperazimycins	Island of Guam	<i>Streptomyces</i> sp. (CNQ-593)	<i>Streptomyceteceae</i>	Human colon carcinoma	Miller et al. 2007
Proximicins	Sea of Japan and Raune Fjord (Norway)	<i>Verrucosipora</i> sp. AB-18-032 ; <i>Verrucosipora</i> strain MG-37	<i>Micromonosporaceae</i>	Gastric adenocarcinoma, Hepatocellular carcinoma, Breast adenocarcinoma	Riedlinger et al. 2004; Fiedler et al. 2008; Schneider et al. 2008
Cyclomarins	Mission Bay, California and Palau	Streptomycete strain CNB-982, <i>Salinispora arenicola</i> CNS-205	<i>Streptomyceteceae</i> , <i>Micromonosporaceae</i>	Human colon carcinoma	Renner et al.1999; Schultz et al. 2008
Arenamides	Great Astrolab Reef in Kandavu Island chain, Fiji	<i>Salinispora arenicola</i> CNT-088	<i>Micromonosporaceae</i>	Human colon carcinoma	Asolkar et al. 2009

Peptides, the polymers of amino acids, are synthesized by non-ribosomal peptide synthetases (NRPS). NRPSs catalyse the condensation of aminoacyl-AMP to produce different peptides which are modified further by activities such as epimerization, methyltransferase, reductase or oxidase.

*Thermoactinomyces* sp. YM3-251 was isolated from sea mud collected at Mecherchar in the Republic of Palau (North Pacific ocean). The strain produced a new cytotoxic substance named mechercharmycin A which are cyclic peptide-like compound, bearing four oxazoles and a thiazole, and mechercharmycin B which are linear peptides. Mechercharmycin A exhibited strong inhibitory activity against Jurkat cells (human leukemia) and A549 cells. Mechercharmycin B did not exhibit antitumor activity against the cell lines. These

results indicate the essentiality of the cyclic structure for the antitumor activity exhibited by mechercharmycin A (Kanoh et al. 2005). Four new structurally unique peptides namely Lucentamycins (A-D), containing 3-methyl-4-ethylideneproline, were produced by *Nocardiopsis lucentensis* CNR-712. The strain was isolated from the sediments of a shallow saline pond from the island of Little San Salvador, in the Bahamas. The only structural differences between these compounds, accounted to the presence of phenyl and indole rings in Lucentamycin A and B, and absence of the moieties in Lucentamycin C and D. *In vitro* cytotoxicity analysis of Lucentamycins A and B showed potent cytotoxicity against HCT-116 cell line with IC<sub>50</sub> values of 0.20 and 11 µM, respectively, whereas lucentamycins C and D were not cytotoxic against HCT-116 cell line. This indicates that the aromatic



ring present is essential for the biological activity of this class of compounds (Cho et al. 2007).

*Streptomyces* sp. CNQ-593 was isolated from a sediment sample collected at a depth of 20 m near the island of Guam. CNQ-593, produced three potent cancer cell cytotoxins, piperazimycins A-C which are cyclic hexadepsipeptides. Initially piperazimycins exhibited significant cytotoxicity with an average GI<sub>50</sub> of 76ng/mL against HCT-116 cell line. In further evaluation, Piperazimycin A also showed potent biological activity against the oncologically diverse 60 cancer cell line panel at the National Cancer Institute, with mean GI<sub>50</sub> value of 100 nM, TGI value of 300 nM and LC<sub>50</sub> value of 2 μM. Overall, piperazimycin A exhibited a nearly 3-fold more potent activity against solid tumors (average LC<sub>50</sub> value of 13.9 μM) than against the leukemia cell lines tested (average LC<sub>50</sub> value of 31.4 μM). In solid tumors, piperazimycin A was most active against the melanoma, central nervous system, prostate cell lines. Piperazimycin A was also active against the colon cancer group, renal cancers, ovarian cancers, non-small cell lung cancers and breast cancers (Miller et al. 2007). A family of structure-related peptide metabolites, named proximicin A, B, and C were reported to be produced by marine actinomycete *Verrucosipora* sp. AB-18-032, *Verrucosipora* sp. MG-37, isolated from sediment collected from the Sea of Japan at a depth of 289 m, and Raune Fjord (Norway) at a depth of 250 m, respectively (Riedlinger et al. 2004; Fiedler et al. 2008). Schneider et al. (2008) reported the structure of proximicins as a unknown γ-amino acid 4-aminofuran-2-carboxylic acid. Proximicins A,B and C exhibited significantly higher growth inhibitory activity towards gastric adenocarcinoma AGS with GI<sub>50</sub> value of 0.6, 1.5, 0.25 μM, respectively and HepG2 cell lines with GI<sub>50</sub> value of 0.82, 9.5, 0.78 μM, respectively, whereas MCF7 cell lines were found to be less sensitive to

proximicins A,B and C. Moreover studies on the effect of proximicins on the cell cycle, revealed an arrest of AGS cells in G0/G1 and an increase in the levels of p53 and p21.

New cyclic heptapeptides, cyclomarins A-C (1-3), were isolated from extracts of a cultured marine *Streptomyces* sp. CNB-982, isolated from a sediment sample collected in Mission Bay, California and cyclomarin D from *Salinispora arenicola* CNS-205, isolated from Palau. (Renner et al. 1999; Schultz et al. 2008). The major metabolite, cyclomarin A was found to be cytotoxic *in vitro* towards cancer cells with mean IC<sub>50</sub> value of 2.6 μM against a panel of human cancer cell lines (Renner et al. 1999) and cyclomarin D displayed moderate cytotoxicity against HCT-116 cell lines with an IC<sub>50</sub> value of 2 μg/mL (Schultz et al. 2008). The extract of a marine actinomycete strain *Salinispora arenicola* CNT-088 yielded three new cyclohexadepsipeptides, arenamides A-C. The strain was obtained from a marine sediment sample collected at a depth of 20 m off the Great Astrolab Reef, in the Kandavu Island chain, Fiji. Arenamides A and B exhibited weak *in vitro* cytotoxicity against HCT-116 cell lines with IC<sub>50</sub> values of 13.2 and 19.2 μg/mL, respectively. Arenamides were reported to be inhibitors of NF-κB, the nuclear transcription factor, thus suppressing carcinogenesis. Stably transfected 293/NF-κB-Luc human embryonic kidney cells, induced by treatment with tumor necrosis factor (TNF) were used to study the effect of arenamides on NF-κB activity. Arenamides A and B blocked TNF-induced activation in a dose and time dependent manner with IC<sub>50</sub> values of 3.7 and 1.7 μM, respectively (Asolkar et al. 2009).

### Polyketide peptides

Table 3 represents the list of compounds from marine actinobacteria with polyketide peptides as chemical structure.

**Table 3-** Compounds with polyketide peptides as chemical structure

Compound	Source	Organism	Family	Cancer	Reference
Salinosporamides	Chub Cay, Bahamas	<i>Salinispora</i> strain CNB-392	<i>Micromonosporaceae</i>	Human colon carcinoma	Feling et al. 2003
Lajollamycin	Scripps Canyon, La Jolla, California	<i>Streptomyces nodosus</i> NPS007994	<i>Streptomyceteceae</i>	Melanoma	Manam et al. 2005

Type I PKS and NRPSs combine to catalyse and allows the formation of the hybrid polyketide-peptides.

*Salinispora* CNB-392 was isolated from a sediment sample collected at a depth of about 1 m from a mangrove environment in Chub Cay,

Bahamas. The strain CNB-392 produced the chemically unique and highly bioactive metabolite salinosporamide A, which inhibited 20S proteasome. The compound was highly cytotoxic *in vitro* toward HCT-116 cell lines (IC<sub>50</sub> ca. 80 ngmL<sup>-1</sup>) and the metabolite also displayed selective activity against NCI 60 cell line panel in which greatest potency was observed against NCI-H226 non-small cell lung cancer, SF-539 CNS cancer, SK-MEL-28 melanoma, and MDA-MB-435 breast cancer (all with LC<sub>50</sub> values less than 10 nm). Salinosporamide A inhibited proteasomal chymotrypsin like proteolytic activity, when tested against purified 20S proteasome (Calbiochem,

cat.no. 539158) with an IC<sub>50</sub> value of 1.3 nm (Feling et al. 2003). A nitro-tetraene spiro- $\beta$ -lactone- $\gamma$ -lactam antibiotic, lajollamycin, was produced by *Streptomyces nodosus* NPS007994 isolated from a marine sediment collected in Scripps Canyon, La Jolla, California. The polyketide-peptide was reported to inhibit the growth of murine melanoma cell line B16-F10, with an EC<sub>50</sub> value of 9.6  $\mu$ M (Manam et al. 2005).

### Isoprenoids

Table 4 represents the list of compounds from marine actinobacteria with isoprenoids as chemical structure.

**Table 4-** Compounds with isoprenoids as chemical structure

Compound	Source	Organism	Family	Cancer	Reference
Altomicidin	Gamo, Japan	<i>Streptomyces sioyaensis</i> SA-1758	<i>Streptomycetaceae</i>	Lymphoid leukemia, Carcinoma	Takahashi et al. 1989a; Takahashi et al. 1989b
Neomarinones	Lagoon, North of San Diego, California	Actinomycete isolate CNH-099	<i>Actinomycetaceae</i>	Human colon carcinoma	Hardt et al. 2000; Kalaitzis et al. 2003
Glaciapyrroles	Alaska	<i>Streptomyces</i> sp. (NPS008187)	<i>Streptomycetaceae</i>	Colorectal adenocarcinoma, Melanoma	Macherla et al. 2005
Drimentine G; Indotertine B	Guangdong province	<i>Streptomyces</i> sp. CHQ-64	<i>Streptomycetaceae</i>	Human colon cancer, Human hepatocellular carcinoma, Human lung cancer, Human ovarian carcinoma;	Che et al. 2012; Che et al. 2013

Isoprenoids forms the largest family, comprising a large number of industrially useful compounds. Isoprenoids are synthesized from the five-carbon precursor molecule isopentenyl diphosphates (IPP) via 2-C-methyl-Derythritol 4-phosphate (MEP) pathway.

*Streptomyces sioyaensis* SA-1758 was isolated from sea mud collected at Gamo, Miyagi Prefecture, Japan. The strain SA-1758 produced a monoterpene-alkaloid, altomicidin (Takahashi et al. 1989a). The compound strongly inhibited the growth of L1210 and carcinoma IMC cell lines with IC<sub>50</sub> values of 0.84 and 0.82  $\mu$ g/mL, respectively

(Takahashi et al. 1989b). Neomarinones are sesquiterpenoid naphtho-quinones with a mixed polyketide-terpenoid origin (Pathirana et al. 1992). A novel marine actinomycete isolate CNH-099 found in a sediment sample collected at 1 m in Batiquitos Lagoon, North of San Diego, California, produced several cytotoxic metabolites related to marinone, such as neomarinone, isomarinone, hydroxydebromomarinone and methoxydebromomarinone. These compounds exhibited moderate *in vitro* cytotoxicity (IC<sub>50</sub> ca. 8  $\mu$ g/mL) against HCT-116 cells. Furthermore, neomarinone generated a mean IC<sub>50</sub> value of 10  $\mu$ M

in the NCI's 60 cancer cell line panel (Hardt et al. 2000; Kalaitzis et al. 2003). Glaciapyrroles (pyrrolosoquiterpenes) were found to be produced by *Streptomyces* sp. NPS008187 that was isolated from a marine sediment collected in Alaska. Glaciapyrroles inhibited both HT-29 and B16-F10 tumor cell growth (Macherla et al. 2005). Actinomycete strain CNQ-525, was isolated from a sediment sample derived from a depth of 152 m near La Jolla, California. The culture broth of the strain yielded three new chlorinated dihydroquinones and a previously reported analogue. These chlorine-containing terpenoid dihydroquinones possessed significant cytotoxicity against HCT-116 cell lines (Soria-Mercado et al. 2005).

*Streptomyces* sp. CHQ-64 was isolated from the reeds rhizosphere soil of mangrove conservation

area in Guangdong province, China. The culture broth yielded indotertine A and B, hybrid isoprenoid with a condensed pentacyclic skeleton, and two related compounds drimentines F, G and H. Drimentines G showed potent cytotoxicity against HCT-8, (human colon cancer cell line), Bel-7402 (human hepatocellular carcinoma cell line), A549 and A2780 (human ovarian carcinoma cell line), with IC<sub>50</sub> values of 2.81, 1.38, 1.01, and 2.54 μM, respectively. Indotertine B exhibited best cytotoxic activities against the HCT-8 and A549 cell lines with IC<sub>50</sub> values of 6.96 and 4.88 μM, respectively (Che et al. 2012; Che et al. 2013).

### Indolocarbazoles

Table 5 represents the list of compounds from marine actinobacteria with indolocarbazoles as chemical structure.

**Table 5-** Compounds with indolocarbazoles as chemical structure

Compound	Source	Organism	Family	Cancer	Reference
ZHD-0501	Jiaozhou Bay, China	<i>Actinomadura</i> sp. 007	<i>Thermomonosporaceae</i>	Human lung adenocarcinoma, Hepatocarcinoma, Pro-myelocytic leukemia	Han et al. 2005
N-formyl staurosporine, N-formyl staurosporine, Selina-4(14),7(11)-diene-8,9-diol	Jiaozhou Bay, Qindao, China	<i>Streptomyces</i> sp isolate QD518	<i>Streptomycetaceae</i>	Bladder, CNS, Colon, Gastric, Head and Neck, Lung, Mammary, Ovarian, Pancreatic, Prostate, Renal cancer	Wu et al. 2006
K252c, Arcyriaflavin A	Coast of Qingdao, China	Actinomycete strain Z2039-2	<i>Actinomycetaceae</i>	Human chronic myelogenous leukemia	Liu et al. 2007
Albidopyrone	Atlantic ocean sediment	<i>Streptomyces</i> sp. NTK 227	<i>Streptomycetaceae</i>		Hohmann et al. 2009a
Chromomycin	Bijiatuan, China	<i>Streptomyces</i> sp. WBF16	<i>Streptomycetaceae</i>	Human gastric cancer, Human liver hepatocellular carcinoma, Human colon cancer, Human lung adenocarcinoma	Lu et al. 2012
Streptokordin	Ayu Trough, Pacific ocean	<i>Streptomyces</i> sp. KORDI-3238	<i>Streptomycetaceae</i>	Human breast cancer, Human renal cancer,	Jeong et al. 2006

				Human skin cancer, Human leukemia	
Mansouramycin C	Jade Bay, southern German North Sea coast	<i>Streptomyces</i> sp. Isolate (Mei37)	<i>Streptomycetaceae</i>	Non-small cell lung cancer, Melanoma, Breast cancer, Prostate cancer	Hawas et al. 2009
Usabamycin A-C	Usa Bay, Kochi Prefecture, Japan	<i>Streptomyces</i> sp. NPS853	<i>Streptomycetaceae</i>	Human epithelial carcinoma	Sato et al. 2011
1(10-aminodecyl) Pyridinium	Bay of Bengal	<i>Amycolatopsis alba</i> (DVR D4)	<i>Pseudonocardiaceae</i>	Cervix cancer, Brain cancer	Dasari et al. 2012
Stretocarbazoles A,B	Sanya, Hainan Province, China	<i>Streptomyces</i> sp. strain FMA	<i>Streptomycetaceae</i>	Human promyelocytic leukemia, Human alveolar adenocarcinoma	Fu et al. 2012
1-ethyl- $\beta$ -carboline-3-carboxylic acid, 1-methyl indole-3-carboxamide, 1-vinyl- $\beta$ -carboline-3-carboxylic acid	Khuean Srinagarindra National Park, Kanchanaburi Province, Thailand	<i>Actinomadura</i> sp. BCC 24717	<i>Thermomonosporaceae</i>	Human epidermoid carcinoma of oral cavity, Human small cell lung cancer	Kornsakulkarn et al. 2013
Spiroindimicins A-D, Indimicin B	South China Sea and Indian ocean	<i>Streptomyces</i> sp. SCSIO 03032	<i>Streptomycetaceae</i>	Human lymphoblastic leukemia, Murine melanoma, Human lung cancer, Human liver cancer, Human breast cancer	Zhang et al. 2014

Indolocarbazoles are distinct family of compounds displaying wide range of biological activities such as antitumor, antibacterial and antifungal. These form unique class of compounds with characteristic structure of indolo[2,3-*a*]pyrrolo[3,4-*c*]carbazole as the core which is derived from two molecules of tryptophan.

*Actinomadura* sp. 007, was isolated from sea sediment sample collected in Jiaozhou Bay, China. The culture broth of the strain yielded a staurosporine analog, ZHD-0501. The compound exhibited cytotoxic activity against A549, BEL-

7402, and pro-myelocytic leukemia HL60 cancer cell lines, P388 cells and mouse mammary cancer tsFT210 cells (Han et al. 2005). Wu et al. (2006) reported the isolation of *Streptomyces* sp. QD518 from the Jiaozhou Bay of Qindao, China. Among the metabolites produced by the actinobacteria, N-formylstaurosporine, N-carboxamido-staurosporine, selina-4(14),7(11)-diene-8,9-diol exhibited *in vitro* anticancer activity against 37 human tumor cell lines derived from solid human tumors comprising bladder, central nervous system, colon, gastric, head and neck, lung, mammary,

ovarian, pancreatic, prostate and renal cancers, as well as cell lines established from human melanoma, pleuramesothelioma and the uteri body. Indolocarbazole alkaloids, namely K252c and arcyliaflavin A were isolated from actinomycete strain Z2039-2, collected on the coast of Qingdao. Cytotoxic effects of the two compounds against K562 cell lines have been evaluated. The indolocarbazoles has also exhibited significant apoptic activity (Liu et al. 2007). *Streptomyces* sp. NTK 227, a strain isolated from Atlantic ocean sediment, was reported to produce secondary metabolite namely albidopyrone. The pyrone containing secondary metabolite exhibited cytotoxic activity by moderately inhibiting protein tyrosine phosphatase B (Hohmann et al. 2009a). A marine-derived actinomycete, identified as *Streptomyces* sp. WBF16, was isolated from sea sediments located in Bijiatuan, in the city of Weihai, China. Chromomycin A2, chromomycin A3, and chromomycin B which are a family of aureolic acids are found to be produced by WBF16. Chromomycin (A2, A3, and B) exhibited cytotoxic activity against human gastric cancer cell line (SGC7901), HepG2, A549, HCT116, human ovarian cancer cell line (COC1), while Chromomycin B alone exhibited strong cytotoxic activity against Human Umbilical Vein Endothelial Cells (HUVEC) (Lu et al. 2012). A new cytotoxic compound, streptokordin was reported to be produced by *Streptomyces* sp. KORDI-3238, which was isolated from a deep-sea sediment sample collected at Ayu Trough, in the western Pacific ocean. *In vitro* cytotoxicity against several human cancer cell lines such as MDA-MB-231, HCT 15, human prostate cancer (PC-3), human lung cancer (NCI-H23), human renal cancer (ACHN), human skin cancer (LOXIMVI), K562 proved that streptokordin exhibited significant cytotoxicity (Jeong et al. 2006). Hawas et al. (2009) reported the isolation of *Streptomyces* sp. Mei37 from the muddy sediment of Jade Bay on the southern German North Sea coast, which produced five isoquinolinequinones among which mansouramycin C as the main quinone. *In vitro* cytotoxicity analysis of the derivatives against 36 human tumor cell lines indicated significant cytotoxicity with pronounced selectivity for non small cell lung cancer, breast cancer, melanoma, and prostate cancer cells. New anthramycin type analogues, usabamycin A, B and C are produced by *Streptomyces* sp. NPS853, isolated from a marine sediment sample collected at

a depth of 20 m near Usa Bay, Kochi Prefecture, Japan. The usabamycins (A-C) exhibited weak cytotoxic activity against HeLa cells *in vitro*, with IC<sub>50</sub> values of 106.6 μM, 103.5 μM, 101.9 μM respectively, and selective inhibition of serotonin (5-hydroxytryptamine) 5-HT<sub>2B</sub> uptake (Sato et al. 2011). *Amycolatopsis alba* (DVR D4) was isolated from the sediments from Bay of Bengal (NTPC area, Visakhapatnam, India). The strain produced a cytotoxic compound identified as 1-(10-aminodecyl) pyridinium salt antibiotic. The compound exhibited significant cytotoxic activity profile against cancer cell lines of HeLa, MCF-7 and brain (U87MG) *in vitro* (Dasari et al. 2012). *Streptomyces* sp. strain FMA was isolated from a mangrove soil collected in Sanya, Hainan Province of China. The ethyl acetate extracts contained two novel indolocarbazoles, streptocarbazoles A and B, which exhibited potent cytotoxicity against HL-60 and A549 cell lines (Fu et al. 2012). *Actinomadura* sp. BCC 24717 was isolated from soil, collected at Khuean Srinagarindra National Park, Kanchanaburi Province, Thailand. The culture broth yielded four new β-carbolines namely methyl 1-(2-methyl carbamate)ethyl- β-carboline-3-carboxylate, methyl 1-(propionic acid)- β-carboline-3-carboxylate, methyl 1-(methyl propionate)- β-carboline-3-carboxylate, 1-ethyl- β-carboline-3-carboxylic acid, and two new indoles namely 1-hydroxymethylindole-3-carboxylic acid, 1-methyl indole-3-carboxamide. The compounds, 1-ethyl-β-carboline-3-carboxylic acid and 1-methyl indole-3-carboxamide exhibited cytotoxicity to Vero African green monkey kidney cell lines (IC<sub>50</sub> 35.91 mg/mL), while the compound, 1-vinyl- β-carboline-3-carboxylic acid was moderately active against KB cells, human small cell lung cancer (NCI-H187), and Vero cells (IC<sub>50</sub> 14.50–23.88 mg/mL) (Kornsakulkarn et al. 2013). Spiroindimicins A-D, were produced by the culture extract of *Streptomyces* sp. SCSIO 03032 which was isolated from South China Sea and Indian ocean. The bisindole alkaloids were evaluated for cytotoxicity against various cell lines. Spiroindimicin B exhibited moderate cytotoxic activities against human lymphoblastic leukemic cell line (CCRF-CEM), B16 and human lung cancer cell line (H460) with IC<sub>50</sub> values of 4, 5, and 12 μg/mL, respectively. Spiroindimicin C showed cytotoxicity against HepG2 and H460 with IC<sub>50</sub> values of 6 and 15 μg/mL. Spiroindimicin D showed moderate inhibitory effects against HepG2, B16, and H460 (Zhang et al. 2012). The strain

SCSIO 03032 was also reported to produce five new bisindole alkaloids, indimicins A-E and lynamicins F and G. Indimicin B exhibited cytotoxicity against MCF-7 cell lines with IC<sub>50</sub> value 10.0 µM (Zhang et al. 2014).

### Other chemical structures

Table 6 represents the list of compounds from marine actinobacteria with other chemical structures.

**Table 6-** Compounds with other chemical structure

Compound	Source	Organism	Family	Cancer	Reference
Benzastatins	Sokcho-city, Kangwon-do, Korea	<i>Streptomyces nitrosporeus</i> 30643	<i>Streptomycetaceae</i>	Neuroblastoma	Kim et al. 1996
Chandrananimycin A,B,C, Questiomycin A, N-acetylquestiomycin, Iodinin and 1,6-phenazinediol	Jiaozhou Bay in China	<i>Actinomadura</i> sp. isolate M048	<i>Thermomonosporaceae</i>	Human colon carcinoma, Melanoma, Human lung carcinoma, Human breast carcinoma, Human kidney tumor, Human uterus cancer	Maskey et al. 2003
Butenolides	TaiPingJiao, QingDao, China	<i>Streptoverticillium luteoverticillatum</i> 11014	<i>Streptomycetaceae</i>	Human leukemia Murine lymphoma	Li et al. 2006
Bohemamine	Island of Guam	<i>Streptomyces</i> sp. CNQ-583	<i>Streptomycetaceae</i>	Human colon adenocarcinoma	Bugni et al. 2006
Streptochlorin	Ayajin Bay, Korea	<i>Streptomyces</i> sp. 04DH110	<i>Streptomycetaceae</i>	Human leukemia	Shin et al. 2007
Streptopyrrolidine	Ayu Trough	<i>Streptomyces</i> sp. KORDI-3973	<i>Streptomycetaceae</i>		Shin et al. 2008b
Caboxamycin	Atlantic ocean, Canary Islands	<i>Streptomyces</i> sp. NTK 937	<i>Streptomycetaceae</i>	Human gastric adenocarcinoma, Hepaticellular carcinoma, Breast carcinoma	Hohmann et al. 2009b
Ammosamides	Bahamas islands	<i>Streptomyces</i> strain CNR-698	<i>Streptomycetaceae</i>	HCT-116 colon carcinoma	Hughes et al. 2009
Dermacozine	Mariana Trench sediment	<i>Dermacoccus abyssii</i> sp. nov., strains MT1.1 and MR1.2,	<i>Dermacoccaceae</i>	Human chronic myelogenous leukemia	Abdel-Mageed et al. 2010
N-(2-hydroxyphenyl)-2-phenazinamine	Arctic ocean	<i>Nocardia dassonvillei</i> BM-17	<i>Nocardiaceae</i>	Human liver hepatocellular carcinoma, Human lung adenocarcinoma, Human colon adenocarcinoma, Human ovarian cancer	Gao et al. 2012

### Benzastatins

*Streptomyces nitrosporeus* 30643 was isolated from the soil collected in Sokcho-city, Kangwon-do,

Korea. The culture broth of the strain 30643 contained benzastatins, rare metabolites, which are free radical scavengers. Benzastatins A and B, have a aminobenzamide skeleton whereas benzastatins C and D have a tetrahydroquinoline skeleton. Cytotoxicity analysis of benzastatins against neuroblastoma and retina hybrid cell line N18-RE-105 showed that benzastatin C exhibited cytotoxicity with an IC<sub>50</sub> value of 38.1  $\mu$ M (Kim et al. 1996).

### Chandrananimycin

*Actinomadura* sp. isolate M048 was derived from the sediment collected from Jiaozhou Bay in China. The crude extract of the strain M048 yielded three novel antibiotics designated as chandrananimycin A, B and C, containing phenoxazinone chromophores and six known metabolites, namely iodinin, 1,6-phenazinediol, questiomycin A, N-acetylquestiomycin, genistein, daidzein. The compounds questiomycin A, N-acetylquestiomycin, Chandrananimycin A, B and C exhibited cytotoxicity with IC<sub>70</sub> values down to 1.4  $\mu$ g/ml, against the human cell lines, CCL HT29 (colon carcinoma), MEXF 514L (melanoma), LXFA 526L (lung carcinoma), LXFL 529L (lung carcinoma), CNCL SF268, LCL H460, MACL MCF-7 (breast carcinoma), PRCL PC3M, RXF 631L (kidney tumor). Iodinin and 1,6-phenazinediol showed antitumour activity against the human cell lines LXFA 629L and LXFL 529L (lung), MAXF 401NL, MEXF 462NL, RXF 944L and UXF 1138L (uterus) with IC<sub>50</sub> values of 3.6 and 3.2  $\mu$ g/ml, respectively (Maskey et al. 2003).

### Butenolides

The bacterial strain *Streptoverticillium luteoverticillatum* 11014 was isolated from underwater sediment collected off the coast of TaiPingJiao, QingDao, China. The fermentation extract of the strain yielded four known butenolides namely (4S)-4,10-dihydroxy-10-methyl-undec-2-en-1,4-olide (compound 1), (4S)-4,10-dihydroxy-10-methyl-dodec-2-en-1,4-olide (compound 2) and two diastereomeric (4S)-4,11-dihydroxy-10-methyl-dodec-2-en-1,4-olides (compound 3 and 4). *In vitro* cytotoxicity analysis by SRB assay revealed that the compounds exhibited cytotoxicity at high concentrations against K562 cell lines (IC<sub>50</sub> values of 8.73, 6.29, and 1.05  $\mu$ mol/mL) and P388 cell lines (IC<sub>50</sub> values of 0.34, 0.19 and 0.18  $\mu$ mol/mL) (Li et al. 2006).

### Bohemamine

Marine sediment sample collected from the island of Guam, was used to isolate the strain CNQ-583, identified as *Streptomyces* species. Pyrrolizidine alkaloids, bohemamine B (1), bohemamine C (2), and 5-chlorobochemamine C (3) were isolated from the extract of the strain. The compounds were tested for inhibition of HCT-116 cell lines but were found to be inactive (Bugni et al. 2006). Zhang et al. (2003) reported that bohemamine and deoxybohemamine to be cell adhesion inhibitors based on LFA-1/ICAM-1 capable of inhibiting adhesion of HL-60 cells to Chinese hamster ovary cells transfected with human ICAM-1, at IC<sub>50</sub> values of 24.3 and 27.2  $\mu$ g/mL.

### Streptochlorin

The bacterial strain designated as *Streptomyces* sp. 04DH110 was isolated from marine sediment collected from Ayajin Bay, on the East Sea of Korea. The strain yielded an antiproliferative compound streptochlorin, with 4-chloro-5-(1H-indol-3-yl) oxazole as a gross structure. In the sulforhodamine B (SRB) assay, streptochlorin exhibited significant *in vitro* growth inhibitory activity against K562 cell lines with IC<sub>50</sub> value of 1.05  $\mu$ g/ml) and immortalized hepatocytes (CHANG) derived from normal human liver (with IC<sub>50</sub> value of 10.9  $\mu$ g/ml) (Shin et al. 2007). Investigating the pro-apoptotic effect of streptochlorin in human leukemic U937 cells, it was found that the pro-apoptotic effect of streptochlorin is mediated through activation of caspases and mitochondria in U937 cells (Park et al. 2008; Shin et al. 2008a). Streptochlorin also exhibited antiangiogenic activity by inhibition of endothelial cell invasion and tube formation stimulated with vascular endothelial cell growth factor, by interrupting the TNF- $\alpha$ -induced NF- $\kappa$ B activation (Choi et al. 2007).

### Streptopyrrolidine

*Streptomyces* sp. KORDI-3973 isolated from a deep-sea sediment sample collected at Ayu Trough, yielded a benzyl pyrrolidine derivative, streptopyrrolidine. The compound exhibited anti-angiogenesis activity and blocked the capillary tube formation without showing cytotoxicity against HUVECs at the concentration of 100  $\mu$ g/ml (Shin et al. 2008b).

### Caboxamycin

A benzoxazole compound named caboxamycin, was reported to be produced from the culture extract of *Streptomyces* sp. NTK 937, which was isolated from an Atlantic ocean deep-sea sediment core collected near the Canary Islands. Cytotoxicity analysis of caboxamycin showed moderate growth inhibitory activity towards AGS, HepG2 and MCF7 cell lines with GI<sub>50</sub> of 7.5, 7.4 and 7.3 µg/mL, respectively (Hohmann et al. 2009b).

#### **Ammosamides**

*Streptomyces* sp. CNR-698, isolated from bottom sediments collected at a depth of 1618 meters in the Bahamas Islands. The fermentation broth of the strain yielded ammosamides A and B, a pyrroloiminoquinone compound with thio-γ-lactam functionality. The compounds exhibited significant *in vitro* cytotoxicity against HCT-116 cell lines, each with IC<sub>50</sub> value of 320 nM. Further the molecular target of the ammosamides was identified as a member of the myosin family, important cellular proteins that are involved in cell cycle regulation, cytokinesis, and cell migration (Hughes et al. 2009).

#### **Dermacozine**

Dermacozines A-G are phenazine alkaloids, isolated from *Dermacoccus abyssi* sp. nov., strains MT1.1 and MT1.2, which was derived from Mariana Trench sediment. The compounds exhibited low cytotoxicity against K562 cell line among which highest cytotoxicity was observed for dermacozines F and G with IC<sub>50</sub> value of 9 and 7 µM, respectively (Abdel-Mageed et al. 2010).

#### **N-(2-hydroxyphenyl)-2-phenazinamine:**

*Nocardia dassonvillei* BM-17 was isolated from a sediment sample collected in the Arctic ocean. The culture broth yielded a novel metabolite N-(2-hydroxyphenyl)-2-phenazinamine (NHP) and six known antibiotics. NHP displayed potent cytotoxicity against HepG2, A549, HCT-116 and COC1 cell lines among which high cytotoxicity was observed against HCT-116 with IC<sub>50</sub> value of 27.82 µg/ml (Gao et al. 2012).

### **REQUIREMENT OF TARGETED RESEARCH**

Screening the culture extracts of actinobacteria for anticancer activity by *in vitro* assays has been the existing method for identification of potential actinobacteria. *In vivo* bioassays also augment the

traditional cell based screening method for anticancer drugs. Random screening may lead to discovering anticancer metabolites from actinobacteria, but focused research through gene based screening methods are still more effective in discovering lead compounds. Though successful, contemporary screening protocols need to be coupled with promising new strategies for efficient cancer treatment. Thus widening the scope of finding novel and efficient drugs to treat cancer involves tremendous knowledge about tumour biology. With enormous progress in significant understanding about cancer progression, utilization of technical capabilities towards targeted research is the fundamental criteria for successful screening of antitumorals from actinobacteria. Achieving effective cancer therapy requires integrative approach of experimentation and underlies exploring molecular targets of cancer. Functional screening of the metabolites not only reveals novel mechanism of action but also prove out to be successful targets of cancer therapy by exploitation of the selective and specific biological activities of the metabolites such as inhibitors of protein-protein interactions and cell cycle targets (Aggarwal et al. 2006). Inhibition of proteasome function has emerged as a powerful strategy for anti-cancer therapy (Crawford et al. 2011). Thus evaluating the ability of the bioactive compounds from marine actinobacteria, to inhibit aromatase and proteasome using various noncellular, cell-based and *in vivo* assays will be an effective screening strategy to develop anticancer metabolite. Optimizing through these specific targets would thus prove out to be efficient method of search for cancer therapeutics.

#### **INSILICO APPROACH**

The successful genome mining is effectively encouraged by focusing on *insilico* recognition of the most promising targets within genome. The correct identification of potential gene clusters for secondary metabolites is thus effectively accomplished using specialized bioinformatics tools and resources. This facilitates the natural product discovery to a greater extent, for the reliable prediction of the genes involved in the secondary metabolism, and gaining insight about the metabolite, further supports in reduction of the timeline involved in the drug discovery process, thus accelerating experimental research. Few of the *insilico* methods which are already in use to



automate the analysis of secondary metabolism in microbes, are listed below in table 7.

**Table 7** – Insilico methods

Tools/ Programs/ Databases	Reference
antiSMASH (antibiotics & Secondary Metabolite Analysis Shell)	Medema et al. 2011; Blin et al. 2013
BAGEL3	de Jong et al. 2010; van Heel et al. 2013
CLUSEAN (Cluster Sequence Analyzer)	Weber et al. 2009
ClustScan	Starcevic et al. 2008
MIDDAS-M	Umemura et al. 2013
NaPDoS (Natural Product Domain Seeker)	Ziemert et al. 2012
NP.searcher (Natural Product <i>searcher</i> )	Li et al. 2009
NRPSPredictor	Röttig et al. 2011
PKMiner	Kim and Yi 2012
SBSPKS (Structure Based Sequence Analysis of Polyketide Synthases)	Anand et al. 2010
SEARCHPKS	Yadav et al. 2003
SMURF (Secondary Metabolite Unknown Regions Finder)	Khaldi et al. 2010

## CONCLUSION

Understanding the dynamics of cancer is the essence to develop adequate therapies. Also, supreme importance is needed in perceiving the molecular mechanisms adapted by tumor cells in the development of resistance against multiple drugs. *Insilico* approach to decipher biosynthetic potential complements the traditional screening methodology and thus augmenting the drug research. With discrete pharmaceutical and biotechnological applications, actinobacteria render them as promising therapeutic agents. The bioactive compounds with distinct chemical structures, from marine actinobacteria are thus promising dominant leads in cancer drug discovery. Harnessing the biological capabilities of marine actinobacteria through systematic and intensive research and exposing new platforms of study could pave way for sustained discovery of potential bioactive compounds for therapeutic use.

## REFERENCES

- Abdel-Mageed WM, Milne BF, Wagner M, Schumacher M, Sandor P, Pathom-aree W. Dermacozines, a new phenazine family from deep-sea dermacocci isolated from a Mariana Trench sediment. *Org Biomol Chem.* 2010; 8 (10): 2352–62.
- Aggarwal BB, Sethi G, Nair A, Ichikawa H. Nuclear factor-B: a Holy Grail in cancer prevention and therapy. *Curr Signal Transduction Therapy.* 2006; 1 (1): 25–52.
- Anand S, Prasad MV, Yadav G, Kumar N, Shehara J, Ansari MZ, Mohanty D. SBSPKS: structure based sequence analysis of polyketide synthases. *Nucleic Acids Res.* 2010; 38: W487–W496.
- Asolkar RN, Maskey RP, Helmke E, Laatsch H. Chalcomycin B, a new macrolide antibiotic from the marine isolate *Streptomyces sp.* B7064. *J Antibiot.* 2002; 55, 893-898.
- Asolkar RN, Jensen PR, Kauffman CA and Fenical W. Daryamides A-C weakly cytotoxic polyketides from a marine derived actinomycete of the genus *Streptomyces* strain CNQ-085. *J Nat Prod.* 2006; 69: 1756–1759.
- Asolkar RN, Freel KC, Jensen PR, Fenical W, Kondratyuk TP, Park E-J. Arenamides A-C, cytotoxic NF- $\kappa$ B inhibitors from the marine Actinomycete *Salinispora arenicola*. *J Nat Prod.* 2009; 72 (3): 396–402.
- Berdy J. Bioactive microbial metabolites. A personal view. *J. Antibiot.* 2005; 58: 1–26.
- Blin K, Medema MH, Kazempour D, Fischbach M, Breitling R, Takano E, Weber T. antiSMASH 2.0 – a versatile platform for genome mining of secondary metabolite producers. *Nucleic Acids Res.* 2013; 41: W204–W212.
- Borrel MN, Pereira E, Fiallo M, Garnier Suillerot A. Mobile ionophores are a novel class of P-glycoprotein inhibitors. *Eur J Biochem.* 2005; 223 (1): 125–33.

- Bugni TS, Woolery M, Kauffman CA, Jensen PR, Fenical W. Bohemamines from a marine-derived *Streptomyces* sp. *J Nat Prod.* 2006; 69: 1626-1628.
- Burg R W, Miller B M, Baker E E, Birnbaum J, Currie S A, Hartman R. Avermectins, new family of potent anthelmintic agents: producing organism and fermentation. *Antimicrob Agents Chemother.* 1979; 15 (3): 361–367.
- Burgess JG. New and emerging analytical techniques for marine biotechnology. *Curr Opin Biotechnol.* 2012; 23: 29–33.
- Carlson J C, Li S, Burr D A, Sherman D H. Isolation and characterization of tirandamycins from a marine-derived *Streptomyces* sp. *J Nat Prod.* 2009; 72: 2076–2079.
- Che Q, Zhu T, Qi X, Mándi A, Kurtán T, Mo X, Li J, Gu Q, Li D. Hybrid Isoprenoids from a Reeds Rhizosphere Soil Derived Actinomycete *Streptomyces* sp. CHQ-64. *Org Lett.* 2012; 14: 3438–3441.
- Che Q, Zhu T, Keyzers RA, Liu X, Li J, Gu Q, Li D. Polycyclic Hybrid Isoprenoids from a Reed Rhizosphere Soil Derived *Streptomyces* sp. CHQ-64. *J Nat Prod.* 2013; 76: 759–763.
- Cho JY, Kwon HC, Williams PG, Kauffman CA, Jensen PR, Fenical W. Actinofuranones A and B, polyketides from a marine-derived bacterium related to the genus *Streptomyces* (Actinomycetales). *J Nat Prod.* 2006; 69 (3): 425–8.
- Cho JY, Williams PG, Kwon HC, Jensen PR, Fenical W. Lucentamycins AD, cytotoxic peptides from the marine-derived actinomycete *Nocardiopsis lucentensis*. *J NatProd.* 2007; 70 (8): 1321–8.
- Choi IK, Shin HJ, Lee HS, Kwon HJ. Streptochlorin, a marine natural product, inhibits NF- $\kappa$ B activation and suppresses angiogenesis *in vitro*. *J Microbiol Biotechnol.* 2007; 17: 1338-1343.
- Crawford LJ, Walker B, Irvine AE. Proteasome inhibitors in cancer therapy. *J Cell Commun Signal.* 2011; 5 (2): 101-110.
- Dasari VRRK, Muthyala MKK, Nikku MY, Donthireddy SRR. Novel Pyridinium compound from marine actinomycete, *Amycolatopsis alba* var. nov., DVR D4 showing antimicrobial and cytotoxic activities *in vitro*. *Microbiol Res.* 2012; 167: 346–51.
- de Jong A, van Heel AJ, Kok J, Kuipers OP. BAGEL2: mining for bacteriocin genomic data. *Nucleic Acids Res.* 2010; 38: W647–W651.
- Feling RH, Buchanan GO, Mincer TJ, Kauffman CA, Jensen PR, Fenical W. Salinosporamide A: a highly cytotoxic proteasome inhibitor from a novel microbial source, a marine bacterium of the new genus *Salinospora*. *Angew Chem Int Ed.* 2003; 42 (3): 355–7.
- Fiedler H-P, Bruntner C, Riedlinger J, Bull AT, Knutsen G, Goodfellow M. Proximicin A, B and C, novel aminofuran antibiotic and anticancer compounds isolated from marine strains of the actinomycete *Verrucosispora*. *J Antibiot.* 2008; 61 (3): 158–63.
- Fu P, Wang SX, Hong K, Li X, Liu PP, Wang Y. Cytotoxic bipyridines from the marine-derived actinomycete *Actinoalloteichus cyanogriseus* WH1-2216-6. *J Nat Prod.* 2011a; 74: 1751–1756.
- Fu P, Liu PP, Li X, Wang Y, Wang SX, Hong K, Zhu WM. Cyclic bipyridine glycosides from the marine-derived actinomycete *Actinoalloteichus cyanogriseus* WH1-2216-6. *Org Lett.* 2011b; 13: 5948–5951.
- Fu P, Yang C, Wang Y, Liu P, Ma Y, Xu L. Streptocarbazoles A and B, two novel indolocarbazoles from the marine derived actinomycete strain *Streptomyces* sp. FMA. *Org Lett.* 2012; 14: 2422–2425.
- Gao X, Lu Y, Xing Y, Ma Y, Lu J, Bao W. A novel anticancer and antifungus phenazine derivative from a marine actinomycete BM-17. *Microbiol Res.* 2012; 167: 616–622.
- Goodfellow M, Williams ST. Ecology of actinomycetes. *Annu. Rev. Microbiol.* 1983; 37: 189–216.
- Gorajana A, Kurada BV, Peela S, Jangam P, Vinjamuri S, Poluri E, Zeeck A. 1-Hydroxy-1-norresistomycin, a new cytotoxic compound from a marine actinomycete, *Streptomyces chibaensis* AUBN1/7. *J Antibiot.* 2005; 58: 526-529.
- Gorajana A, Vinjamuri S, Kurada BV, Peela S, Jangam P, Poluri E. Resistoflavine, cytotoxic compound from a marine actinomycete, *Streptomyces chibaensis* AUBN1/7. *Microbiol Res.* 2007; 162 (4): 322–7.
- Han XX, Cui CB, Gu QQ, Zhu WM, Liu HB, Gu JY, Osada H. ZHD-0501, a novel naturally occurring staurosporine analog from *Actinomadura* sp. 007. *Tetrahedron Lett.* 2005; 46: 6137–6140.
- Hardt IH, Jensen PR, Fenical W. Neomarinone, and new cytotoxic marinone derivatives, produced by a marine filamentous bacterium (Actinomycetales). *Tetrahedron Lett.* 2000; 41 (13): 2073–6.
- Hawas UW, Shaaban M, Shaaban KA, Speitling M, Maier A, Kelter G. Mansouramycins A-D, cytotoxic isoquinolinequinones from a marine Streptomycete. *J Nat Prod.* 2009; 72 (12): 2120–4.
- Hayakawa Y, Shirasaki S, Kawasaki T, Matsuo Y, Adachi K, Shizuri Y. Structures of new cytotoxic antibiotics, piericidins C7 and C8. *J Antibiot.* 2007a; 60 (3): 201–3.
- Hayakawa Y, Shirasaki S, Shiba S, Kawasaki T, Matsuo Y, Adachi K. Piericidins C7 and C8, new cytotoxic antibiotics produced by a marine *Streptomyces* sp. *J Antibiot.* 2007b; 60 (3): 196–200.
- Helaly S E, Pesic A, Fiedler H P, Süßmuth R D. Elaiomycins B and C: alkylhydrazide antibiotics from *Streptomyces* sp. BK 190. *Org Lett.* 2011; 13 (5): 1052–1055.
- Hohmann C, Schneider K, Bruntner C, Brown R, Jones AL, Goodfellow M. Albidopyrone, a new  $\alpha$ -pyrone-

- containing metabolite from marine-derived *Streptomyces* sp. NTK 227. *J Antibiot.* 2009a; 62 (2): 75–9.
- Hohmann C, Schneider K., Bruntner C, Irran E, Nicholson G, Bull AT, Jones AL, Brown R, Stach JE, Goodfellow M, Beil W, Krämer M, Imhoff JF, Süßmuth RD, Fiedler HP. Caboxamycin, a new antibiotic of the benzoxazole family produced by the deep-sea strain *Streptomyces* sp. NTK 937. *J Antibiot.* 2009b; 62: 99-104.
- Hughes CC, MacMillan JB, Gaudêncio SP, Jensen PR, Fenical W. The ammosamides: structures of cell cycle modulators from a marine-derived *Streptomyces* species. *Angew Chem Int Ed Engl.* 2009; 48: 725-727.
- Jensen PR, Williams PG, Oh DC, Zeigler L and Fenical W. Species-specific secondary metabolite production in marine actinomycetes of the genus *Salinispora*. *Appl Environ Microbiol.* 2007; 73: 1146–1152.
- Jeong SY, Shin HJ, Kim TS, Lee HS, Park SK, Kim HM. Streptokordin, a new cytotoxic compound of the methylpyridine class from a marine-derived *Streptomyces* sp. KORDI-3238. *J Antibiot.* 2006; 59 (4): 234–40.
- Kalaitzis JA, Hamano Y, Nilsen G, Moore BS. Biosynthesis and structural revision of neomarinone. *Org Lett.* 2003; 5 (23): 4449–52.
- Kanoh K, Matsuo Y, Adachi K, Imagawa H, Nishizawa M, Shizuri Y. Mechercharmycins A and B, cytotoxic substances from marine-derived *Thermoactinomyces* sp. YM3-251. *J Antibiot.* 2005; 58 (4): 289–92.
- Khalidi N, Seifuddin FT, Turner G, Haft D, Nierman WC, Wolfe KH, Fedorova ND. SMURF: Genomic mapping of fungal secondary metabolite clusters. *Fungal Genet Biol.* 2010; 47: 736–741.
- Kim WG, Kim JP, Kim CJ, Lee KH, Yoo ID. Benzastatins A, B, C, D: New free radical scavengers from *Streptomyces nitrosporeus* 30643. I. Taxonomy, fermentation, isolation and physico-chemical properties and biological activities. *J Antibiot.* 1996; 49: 20-25.
- Kim J, Yi GS. PKMiner: a database for exploring type II polyketide synthases. *BMC Microbiol.* 2012; 12: 169.
- Kornsakulkarn J, Saepua S, Boonruangprapa T, Suphothina S, Thongpanchang C. New  $\beta$ -carboline and indole alkaloids from Actinomycete *Actinomadura* sp. BCC 24717. *Phytochem. Lett.* 2013; 6: 491–494.
- Kwon HC, Kauffman CA, Jensen PR, Fenical W. Marinomycins A-D, antitumor antibiotics of a new structure class from a marine actinomycete of the recently discovered genus "*Marinispora*". *J. Am. Chem. Soc.* 2006; 128: 1622-1632.
- Lee J G, Yoo I D, Kim W G. Differential antiviral activity of benzastatin C and its dechlorinated derivative from *Streptomyces nitrosporeus*. *Biol Pharm Bull.* 2007; 30 (4): 795–797.
- Li F, Maskey RP, Qin S, Sattler I, Fiebig HH, Maier A. Chinikomycins A and B: Isolation, structure elucidation, and biological activity of novel antibiotics from a marine *Streptomyces* sp. isolate M045. *J Nat Prod.* 2005; 68 (3): 349–53.
- Li DH, Zhu TJ, Liu HB, Fang YC, Gu QQ, Zhu WM. Four butenolides are novel cytotoxic compounds isolated from the marine-derived bacterium, *Streptoverticillium luteoverticillatum* 11014. *Arch Pharm Res.* 2006; 29: 624-626.
- Li MH, Ung PM, Zajkowski J, Garneau-Tsodikova S, Sherman DH. Auto-mated genome mining for natural products. *BMC Bioinformatics.* 2009; 10: 185.
- Liu R, Zhu T, Li D, Gu J, Xia W, Fang Y, Liu H, Zhu W, Gu Q. Two indolocarbazole alkaloids with apoptosis activity from a marine-derived actinomycete Z2039-2. *Arch Pharm Res.* 2007; 30: 270-274.
- Lorico A, Long BH. Biochemical characterisation of elsamicin and other coumarin-related antitumour agents as potent inhibitors of human topoisomerase II. *Eur. J. Cancer.* 1993; 29A: 1985-1991.
- Lu J, Ma Y, Liang J, Xing Y, Xi T, Lu Y. Aureolic acids from a marine-derived *Streptomyces* sp. WBF16. *Microbiol Res.* 2012; 167: 590–5.
- Macherla VR, Liu J, Bellows C, Teisan S, Nicholson B, Lam KS. Glaciapyrroles A, B, and C, pyrrolonesquiterpenes from a *Streptomyces* sp. isolated from an Alaskan marine sediment. *J Nat Prod.* 2005; 68 (5): 780–3.
- Manam RR, Teisan S, White DJ, Nicholson B, Grodberg J, Neuteboom ST, Lam KS, Mosca DA, Lloyd GK, Potts BC. Lajollamycin, a nitro-tetraene spiro-beta-lactone-gamma-lactam antibiotic from the marine actinomycete *Streptomyces nodosus*. *J Nat Prod.* 2005; 68: 240-243.
- Martin GD, Tan LT, Jensen PR, Dimayuga RE, Fairchild CR, Raventos-Suarez C, Fenical W. Marmycins A and B, cytotoxic pentacyclic C-glycosides from a marine sediment derived actinomycete related to the genus *Streptomyces*. *J Nat Prod.* 2007; 70: 1406-1409.
- Maskey RP, Li FC, Qin S, Fiebig HH, Laatsch H. Chandrananimycins AC: Production of novel anticancer antibiotics from a marine *Actinomadura* sp. isolate M048 by variation of medium composition and growth conditions. *J Antibiot.* 2003; 56: 622–629.
- Maskey R P, Helmke E, Kayser O, Fiebig H H, Maier A, Busche A. Anti-cancer and antibacterial trioxacarcins with high anti-malaria activity from a marine Streptomycete and their absolute stereochemistry. *J Antibiot.* 2004; 57 (12): 771–779.
- McGovren JP, Neil GL, Crampton SL, Robinson MI, Douros JD. Antitumor activity and preliminary drug disposition studies on chartreusin (NSC 5159). *Cancer Res.* 1977; 37: 1666-1672.

- Medema MH, Blin K, Cimermancic P, de Jager V, Zakrzewski P, Fischbach MA, Weber T, Takano E, Breitling R. antiSMASH: rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. *Nucleic Acids Res.* 2011; 39: W339–W346.
- Miller ED, Kauffman CA, Jensen PR, Fenical W. Piperazimycins: cytotoxic hexadepsipeptides from a marine-derived bacterium of the genus *Streptomyces*. *J Org Chem.* 2007; 72 (2): 323–30.
- Mitchell SS, Nicholson B, Teisan S, Lam KS, Barbara C. Aureoverticillactam, a novel 22-atom macrocyclic lactam from the marine actinomycete *Streptomyces aureoverticillatus*. *J Nat Prod.* 2004; 67 (8): 1400–2.
- Moore B S, Trischman J A, Seng D, Kho D, Jensen P R, Fenical W. Salinamides, anti-inflammatory depsipeptides from a marine streptomycete. *J Org Chem.* 1999; 64 (4): 1145–1150.
- Okazaki T, Kitahara T, Okami Y. Studies on marine microorganisms. IV. A new antibiotic SS-228 Y produced by *Chainia* isolated from shallow sea mud. *J. Antibiot.* 1975; 28: 176-184.
- Park C, Shin HJ, Kim GY, Kwon TK, Nam TJ, Kim SK, Cheong J, Choi IW, Choi YH. Induction of apoptosis by streptochlorin isolated from *Streptomyces* sp. in human leukemic U937 cells. *Toxicol In vitro.* 2008; 22: 1573-1581.
- Pathirana C, Jensen PR, Fenical W. Marinone and debromomarinone: antibiotic sesquiterpenoid naphthoquinones of a new structure class from a marine bacterium. *Tetrahedron Lett.* 1992; 33 (50): 7663–6.
- Pérez M, Crespo C, Schleissner C, Rodríguez P, Zúñiga P, Reyes F, Tartrolon D, a cytotoxic macrodiolide from the marine-derived actinomycete *Streptomyces* sp. MDG-04-17-069. *J Nat Prod.* 2009; 72: 2192–2194.
- Renner MK, Shen Y-C, Cheng X-C, Jensen PR, Frankmoelle W, Kauffman CA. New anti-inflammatory cyclic peptides produced by a marine bacterium (*Streptomyces* sp.). *J Am Chem Soc.* 1999; 121 (49): 11273–6.
- Riedlinger J, Reicke A, Zähler H, Krismer B, Bull AT, Maldonado LA. Abyssomicins, inhibitors of the para-aminobenzoic acid pathway produced by the marine *Verrucosipora* strain AB-18-032. *J Antibiot.* 2004; 57 (4): 271–9.
- Röttig M, Medema MH, Blin K, Weber T, Rausch C, Kohlbacher O. NRP-Spredictor2 – a web server for predicting NRPS adenylation domain specificity. *Nucleic Acids Res.* 2011; 39: W362–W367.
- Sato S, Iwata F, Yamada S, Kawahara H, Katayama M. Usabamycins A–C: new anthramycin-type analogues from a marine-derived actinomycete. *Bioorg Med Chem Lett.* 2011; 21 (23): 7099–101.
- Schneider K, Keller S, Wolter FE, Röglin L, Beil W, Seitz O, Nicholson, G, Bruntner C, Riedlinger J, Fiedler HP, Süßmuth RD. Proximicins A, B, and C-antitumor furan analogues of netropsin from the marine actinomycete *Verrucosipora* induce upregulation of p53 and the cyclin kinase inhibitor p21. *Angew Chem Int Ed Engl.* 2008; 47: 3258-3261.
- Schultz AW, Oh D-C, Carney JR, Williamson RT, Udvary DW, Jensen PR. Biosynthesis and structures of cyclomarins and cyclomarazines, prenylated cyclic peptides of marine actinobacterial origin. *J Am Chem Soc.* 2008; 130 (13): 4507–16.
- Shin HJ, Jeong HS, Lee HS, Park SK, Kim HM, Kwon HJ. Isolation and structure determination of streptochlorin, an antiproliferative agent from a marine-derived *Streptomyces* sp. 04DH110. *J Microbiol Biotechnol.* 2007; 17: 1403-1406.
- Shin DY, Shin HJ, Kim GY, Cheong J, Choi IW, Kim SK, Moon SK, Kang HS, Choi YH. Streptochlorin isolated from *Streptomyces* sp. Induces apoptosis in human hepatocarcinoma cells through a reactive oxygen species-mediated mitochondrial pathway. *J Microbiol Biotechnol.* 2008a; 18: 1862-1868.
- Shin HJ, Kim TS, Lee H-S, Park JY, Choi I-K, Kwon HJ. Streptopyrrolidine, an angiogenesis inhibitor from a marine-derived *Streptomyces* sp. KORDI-3973. *Phytochemistry.* 2008b; 69 (12):2363–6.
- Soria-Mercado IE, Prieto-Davo A, Jensen PR, Fenical W. Antibiotic terpenoid chlorodihydroquinones from a new marine actinomycete. *J Nat Prod.* 2005; 68 (6): 904–910.
- Starcevic A, Zucko J, Simunkovic J, Long PF, Cullum J, Hranueli D. ClustScan: an integrated program package for the semi-automatic annotation of modular biosynthetic gene clusters and in silico prediction of novel chemical structures. *Nucleic Acids Res.* 2008; 36: 6882–6892.
- Stritzke K, Schulz, Laatsch H, Helmke E, Beil W. Novel caprolactones from a marine streptomycete. *J Nat Prod.* 2004; 67: 395–401.
- Takahashi A, Ikeda D, Nakamura H, Naganawa H, Kurasawa S, Okami Y, Takeuchi T, Iitaka Y. Altemicidin, a new acaricidal and antitumor substance. II. Structure determination. *J Antibiot.* 1989a; 42: 1562-1566.
- Takahashi A, Kurasawa S, Ikeda D, Okami Y, Takeuchi T. Altemicidin, a new acaricidal and antitumor substance. I. Taxonomy, fermentation, isolation and physico-chemical and biological properties. *J Antibiot.* 1989b; 42: 1556-1561.
- Umamura M, Koike H, Nagano N, Ishii T, Kawano J, Yamane N, Kozono I, Hori-moto K, Shin-Ya K., Asai K, Yu J, Bennett JW, Machida M. MIDDAS-M: motif-independent de novo detection of secondary metabolite gene clusters through the integration of genome sequencing and transcriptome data. *PLoS ONE.* 2013; 8: e84028.
- van Heel AJ, de Jong A, Montalban-Lopez M, Kok J, Kuipers OP. BAGEL3: automated identification of genes encoding bacteriocins and (non-)bactericidal

- post translationally modified peptides. *Nucleic Acids Res.* 2013; 41, W448–W453.
- Waksman SA. The *Actinomycetes*: Isolation, identification, cultivation and preservation. Baltimore, Williams & Wilkins Company. 1959; 17-28.
- Watve MG, Tickoo R, Jog MM, and Bhole BD. How many antibiotics are produced by the genus *Streptomyces*? *Arch Microbiol.* 2001; 176: 386–390.
- Weber T, Rausch C, Lopez P, Hoof I, Gaykova V, Huson DH, Wohlleben W. CLUSEAN: a computer-based framework for the automated analysis of bacterial secondary metabolite biosynthetic gene clusters. *J Biotechnol.* 2009; 140: 13–17.
- William PG, Asolkar RN, Kondratyuk T, Pezzuto JM, Jensen PR and Fenical W. Saliniketals A and B, bicyclic polyketides from the marine actinomycete *Salinispora arenicola*. *J Nat Prod.* 2007a; 70: 83–88.
- Williams PG, Miller ED, Asolkar RN, Jensen PR, Fenical W. Arenicolides A-C, 26- membered ring macrolides from the marine actinomycete *Salinispora arenicola*. *J Org Chem.* 2007b; 72: 5025-5034.
- Wu SJ, Fotso S, Li F, Qin S, Kelter T, Fiebig HH. 39-N-carboxamidostaurosporine and selina-4(14),7(11)-diene-8,9-diol, new metabolites from a marine *Streptomyces* sp. *J Antibiot.* 2006; 59: 331–7.
- Wu SJ, Fotso S, Li F, Qin S, Laatsch H. Amorphane sesquiterpenes from a marine *Streptomyces* sp. *J Nat Prod.* 2007; 70 (2): 304–6.
- Yadav G, Gokhale RS, Mohanty D. SEARCHPKS: a program for detection and analysis of polyketide synthase domains. *Nucleic Acids Res.* 2003; 31: 3654–3658.
- Zhang Q, Schrader KK, ElSohly HN, Takamatsu S. New cell-cell adhesion inhibitors from *Streptomyces* sp. UMA-044. *J Antibiot.* 2003; 56: 673-681.
- Zhang W, Liu Z, Li S, Yang T, Zhang Q, Ma L, Tian X, Zhang H, Huang C, Zhang S. Spiroindimicins A–D: New Bisindole Alkaloids from a Deep-Sea-Derived Actinomycete. *Org Lett.* 2012; 14: 3364–3367.
- Zhang W, Ma L, Li S, Liu Z, Chen Y, Zhang H, Zhang G, Zhang Q, Tian X, Yuan C. Indimicins A–E, Bisindole Alkaloids from the Deep-Sea-Derived *Streptomyces* sp. SCSIO03032. *J Nat Prod.* 2014; 77: 1887–1892.
- Ziemert N, Podell S, Penn K, Badger JH, Allen E, Jensen PR. The natural product domain seeker NaPDoS: a phylogeny based bioinformatic tool to classify secondary metabolite gene diversity. *PLoS ONE.* 2012; 7: e34064.

Received: January 15, 2016;  
Accepted: April 25, 2016