

Advance Publication

Experimental Animals

Received: 2018.11.28

Accepted: 2019.4.12

J-STAGE Advance Published Date: 2019.5.21

1 **Article Category: Model Animals**

2

3 **The promise of zebrafish as a model of metabolic syndrome**

4

5 Khaled Benchoula¹, Alfi Khatib^{2*}, Ashika Jaffar³, Qamar Udin Ahmed², Wan Mohd Azizi
6 Wan Sulaiman¹, Ridhwan Abd Wahab⁴, and Hesham R. El-Seedi^{5,6}

7

8

9 ¹Department of Basic Medical Sciences, Kulliyyah of Pharmacy, International Islamic
10 University Malaysia, Sultan Ahmad Shah Street, Kuantan 25200, Pahang, Malaysia

11 ²Department of Pharmaceutical Chemistry, Kulliyyah of Pharmacy, International Islamic
12 University Malaysia, Sultan Ahmad Shah Street, Kuantan 25200, Pahang, Malaysia

13 ³School of Biosciences & Technology, VIT University, Vellore 632014, India

14 ⁴Kulliyyah of Allied Health Science, International Islamic University Malaysia, Sultan Ahmad
15 Shah Street, Kuantan 25200, Pahang, Malaysia

16 ⁵Pharmacognosy Group, Department of Medicinal Chemistry, Biomedical Centre, Uppsala
17 University, Box 574, SE-751 23 Uppsala, Sweden.

18 ⁶Alrayan Medical colleges, Medina 42541, Kingdom of Saudi Arabia

19

20

21 *Corresponding author: alfikhatib@iium.edu.my.

22 Department of Pharmaceutical Chemistry, Kulliyyah of Pharmacy International Islamic
23 University Malaysia. Sultan Ahmad Shah Street, Bandar Indera Mahkota, 25200 Kuantan,
24 Pahang, Malaysia

25

26

27

Abstract

Metabolic syndrome is a cluster including hyperglycaemia, obesity, hypertension, and hypertriglyceridaemia as a result of biochemical and physiological alterations and can increase the risk of cardiovascular disease and diabetes. Fundamental research on this disease requires validated animal models. One potential animal model that is rapidly gaining in popularity is zebrafish (*Danio rerio*). The use of zebrafish as an animal model conveys several advantages, including high human genetic homology, transparent embryos and larvae that allow easier visualization. This review discusses how zebrafish models contribute to the development of metabolic syndrome studies. Different diseases in the cluster of metabolic syndrome, such as hyperglycaemia, obesity, diabetes, and hypertriglyceridaemia, have been successfully studied using zebrafish; and the model is promising for hypertension and cardiovascular metabolic-related diseases due to its genetic similarity to mammals. Genetic mutation, chemical induction, and dietary alteration are among the tools used to improve zebrafish models. This field is expanding, and thus, more effective and efficient techniques are currently developed to fulfil the increasing demand for thorough investigations.

Key words: zebrafish; metabolic syndrome; obesity; diabetes; and hypertension.

1 **Introduction**

2 Since the 1970s, the use of *Danio rerio*, commonly known as zebrafish, as an animal model
3 has spread widely among various universities and research centres. The model has shed new
4 light into various research fields. George Streisinger, a molecular biologist from the
5 University of Oregon, was the first scientist to use zebrafish as a tool to study the nervous
6 system. This vertebrate animal was found to be less complicated than mice [16]. The
7 zebrafish is an animal model that was established to be an ideal experimental model for
8 several biological disorders [20].

9 The use of zebrafish as an animal model provides several advantages. Housing is less
10 cumbersome relative to other animal models, allowing cost efficiency. The rapid egg
11 production and early developmental morphology of this small organism make it one of the
12 favourite animal models for drug screening. One mating can produce approximately 300 eggs
13 in each conjugation in one night. The zebrafish embryo can reach the early larval stage after 3
14 days' post-fertilisation (dpf) and become an adult at 3 months' post-fertilisation. Their small
15 eggs can be placed into the wells of plates and survive for approximately 7 days without
16 feeding or changing the water. These characteristics are beneficial, as several drugs can be
17 screened in a short period of time. Furthermore, the size of this animal, which is relatively
18 small, reduces the quantity of tested drug/plant extract, which makes it possible to test a small
19 amount of compound(s). Zebrafish share 70% homology to mammals and 84% similarity to
20 mammalian disease genes. Gene similarity leads to the use of zebrafish as a model for
21 studying gene functions and mammalian disorders, such as metabolic syndromes [23, 21].
22 Oka et al. [48] reported that the zebrafish shares similar pathological characteristics with
23 humans since the same genes are affected, including *IL-6*, *IL-1 β* and *APOH* in the
24 coagulation pathway and *SREBF1*, *PPAR α/γ* , *NR1H3* and *LEP* in lipid metabolism. In
25 particular, zebrafish embryos and larvae have similarly coded genes for carbohydrate

1 metabolism to mammals [78]. In addition, zebrafish larvae have adipocytes that store lipids
2 and other specific cells involved in lipid metabolism [26]. The pancreas of zebrafish also has
3 a structure similar to that of mammals, with two parts: endocrine and exocrine. The endocrine
4 cells of zebrafish were found to have three types of cells, α , β , and δ , which release insulin,
5 glucagon and somatostatin, respectively. The exocrine gland contains three types of cells,
6 ductal cells, acinar cells, and centroacinar cells [8, 42]. This discovery has initiated the use of
7 zebrafish as a model for studying different metabolic disorders associated with glucose
8 metabolism.

9 Nevertheless, apart from these advantages, the solubility of testing compounds is an
10 obstacle for certain studies using zebrafish embryo, whereby the tested compounds or drug
11 are preferably water soluble, or at least soluble in 1% DMSO or acetone [3]. However, it
12 will not be a limiting factor when an adult zebrafish is used since force-feeding can be
13 applied and enabling the evaluation of whole part of the sample. Apart from it, the volume of
14 zebrafish blood is very limited; however, it can be overcome through the use of high-end
15 analytical instrument such liquid-chromatography tandem mass-spectrometry (LC-MS) for
16 analysis of the metabolites. This technique is very sensitive enabling analysis of small
17 quantity of the sample. LC-MS time of flight (LC-MS-TOF) has been utilized extensively in
18 metabolomics and proteomics approach using zebrafish [71, 12]. Finally, the small size of
19 zebrafish causes difficulties in dissection and thus requires more effort and skill to extract the
20 organs compared to other conventional animal models. Although certain organs in zebrafish
21 are very fragile, the isolation of those organs for histology is possible [80].

22

23

24

1 **Metabolic syndrome**

2 In 1988, Gerald Reaven was the first scientist who gave the term syndrome X to various
3 clusters of abnormal disorders [17]. Since then, syndrome X has been the focus of many
4 researchers and defined differently. Later, the term syndrome X was changed to metabolic
5 syndrome, although the definition of metabolic syndrome varies and is still unclear.
6 According to the International Diabetes Federation (IDF) definition, the patient at risk of
7 metabolic syndrome has been diagnosed with obesity and at least two of the four following
8 disorders: high fasting blood glucose (≥ 100 mg/dL), high blood pressure (systolic BP ≥ 130
9 or diastolic BP ≥ 85 mmHg), high triglycerides (≥ 150 mg/dL), and a decrease in high-
10 density lipoprotein (HDL < 40 mg/dL in males, < 50 mg/dL in females) [69]. However, the
11 World Health Organization (WHO) has suggested different definitions, i.e., metabolic
12 syndrome is an abnormal glucose level with at least two of the following health problems:
13 high arterial pressure ($\geq 140/90$ mmHg), high plasma triglyceride (≥ 150 mg/dL), abdominal
14 obesity body mass index (BMI > 30 kg/m²) and microalbuminuria [67]. Notably, the
15 European Group for the Study of Insulin Resistance (EGIR) changed the term to insulin
16 resistance syndrome. According to the EGIR, a person with insulin resistance syndrome has a
17 high level of plasma insulin with two of the following factors: raised plasma glucose but not
18 diabetic, raised triglycerides (≥ 150 mg/dL) and/or reduced high-density lipoproteins (HDL $<$
19 110 mg/dL), high blood pressure ($> 130/85$ mmHg), and hypertriglyceridaemia (≥ 150
20 mg/dL) with a low amount of HDL in both genders. According to all mentioned definitions,
21 the four diseases (diabetes mellitus, high blood pressure, obesity, and high triglyceride levels)
22 are clearly shown to be related to each other [27].

23

24

1 **Zebrafish as a model to study metabolic syndrome**

2 *Zebrafish as a diabetic model*

3 Diabetes mellitus (DM) is a prolonged state of high blood glucose levels. A great deal of
4 complexity is associated with this metabolic illness and is primarily manifested among DM
5 patients [79]. The hormones that are responsible for maintaining the blood glucose level
6 within the optimal range are insulin, glucagon, and somatostatin. These hormones are
7 released from the endocrine part of the pancreas [55]. The similarity between the zebrafish
8 and human pancreas has been avidly studied. The zebrafish pancreas can be divided into two
9 important parts: the exocrine and endocrine segments. The exocrine segment contains three
10 major types of cells: ductal, acinar, and centroacinar cells. In contrast, the endocrine part is
11 separated into islets that contain three types of cells: beta cells that produce insulin, alpha
12 cells that secrete glucagon, and delta cells that give rise to somatostatin [8, 42]. Thus, from
13 this point, humans and zebrafish can be concluded to be morphologically comparable.
14 Similarly, the normal blood glucose level in zebrafish (50 – 75 mg/dL) is close to the normal
15 human blood glucose level range (70 – 120 mg/dL) [75].

16 PCR analyses of both adult and larval zebrafish showed that the same genes that are
17 regulated by carbohydrates have been detected in mammals. The pancreas of zebrafish is
18 largely composed of exocrine tissue organized in acini. Likewise, it has the same function
19 and structure as the mammalian counterpart in glucose homeostasis [28]. Zebrafish have the
20 ability to transcribe all genes related to gluconeogenesis and lipolysis after 4 dpf, such as
21 cytosolic phosphoenol pyruvate carboxykinase, glucose-6-phosphatase, fatty acid synthase,
22 acetyl-coA carboxylase, glucose-6-phosphate dehydrogenase, glycogen synthase, and
23 glycogen phosphorylase [13]. At 4 dpf, pancreas is still immature but it is functioning
24 [40]. To exploit this advantage, the larvae were treated with an overdose of insulin, resulting

1 in insulin resistance [38]. Numerous investigations have documented the relationship
2 between insulin resistance and obesity in metabolic syndrome. Forn-Cuní et al. [14] reported
3 the intersection between the non-alcoholic fatty liver of zebrafish and humans. Both zebrafish
4 liver genes of *atp5e* and *atp5gb3b* play a significant role in insulin resistance associated with
5 obesity. Moreover, Malle et al. [35] reported that systemic monovalent 1 (MV1) can activate
6 NF- κ B-inducing kinase (NIK). The NF- κ B pathway is associated with β cell dysfunction,
7 which causes insulin resistance. Song and Cone [66] reported that overexpression of agouti-
8 related protein (*AgRP*) increases the energy storage rate, causing overweight zebrafish.

9 Essentially, the most difficult part of inducing diabetes in zebrafish is the rapid
10 recovery of β cells. Because of β cell damage, centroacinar cells are stimulated to multiply
11 and separate into β cells. In addition, α cells can also contribute to supplanting β cells [4].
12 Although numerous studies discussed diabetes in zebrafish before 2007, the primary trial for
13 the ablation (removal) of β cells from the pancreas of zebrafish was carried out by Pisharath
14 et al. [51]. Indeed, the objective of this study was primarily to establish a zebrafish diabetic
15 model to test the ability of zebrafish as a diabetic model.

16 Table 1 presents the various studies that have used zebrafish as a model for obesity
17 and diabetes, which are related directly to metabolic syndrome. The food, environmental, and
18 chemical inducers and their doses are indicated in Table 1. Pisharath et al. [51] submerged
19 zebrafish larvae in metronidazole with the goal of ablating the β cells. The outcome
20 demonstrated that at 10 mM, the drug can completely decimate the β cells. Around the same
21 time, another study took one step forward by the work carried out by Elo et al. [13], who
22 demonstrated that the antidiabetic drug glipizide can decrease the glucose level in adult
23 zebrafish after exposure to 25% glucose in Me₂SO. The main diabetic model utilising
24 zebrafish was conceived by Gleeson et al. [19] when they screened diverse groups of
25 glucose-utilising adult zebrafish. The blood glucose level was significantly increased after

1 immersing the adult zebrafish in the glucose solution. Connaughton et al. [10] increased the
2 duration of the hyperglycaemic state of adult zebrafish from 2 to 8 months by increasing the
3 glucose solution from 1 to 3% and the exposure time to two months. Similarly, alloxan
4 monohydrate has demonstrated a successful outcome in raising the blood glucose level.
5 Submerging the fish in 300 mg alloxan/100 mL H₂O and 400 mg alloxan/100 mL H₂O
6 solution for 30 minutes as well as treatment with 1% glucose solution led to an augmentation
7 of blood glucose level [64].

8 Routinely, to induce diabetes in rodents, researchers utilised diabetogenic
9 medications, for example, alloxan or streptozotocin (STZ). Olsen et al. [49] developed
10 another convention when 0.3% STZ solution was infused into zebrafish at 350 mg/kg body
11 weight. The STZ-injected fish showed altered glucose levels, certain diabetes complications
12 reported in this investigation, and increased serum non-enzymatic glycated protein levels
13 compared to the healthy fish. However, insulin levels increased by 80%. More profound, a
14 high blood glucose level influences the limb regeneration of zebrafish. This outcome can be
15 seen following two weeks of high-glucose exposure. Later, the same research group
16 recognised 71 factors hindering the caudal balance recovery of diabetic zebrafish [58].
17 Although the two investigations focused on multiple aspects of the zebrafish diabetic model,
18 a need for a clear STZ injection procedure was demonstrated. Therefore, Intine et al. [25]
19 gave a full clarification of the protocol for establishing a diabetes model by infusing 5
20 injections (350 mg/kg BW) of STZ over 3 weeks. Additionally, 5% of mortality was found in
21 this investigation. Moss et al. [45] also used STZ to observe the regeneration process of
22 zebrafish pancreas. This study further demonstrated that STZ can effectively destroy β cells.

23 Typically, high-fat and glucose regimens are the typical approaches used to induce
24 type 2 diabetes in rodents. Accordingly, a similar method was used to induce type 2 diabetes
25 in zebrafish. The overfeeding of adult zebrafish with a high-fat diet promotes certain health

1 issues, including hyperglycaemia and fat accumulation in the liver [34]. Similarly,
2 overfeeding with normal zebrafish food for 8 weeks/6 feeds/day promoted increases in
3 glucose and insulin. In addition, overfeeding induces impaired tolerance to glucose.
4 Treatment with metformin and glimepiride could reverse the damaging effect of overfeeding
5 in the body of zebrafish [76]. Furthermore, the high-glucose diet increases the levels of
6 insulin, glucagon, and phosphoenolpyruvate carboxykinase. These 3 metabolites are directly
7 correlated with type 2 diabetes [13].

8 Notably, the non-enzymatic glycated proteins are initially more pronounced when the
9 glucose level is elevated in the circulatory system. Fructosamine is one of these glycated
10 proteins whose level was raised by 41% in a diabetic zebrafish model. The focal neurological
11 system complexity related to diabetes was a part of this investigation [6]. This conclusion
12 was reached after an experiment in which the fish were immersed in 111 mM glucose for 14
13 days. The results demonstrated that the action of acetylcholinesterase had increased in the
14 diabetic fish, unlike the control aggregate, and acetylcholinesterase was the main focus of this
15 research to investigate the impact of diabetes on the cerebrum. This protein is situated in the
16 myoneural intersection and separates acetylcholine and different esters. Acetylcholinesterase
17 alteration in diabetes can cause certain neurological maladies [7]. A similar convention was
18 employed by Alvarez et al. [1], who examined the impact of hyperglycaemia and its relation
19 to retinopathy. The retinal vessels in 2-year-old fish were apparently thickened by
20 aggravation of retinal vein obstruction. Related to diabetic complications, hyperglycaemia
21 influences the level of cortisol in zebrafish hatchlings. Therefore, the diabetic zebrafish
22 model provides better insight into the disorder. However, further diabetes studies in zebrafish
23 are still warranted to derive further key information [52].

24

1 *Zebrafish as an obesity and hyperlipidaemia model*

2 Obesity is a metabolic disorder that is becoming highly prevalent in developed and
3 developing countries. This lipid metabolism disorder disrupts the energy balance and is
4 associated with genes and environment [54]. Previously, obesity was most likely associated
5 with different genes, such as those encoding the leptin receptor, melanocortin-4-receptor
6 *Mc4r*, pro-hormone convertase 1, and pro-opiomelanocortin genes [31]. The increasing
7 prevalence of obesity has switched the focus of genotyping from a single gene to polygenic
8 obesity [2]. Genome-wide association studies (GWASs) aided in creating the human obesity
9 genome map, and the 12th update indicated the involvement of 253 quantitative trait loci [54].
10 The recent trend of obesity has brought the role of the environment into the light. Therefore,
11 obesity in humans is often researched based on monozygotic and dizygotic twin studies [11].

12 Energy balance and metabolism are related to the leptin and insulin pathways, and are
13 thus intercalated with other metabolic disorders. The brain and hormonal activities also
14 affected food uptake and in turn, contributed to metabolic disorder [9]. Obesity is detrimental
15 on a large scale as it increases the risk of other metabolic disorders such as hyperlipidaemia,
16 cardiovascular diseases, type 2 diabetes, and cancer [15].

17 The increase in obesity has further highlighted the need to develop animal models.
18 Monogenic and polygenic mutations have been used in rodent models. Mouse and rat models
19 are the most common obesity models with a single gene causing a mutation in leptin, the
20 leptin receptor, and insulin signalling using *ob/ob* and *db/db* mice [22], as well as Zucker
21 *fa/fa* and Wistar-Kyoto rat models [39].

22 Polygenic models have been studied through diet-induced and age-related obesity
23 [29]. Western human and high-fat diets have been found to affect leptin and insulin signalling
24 in rats. These protocols show the role of the hypothalamus during obesity development [18].

1 Various mouse models, including New Zealand obese mice, elucidated the type 2 diabetes
2 state and revealed its linkage to metabolic disorders [61]. The age-related obesity model is
3 more suitable for mice, because it mimics the late onset of obesity in humans [77]. In the
4 human obesity genome map, 119 genes have been recognised to be associated with obesity.
5 Therefore, various genetic modification models in animals have been made [74]. The rodents
6 are also often exposed to mutagens by radiation to alter genes related to adiposity for a faster
7 random model, but this procedure is often not economical and does not help characterise
8 energy homeostasis as a basis of measurement [24].

9 Non-rodent models, namely, chickens and pigs, are less common but have also been
10 induced through high-fat diets [44]. Obesity is often diagnosed in dogs, which could be an
11 upcoming model [50]. Non-human primates are evolutionarily closer, defining a role of
12 epigenetics; thus, they have also helped in such studies, but a further exploration into the role
13 of the brain in energy homeostasis and food intake was not investigated. Ethical
14 complications, cumbersome processes, and high cost and maintenance have caused
15 researchers to consider other models. Zebrafish, as a member of the lower metazoan family,
16 are similar in terms of organs and adipose tissues [62]. It also has similar signalling pathways
17 of leptin [43] and melanocortin [21] linked to the neuronal endocrine pathways [48].

18 As observed in humans, the mutation of *Mc4r* results in the obese state, whereby the
19 same gene also plays a prime role in zebrafish. The zebrafish model has been used to study
20 the genes involved in obesity, revealing the overexpression of *zAgRP*, which acts as an
21 antagonist to the melanocortin receptor and in turn causes adipocyte hypertrophy [66]. The
22 overexpression of genes related to insulin signalling *Akt1* is able to induce obesity. This
23 finding has allowed a novel anti-obesity study at the larval zebrafish stage [57]. Furthermore,
24 chemical mutagens such as rosiglitazone, phenylephrine, and T0070907 at varying
25 concentrations, along with diet, have been used to create a larval zebrafish model for studying

1 this metabolic disorder successfully, as shown in Table 1. Beyond chemical mutagens,
2 interesting new insights into the environmental role in obesity have been found, and a group
3 of researchers have obtained promising results by altering the circadian rhythm [68]. Kopp et
4 al. [32] found that prolonged exposure of the larval zebrafish to light increases the number of
5 adipocytes by seven-fold, but it did not affect fatty acids.

6 As seen in rodents, the zebrafish model of diet-induced obesity is also an upcoming
7 option focusing on the external factors contributing to obesity. Overfeeding of high-fat foods,
8 such as artemia, 60 mg/day for 8 weeks, to adult zebrafish has been proven to play a role in
9 the lipid metabolism pathway. This finding was deduced by comparing diet-induced obesity
10 mouse and rat models and humans [48]. A fat-rich diet model of adult zebrafish has also been
11 used for natural anti-obesity compound studies. Meguro et al. [41] tested the effect of green
12 tea extracts on the diet-induced obesity model. The zebrafish model of obesity is an excellent
13 alternative to the previous study models and is expected to be exploited further, as studies
14 will utilise this model because it is beneficial, informative, and cost-effective.

15

16 *Zebra fish as a high blood pressure model*

17 High blood pressure or hypertension is a noteworthy general medical issue.
18 Hypertension is characterised by high volumes of blood in the blood supply routes, where the
19 blood flows between the heart and every organ in the body. Blood pressure can be grouped
20 into systolic and diastolic pressures. The estimation of the pulse is communicated in terms of
21 systolic pressure (when the heart pumps the blood) over diastolic pressure (between the heart
22 beats). Hypertension management is not completely understood; consequently, further studies
23 are greatly needed, which has given rise to a newly found commitment to manage
24 hypertension in a better way.

1 The renin–angiotensin–aldosterone system (RAAS) is the hormone system that
2 regulates blood pressure. As the blood flows, high sodium levels affect the circulation. In this
3 circumstance, the renin that is discharged from the kidney alters the conversion of
4 angiotensin I (Ang I) to angiotensin II (Ang II) by the enzyme called angiotensin-converting
5 enzyme (ACE), which is secreted from the lung [63]. The increase in Ang II causes the
6 tubular epithelial cells of the adrenal gland to discharge more aldosterone, promoting the
7 reabsorption of sodium particles from tubular liquid and excreting potassium particles in the
8 urine. In vascular endothelial cells, Ang II can cause vasoconstriction of arterioles by
9 reducing the synthesis of nitric oxide (NO) [70, 60]. ACE plays another role in inhibiting the
10 production of NO by breaking down bradykinin, which is required for the synthesis of NO
11 [69]. The pituitary gland is also affected by Ang II through its binding to angiotensin II
12 receptor type-1 (AT₁), causing the reabsorption of water in kidneys and triggering the
13 stimulus for thirst in the cerebrum. This process results in urging the individual to drink more
14 water, contributing to the decreased concentration of salt in the blood [53]. The renin gene
15 was reported in the hereditary material of zebrafish, and it begins to appear 24-hours post-
16 fertilisation and exhibits 53% similarity to human renin [56].

17 NOSTRIN is another gene associated indirectly with high blood pressure and is
18 expressed in zebrafish. The knockdown of this gene can affect the retinal blood vessels by
19 increasing the clearance of protein from the serum in the vessels, resulting in obvious damage
20 to both glomerular endothelial cells and the glomerular basement membrane in the kidney.
21 Furthermore, this damage alters salt absorption into blood [30].

22 Another system regulating blood pressure was reported by Marek-Trzonkowska et al.
23 [37], who demonstrated a connection between hypertension and angiogenesis. Angiogenesis
24 is the development of new blood vessels, and the initiation of this procedure could reduce
25 blood pressure [46]. The restraint of delta-like 4 (DII4)-Notch by fortification of microRNA-

1 30a (miR-30a) can prompt angiogenesis in zebrafish [65]. The flowers of *Panax notoginseng*
2 were found to be able to repair the damaged blood vessels by accelerating angiogenesis in
3 zebrafish embryos [73]. This plant is utilised as a part of traditional Chinese medicine for the
4 treatment of hypertension [72]. In a related study, Lai et al. [33] discovered a new gene that
5 can control angiogenesis. They found that the knockdown of the WNK lysine-deficient
6 protein kinase 1 (*WNK1*) gene in zebrafish can affect the angiogenesis process.
7 Overexpression of this gene was reported to increase blood pressure in humans [47].
8 Recently, a coiled-coil domain containing 80 (*CCDC80*) was identified as the gene associated
9 with pulmonary arterial hypertension, and this gene was discovered to exist in the genetic
10 material of zebrafish. The knockout of this gene resulted in inhibition of NO synthesis,
11 further contributing to the increase in the size of the arterial blood supply. The effect of NO
12 synthesis inhibition can be seen through the decrease in diameter of the ventral artery.
13 Moreover, the regulation of NO synthesis relies upon the cGMP-dependent protein kinase.
14 These outcomes vividly demonstrate that the zebrafish has a similar pulmonary artery
15 regulatory process to that observed in mammals [59]. Although zebrafish cardiovascular
16 models-related metabolic syndrome has not been completely established, the evidential
17 genetic similarity of zebrafish to mammals will support the establishment of the model
18 through genetic mutation, chemical induction, and dietary alteration.

19

20 **Conclusion**

21 The zebrafish model is on the rise due to its good reproducibility, simplicity, and cost
22 effectiveness. Models for metabolic syndrome, including hyperglycaemia, obesity, and
23 hyperlipidaemia, have been developed through different methods of induction, such as
24 genetic mutation, chemical induction, or dietary alteration. Although zebrafish model related

1 to hypertension and cardiovascular disease has not been completely established, it is
2 promising to be developed due to its genetic similarity to mammals.

3

4 **Acknowledgement**

5 The authors wish to thank the International Islamic University Malaysia for Publication
6 Research Initiative Grant Scheme funding (PRIGS18-027-0027) and extend their
7 appreciation to VR (Stockholm, Sweden) Grant number 2016-05908 for generous financial
8 support awarded to H.R.E.

9

10 **References**

- 11 1. Alvarez, Y., Chen, K., Reynolds, A. L., Waghorne, N., O'Connor, J. J. and Kennedy,
12 B. N. 2010. Predominant cone photoreceptor dysfunction in a hyperglycaemic model
13 of non-proliferative diabetic retinopathy. *Dis. Model. Mech.* 3: 236–45.
- 14 2. Apal Sammy, Y. D. and Mohamed, Z. 2015. Obesity and genomics: role of technology
15 in unraveling the complex genetic architecture of obesity. *Hum. Genet.* 134: 361–74.
- 16 3. Beekhuijzen, M., de Koning, C., Flores-Guillén, M. E., de Vries-Buitenweg, S.,
17 Tobor-Kaplon, M., van de Waart, B. and Emmen, H. 2015. From cutting edge to
18 guideline: a first step in harmonization of the zebrafish embryotoxicity test (ZET) by
19 describing the most optimal test conditions and morphology scoring system. *Reprod.*
20 *Toxicol.* 56: 64–76.
- 21 4. Beer, R. L., Parsons, M. J. and Rovira, M. 2016. Centroacinar cells: at the center of
22 pancreas regeneration. *Dev. Biol.* 413: 8–15.

- 1 5. Bray, M. S. and Young, M. E. 2007. Circadian rhythms in the development of
2 obesity: potential role for the circadian clock within the adipocyte. *Obes. Rev.* 8: 169–
3 81.
- 4 6. Capiotti, K. M., Antonioli, R., Kist, L. W., Bogo, M. R., Bonan, C. D. and Da Silva,
5 R.S. 2014. Persistent impaired glucose metabolism in a zebrafish hyperglycemia
6 model. *Biochem. Mol. Biol.* 2014. 171: 58–65.
- 7 7. Capiotti, K. M., Siebel, A. M., Kist, L. W., Bogo, M. R., Bonan, C. D. and Da Silva,
8 R. S. 2016. Hyperglycemia alters E-NTPDases, ecto-5'-nucleotidase, and ectosolic
9 and cytosolic adenosine deaminase activities and expression from encephala of adult
10 zebrafish (*Danio rerio*). *Purinergic. Signal.* 12: 211–20.
- 11 8. Chen, S., Li, C., Yuan, G. and Xie, F. 2007. Anatomical and histological observation
12 on the pancreas in adult zebrafish. *Pancreas* 34: 120–125.
- 13 9. Claflin, K. E. and Grobe, J. L. 2015. Control of energy balance by the brain renin-
14 angiotensin system. *Curr. Hypertens. Rep.* 17: 38.
- 15 10. Connaughton, V. P., Baker, C., Fonde, L., Gerardi, E. and Slack, C. 2016. Alternate
16 immersion in an external glucose solution differentially affects blood sugar values in
17 older versus younger zebrafish adults. *Zebrafish* 13: 87–94.
- 18 11. Dario, A. B., Ferreira, M. L., Refshauge, K. M., Lima, T. S., Ordoñana, J. R. and
19 Ferreira, P. H. 2015. The relationship between obesity, low back pain, and lumbar
20 disc degeneration when genetics and the environment are considered: a systematic
21 review of twin studies. *Spine. J.* 15: 1106–17.
- 22 12. De Sotto, R. B., Medriano, C. D., Cho, Y., Kim, H., Chung, I.-Y., Seok, K.-W., Song,
23 K. G., Hong, S. W., Park, Y., Kim, D. 2017. Sub-lethal pharmaceutical hazard
24 tracking in adult zebrafish using untargeted LC–MS environmental metabolomics. *J.*
25 *Hazard. Mater.* 339: 63-72.

- 1 13. Elo, B., Villano, C. M., Govorko, D. and White, L. A. 2007. Larval zebrafish as a
2 model for glucose metabolism: Expression of phosphoenolpyruvate carboxykinase as
3 a marker for exposure to anti-diabetic compounds. *J. Mol. Endocrinol.* 38: 433–40.
- 4 14. Forn-Cuní, G., Varela, M., Fernández-Rodríguez, C. M., Figueras, A. and Novoa, B.
5 2015. Liver immune responses to inflammatory stimuli in a diet-induced obesity model
6 of zebrafish. *J. Endocrinol.* 224: 159–70.
- 7 15. Furukawa, S., Fujita, T., Shimabukuro, M., Iwaki, M., Yamada, Y., Nakajima, Y.,
8 Nakayama, O., Makishima, M., Matsuda, M. and Shimomura, I. 2017. Increased
9 oxidative stress in obesity and its impact on metabolic syndrome. *J. Clin. Invest.* 114:
10 1752–61.
- 11 16. Gemberling, M., Bailey, T. J., Hyde, D. R. and Poss, K. D. 2013. The zebrafish as a
12 model for complex tissue regeneration. *Trends. Genet.* 29: 611–20.
- 13 17. Gerald, M. R. 2001. Syndrome X: A Short History. *Ochsner. J.* 3: 124- 125.
- 14 18. Git, K. C. G. and Adan, R. A. H. 2015. Leptin resistance in diet-induced obesity: the
15 role of hypothalamic inflammation. *Obes. Rev.* 16:207–24.
- 16 19. Gleeson, M., Connaughton, V. and Arneson, L. S. 2007. Induction of hyperglycaemia
17 in zebrafish (*Danio rerio*) leads to morphological changes in the retina. *Acta Diabetol.*
18 44: 157–63.
- 19 20. Graham, J. L., and Peter, D. C., 2007. Animal models of human disease: zebrafish
20 swim into view. *Nat. Rev. Genet.* 8: 353–367.
- 21 21. Guillot, R., Cortés, R., Navarro, S., Mischitelli, M., García-Herranz, V., Sánchez, E.,
22 Cal, L., Navarro, J. C., Míguez, J. M., and Afanasyev, S. 2016. Behind
23 melanocortinantagonist overexpression in the zebrafish brain: a behavioral and
24 transcriptomic approach. *Horm. Behav.* 82: 87–100.

- 1 22. Hao, Z., Münzberg, H., Rezai-Zadeh, K., Keenan, M., Coulon, D., Lu, H., Berthoud,
2 H. R. and Ye, J. 2015. Leptin deficient ob/ob mice and diet-induced obese mice
3 responded differently to Roux-en-Y bypass surgery. *Int. J. Obes.* 39: 798–805.
- 4 23. Harper, C. and Lawrence, C. 2016. The laboratory zebrafish. CRC Press, Boca Raton,
5 FL.
- 6 24. Imaoka, T., Nishimura, M., Daino, K., Morioka, T., Nishimura, Y., Uemura, H.,
7 Akimoto, K., Furukawa, Y., Fukushi, M. and Wakabayashi, K. 2016. A rat model to
8 study the effects of diet-induced obesity on radiation-induced mammary
9 carcinogenesis. *Radiat. Res.* 185: 505–15.
- 10 25. Intine, R. V., Olsen, A. S. and Sarras, M. P. 2013. A Zebrafish Model of Diabetes
11 Mellitus and Metabolic Memory. *J. Vis. Exp.* 72: 1–7.
- 12 26. Jennifer, L., Anderson, J. L., Carten, J. D. and Farber, S. A. 2011. Zebrafish Lipid
13 Metabolism: From Mediating Early Patterning to the Metabolism of Dietary Fat and
14 Cholesterol. *Method. Cell. Biol.* 2011. 111–141.
- 15 27. Khosravi-Boroujeni, H., Ahmed, F., Sadeghi, M., Roohafza, H., Talaei, M.,
16 Dianatkhah, M., Pourmogaddas, A. and Sarrafzadegan, N. 2015. Does the impact of
17 metabolic syndrome on cardiovascular events vary by using different definitions?
18 *BMC Public Health* 15: 1313.
- 19 28. Kimmel, R. A. and Meyer, D. 2016. Zebrafish pancreas as a model for development
20 and disease. *Method. Cell. Biol.* 134: 431–61.
- 21 29. King, A. and Bowe, J. 2016. Animal models for diabetes: Understanding the
22 pathogenesis and finding new treatments. *Biochem. Pharmacol.* 99: 1–10.
- 23 30. Kirsch, T., Kaufeld, J., Korstanje, R., Hentschel, D. M., Staggs, L., Bollig, F., Beese,
24 M., Schroder, P., Boehme, L. and Haller, H. 2013. Knockdown of the hypertension-

- 1 associated gene NOSTRIN alters glomerular barrier function in zebrafish (*Danio*
2 *rerio*) novelty and significance. *Hypertension* 62: 726–30.
- 3 31. Klaauw van der, A. A. and Farooqi, I. S. 2015. The hunger genes: pathways to
4 obesity. *Cell* 161: 119–32.
- 5 32. Kopp, R., Billecke, N., Legradi, J., den Broeder, M., Parekh, S. H. and Legler, J.
6 2016. Bringing obesity to light: Rev-erb [alpha], a central player in light-induced
7 adipogenesis in the zebrafish? *Int. J. Obes.* 40: 824–33.
- 8 33. Lai, J. G., Tsai, S. M., Tu, H. C., Chen, W. C., Kou, F. J., Lu, J. W., Wang, H-D.,
9 Huang, C-L. and Yuh, C-H. 2014. Zebrafish WNK lysine deficient protein kinase 1
10 (*wnk1*) affects angiogenesis associated with VEGF signaling. *PLoS One.* 9: e106129.
- 11 34. Landgraf, K., Schuster, S., Meusel, A., Garten, A., Riemer, T., Schleinitz, D., Kiess,
12 W. and Körner, A. 2017. Short-term overfeeding of zebrafish with normal or high-fat
13 diet as a model for the development of metabolically healthy versus unhealthy
14 obesity. *BMC. Phys.* 17: 4.
- 15 35. Malle, E. K., Zammit, N. W., Walters, S. N., Koay, Y. C., Wu, J., Tan, B. M.,
16 Villanueva, J. E., Brink, R., Loudovaris, T., Cantley, J., McAlpine, S. R., Hesselton,
17 D. and Grey, S. T. 2015. Nuclear factor kappa β -inducing kinase activation as a
18 mechanism of pancreatic beta cell failure in obesity. *J. Exp. Med.* 212: 1239–54.
- 19 36. Mamenko, M., Zaika, O. and Pochynyuk, O. 2014. Direct regulation of ENaC by
20 bradykinin in the distal nephron. Implications for renal sodium handling. *Curr. Opin.*
21 *Nephrol. Hypertens.* 23: 122-129.
- 22 37. Marek-Trzonkowska, N., Kwieczyńska, A., Reiwer-Gostomska, M., Koliński, T.,
23 Molisz A. and Siebert, J. 2015. Arterial hypertension is characterized by imbalance of
24 pro-angiogenic versus anti-angiogenic factors. *PLoS One.* 10: e0126190.

- 1 38. Marín-Juez, R., Jong-Raadsen, S., Yang, S. and Spaink, H. P. 2014. Hyperinsulinemia
2 induces insulin resistance and immune suppression via Ptpn6/Shp1 in zebrafish. *J.*
3 *Endocrinol.* 222: 229–41.
- 4 39. Marques, C., Meireles, M., Norberto, S., Leite, J., Freitas, J., Pestana, D., Faria, A.
5 and Calhau, C. 2016. High-fat diet-induced obesity Rat model: a comparison between
6 Wistar and Sprague-Dawley Rat. *Adipocyte* 5: 11–21.
- 7 40. Matsuda, H., Mullapudi, S. T., Zhang, Y., Hesselton, D., and Stainier, D. Y. R. 2017.
8 Thyroid hormone coordinates pancreatic islet maturation during the zebrafish larval to
9 juvenile transition to maintain glucose homeostasis. *Diabetes* 66: 2623-2635.
- 10 41. Meguro, S., Hasumura, T. and Hase, T. 2015. Body fat accumulation in zebrafish is
11 induced by a diet rich in fat and reduced by supplementation with green tea extract.
12 *PLoS One.* 10: e0120142.
- 13 42. Menke, A. L., Spitsbergen, J. M., Wolterbeek, A. P. M. and Woutersen, R. A. 2011.
14 Normal Anatomy and Histology of the Adult Zebrafish. *Toxicol. Pathol.* 39: 759–775.
- 15 43. Michel, M., Page-McCaw, P. S., Chen, W. and Cone, R. D. 2016. Leptin signaling
16 regulates glucose homeostasis, but not adipostasis, in the zebrafish. *Proc. Natl. Acad.*
17 *Sci.* 113: 3084–9.
- 18 44. Mittwede, P. N., Clemmer, J. S., Bergin, P. F. and Xiang, L. 2016. Obesity and
19 critical illness: insights from animal models. *Shock.* 45:349–58.
- 20 45. Moss, J. B., Koustubhan, P., Greenman, M., Parsons, M. J., Walter, I. and Moss, L. G.
21 2009. Regeneration of the pancreas in adult zebrafish. *Diabetes* 58: 1844–51.
- 22 46. Mourad, J. J. G., Guetz, H. D. and Debbabi, D. B. I. 2008. Blood pressure rise
23 following angiogenesis inhibition by bevacizumab. A crucial role for
24 microcirculation. *Ann. Oncol.* 19: 927–934.

- 1 47. Murthy, M., Kurz, T. and O'Shaughnessy, K. M. 2016. ROMK expression remains
2 unaltered in a mouse model of familial hyperkalemic hypertension caused by the
3 CUL3 Δ 403-459 mutation. *Physiol. Rep.* 4: e12850.
- 4 48. Oka, T., Nishimura, Y., Zang, L., Hirano, M., Shimada, Y., Wang, Z., Umemoto, N.,
5 Kuroyanagi, J., Nishimura, N. and Tanaka, T. 2010. Diet-induced obesity in zebrafish
6 shares common pathophysiological pathways with mammalian obesity. *BMC Physiol.*
7 10: 21.
- 8 49. Olsen, A. S. A. S, Sarras, M. P., Intine, R. V., Sarras, Jr. M. P. and Intine, R. V. 2010.
9 Limb regeneration is impaired in an adult zebrafish model of diabetes mellitus.
10 *Wound Repair Regen.* 18:532–42.
- 11 50. Osto, M. and Lutz, T. A. 2015. Translational value of animal models of obesity—
12 Focus on dogs and cats. *Eur. J. Pharmacol.* 759: 240–52.
- 13 51. Pisharath, H., Rhee, J. M., Swanson, M. A., Leach, S. D. and Parsons, M. J. 2007.
14 Targeted ablation of beta cells in the embryonic zebrafish pancreas using *E. coli*
15 nitroreductase. *Mech. Dev.* 124: 218–29.
- 16 52. Powers, J. W., Mazilu, J. K., Lin, S. and McCabe, E. R. B. 2010. The effects of
17 hyperglycemia on adrenal cortex function and steroidogenesis in the zebrafish. *Mol.*
18 *Genet. Metab.* 101: 421–2.
- 19 53. Premer, C., Lamondin, C., Mitzey, A., Speth, R. C. and Brownfield, M. S. 2013.
20 Immunohistochemical Localization of, and Angiotensin II Receptor Subtypes in the
21 Rat Adrenal, Pituitary, and Brain with a Perspective Commentary. *Int. J. Hypertens.*
22 2013: 1–22.
- 23 54. Rao, K. R., Lal, N., and Giridharan, N. V. 2014. Genetic & epigenetic approach to
24 human obesity. *Indian J. Med. Res.* 140: 589–603.

- 1 55. Rashmi Chandra, R. and Liddle, R. A. 2013. Modulation of Pancreatic Exocrine and
2 Endocrine Secretion. *Curr. Opin. Gastroenterol.* 29: 517–522.
- 3 56. Rider, S. A., Mullins, L. J., Verdon, R. F., MacRae C. A. and Mullins, J. J. 2015.
4 Renin expression in developing zebrafish is associated with angiogenesis and requires
5 the Notch pathway and endothelium. *Am. J. Physiol. Physiol.* 309: F531–9.
- 6 57. Rocha, F., Dias, J., Engrola, S., Gavaia, P., Geurden, I., Dinis, M. T. and Panserat, S.
7 2014. Glucose overload in yolk has little effect on the long-term modulation of
8 carbohydrate metabolic genes in zebrafish (*Danio rerio*). *J. Exp. Biol.* 217: 1139–49.
- 9 58. Sarras, M., Alexey, L., Olsen, A. and Intine, R. 2013. Impaired tissue regeneration
10 corresponds with altered expression of developmental genes that persists in the
11 metabolic memory state of diabetic zebrafish. *Wound Repair Regen.* 21: 320–8.
- 12 59. Sasagawa, S., Nishimura, Y., Sawada, H., Zhang, E., Okabe, S., Murakami, S.,
13 Ashikawa, Y., Yuge, M., Kawaguchi, K. and Kawase, R. 2016. Comparative
14 transcriptome analysis identifies CCDC80 as a novel gene associated with pulmonary
15 arterial hypertension. *Front. Pharmacol.* 7: 1–13.
- 16 60. Sauter, N. S., Thienel, C., Plutino, Y., Kampe, K., Dror, E., Traub, S., Timper, K.,
17 Bédard, B., Pattou, F. and Kerr-Conte, J. 2015. Angiotensin II induces interleukin-1 β -
18 mediated islet inflammation and β -cell dysfunction independently of vasoconstrictive
19 effects. *Diabetes* 64 : 1273–83.
- 20 61. Schallschmidt, T., Osthold, S., Schulte, Y., Damen, M., Stermann, T., Chadt, A. and
21 Al-Hasani, H. 2015. Identification of novel susceptibility genes for diabetes-and
22 obesity-related traits in a backcross of obese NZO with lean C3 H mice. *Diabetol.*
23 *Und. Stoffwechsel.* 10: P121.

- 1 62. Schlegel, A. and Stainier, D. Y. R. 2007. Lessons from “Lower” Organisms: What
2 Worms, Flies, and Zebrafish Can Teach Us about Human Energy Metabolism. *PLoS*
3 *Genet.* 3: e199.
- 4 63. Shibata, S. and Fujita, T. 2017. Renin Angiotensin Aldosterone System Blockers. pp
5 230-241. *In: Hypertension: A companion to Braunwald’s heart disease*, Elsevier,
6 Amsterdam, the Netherlands.
- 7 64. Shin, E., Hong, B. N. and Kang, T. H. 2012. An optimal establishment of an acute
8 hyperglycemia zebrafish model. *African J. Pharm. Pharmacol.* 6: 2922–8.
- 9 65. Siekmann, A. F. and Lawson, N. D. 2007. Notch signalling limits angiogenic cell
10 behaviour in developing zebrafish arteries. *Nature* 445: 781–784
- 11 66. Song, Y. and Cone, D. 2007. Creation of a genetic model of obesity in a teleost.
12 *FASEB J.* 21: 2042–9.
- 13 67. Tehrani, F. R. and Behboudi-Gandevani, S. 2015. Polycystic ovary syndrome. Intech
14 Open Limited, London, UK.
- 15 68. Tingaud-Sequeira A., Ouadah N., and Babin P. J. 2011. Zebrafish obesogenic test: a
16 tool for screening molecules that target adiposity. *J. Lipid. Res.* 52: 1765–72.
- 17 69. Vancampfort D., Stubbs B., Mitchell, A. J., De Hert, M., Wampers, M., Ward, P. B.,
18 Rosenbaum, S. and Correll, C. U. 2015. Risk of metabolic syndrome and its
19 components in people with schizophrenia and related psychotic disorders, bipolar
20 disorder and major depressive disorder: a systematic review and meta-analysis. *World*
21 *Psychiatry* 14: 339–47.
- 22 70. Verbrugge, F. H., Tang, W. H. W., and Mullens, W. 2015. Renin-angiotensin-
23 aldosterone system activation during decongestion in acute heart failure. *JACC: Heart*
24 *Failure* 3: 108-111.

- 1 71. Wang, N., MacKenzie, L., De Souza, A. G., Zhong, H., Goss, G. and Liang, Li.
2 2007. Proteome Profile of Cytosolic Component of Zebrafish Liver Generated by
3 LC-ESI MS/MS Combined with Trypsin Digestion and Microwave-Assisted Acid
4 Hydrolysis. *J. Proteome Res.* 6: 263–272.
- 5 72. Wang, T., Guo, R., Zhou, G., Zhou, X., Kou, Z., Sui, F., Li, C., Tang, L., and Wang,
6 Z. 2016. Traditional uses, botany, phytochemistry, pharmacology and toxicology of
7 *Panaxnotoginseng* (Burk.) FH Chen: a review. *J. Ethnopharmacol.* 188: 234–58.
- 8 73. Xie, R. F., Yang, B. R., Cheng, P. P., Wu, S., Li, Z. C., Tang, J. Y., Li, S., Tang, N.,
9 Lee, S. M. Y., and Wang, Y-H. 2015. Study on the HPLC chromatograms and pro-
10 angiogenesis activities of the flowers of *Panax notoginseng*. *J. Liq. Chromatogr.*
11 *Relat. Technol.* 38: 1286–95.
- 12 74. Yazdi, F. T., Clee, S. M., and Meyre, D. 2015. Obesity genetics in mouse and human:
13 back and forth, and back again. *PeerJ* 3: e856.
- 14 75. Zang, L., Shimada, Y., Nishimura, Y., Tanaka, T., and Nishimura, N. 2015. Repeated
15 blood collection for blood tests in adult zebrafish. *J. Vis. Exp.* e53272–e53272.
- 16 76. Zang, L., Yasuhito Shimada, Y. and Nishimura, N. 2017. Development of a Novel
17 Zebrafish Model for Type 2 Diabetes Mellitus. *Sci. Rep.* 7: 1-11.
- 18 77. Zhang, Y., Fischer, K. E., Soto, V., Liu, Y., Sosnowska, D., Richardson, A., and
19 Salmon, A. B. 2015. Obesity-induced oxidative stress, accelerated functional decline
20 with age and increased mortality in mice. *Arch. Biochem. Biophys.* 576:39–48.
- 21 78. Zhang, Y., Qin, C., Yang, L., Lu, R., Zhao, X. and Nie, G. 2018. A comparative
22 genomics study of carbohydrate/glucose metabolic genes: from fish to mammals.
23 *BMC Genet.* 19: 1-14.
- 24 79. Zheng, Y. Y., Ley, S. H. and Hu, F. B. 2018. Global aetiology and epidemiology of
25 type 2 diabetes mellitus and its complications. *Nat. Rev. Endocrinol.* 14: 88-98

- 1 80. Zodrow, J. M., Stegeman, J. J. and Tanguay, R. L. 2004. Histological analysis of
2 acute toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in zebrafish. *Aquat.*
3 *Toxicol.* 66: 25-38.

Table 1: Reported zebrafish models developed for obesity, hyperlipidemia, diabetes, and hyperglycemia

Disorders	Establishment of the model	Reference
Obesity/ hyperlipidemia	The feeding of adult zebrafish for 3 times per day with freshly hatched artemia (60 mg cysts/fish/day) for 8 weeks	48
	Feeding of adult zebrafish for 6 weeks with high fat diet (HFD) which contains 20% lard, and 80% basal diet. Each tank of 8 fish received 80 mg/day of HFD	41
	Continuous light exposure of larvae zebrafish for 24 hours. Light source used= Beams Work Power LED 200 (10.000K daylight, 200 lumen)	32
	Immersion of larvae zebrafish in 1 μ M rosiglitazone or 10 μ M T0070907 or 20 μ M phenylephrine	68
Diabetes/ hyperglycemia	Immersion of adult zebrafish in 1% water glucose solution for 30 mins, followed by immersion in water for 1 hour, and subsequently immersed in 300 mg alloxan/100 ml H ₂ O for 30 mins	64
	Single injection of adult zebrafish with 1 g/kg of streptozotocin	45
	Multiple injections of streptozotocin to the adult zebrafish: Each injection= 350 mg/kg. Week 1= 3 injections (day 1, 3, 5), week 2= 1 injection (day 12), week 3= 1 injection (day 19)	25
	Immersion of adult zebrafish in 25% glucose in Me ₂ SO	13
	Immersion of adult zebrafish in a 111 mM glucose solution for 14 days	6
	Immersion of larvae zebrafish in 10 mM metronidazole	51
	Feeding adult male AB strain zebrafish with high fat diets for 8 weeks for three times per day. The food was mixed artemia (5 mg artemia) with egg yolk powder contain 59% fat, 32% proteins, and 2% carbohydrates.	34
	Feeding the adult zebrafish with normal food which contain for 8 weeks for 6 feeds per day contain 120 mg for each feed. The food contains 11% crude fat, 51% crude protein, 2.3% crude calcium, 1.5% phosphorous, 15% ash, 3% crude fiber, and 6.5% moisture	76