

ANTIMAGIC LABELING OF DIGRAPHS

M. NALLIAH

School of Advanced Sciences, Department of Mathematics
VIT University, Vellore-632 014, INDIA.

Email: nalliahklu@gmail.com

Abstract. An antimagic labeling of a digraph D with p vertices and q arcs is a bijection f from the set of all arcs to the set of positive integers $\{1, 2, 3, \dots, q\}$ such that all the p oriented vertex weights are distinct, where an oriented vertex weight is the sum of the labels of all arcs entering that vertex minus the sum of the labels of all arcs leaving it. A digraph D is called antimagic if it admits an antimagic labeling. In this paper we investigate the existence of antimagic labelings of some few families of digraphs using hooked Skolem sequences.

Key words and Phrases: Antimagic labeling, hooked Skolem sequence, symmetric digraph.

Abstrak. Suatu pelabelan antimagic dari sebuah digraf D dengan p buah titik dan q buah busur adalah suatu bijeksi f dari himpunan semua busur ke himpunan bilangan bulat positif $\{1, 2, 3, \dots, q\}$ sedemikian sehingga semua bobot titik berarah p berbeda, dimana suatu bobot titik berarah adalah jumlahan label-label dari semua busur yang menuju titik tersebut dikurangi jumlahan label-label dari semua busur yang keluar dari titik tersebut. Sebuah digraf D dikatakan antimagic jika mempunyai pelabelan antimagic. Dalam paper ini kami menunjukkan keberadaan pelabelan antimagic dari beberapa keluarga digraf menggunakan barisan Skolem terkait.

Kata kunci: Pelabelan antimagic, barisan Skolem terkait, digraf simetris.

1. INTRODUCTION

By graph $G = (V, E)$ we mean a finite, undirected graph with neither loops nor multiple edges. The order and size of G are denoted by n and m , respectively. For graph theoretical terminology we refer to Chartrand and Lesniak [1].

2000 Mathematics Subject Classification: 05C78.

Received: 23-09-2015, revised: 04-03-2016, accepted: 29-04-2016.

A *digraph* D consists of a finite nonempty set V of objects called *vertices* and a set E of ordered pairs of distinct vertices. Each element of E is an *arc* or a directed edge. If a digraph D has the property that for each pair u, v of distinct vertices of D , at most one of (u, v) or (v, u) is an arc of D , then D is an *oriented graph*. An oriented graph can also be obtained by assigning a direction to (that is, orienting) each edge of a graph G . The digraph D is then referred to as an *orientation* of G . A digraph H is called a *subdigraph* of a digraph D if $V(H) \subseteq V(D)$ and $E(H) \subseteq E(D)$.

A digraph D is *symmetric* if whenever (u, v) is an arc of D , then (v, u) is an arc of D as well.

The underlying graph of a digraph D is obtained by removing all directions from the arcs of D and replacing any resulting pair of parallel edges by a single edge. Equivalently, the underlying graph of a digraph D is obtained by replacing each arc (u, v) or a pair $(u, v), (v, u)$ of arcs by the edge uv . For any graph G the digraph obtained by replacing every edge uv of G by a pair of symmetric arcs uv and vu is denoted by S^* and is called the *symmetric digraph* of G .

A labeling of a graph G is a mapping that assigns integers to the vertices or edges or both, subject to certain conditions. The labeling is called a vertex labeling or an edge labeling or a total labeling according to if as the domain of the mapping is V or E or $V \cup E$.

D. Hefetz, T. Mutze, and J. Schwartz [3] introduced the concept of antimagic labeling of a digraph.

An *antimagic labeling* of a digraph D with p vertices and q arcs is a bijection from the set of arcs of D to $\{1, 2, 3, \dots, q\}$ such that all p oriented vertex weights are distinct, where an oriented vertex weight is the sum of the labels of all arcs entering that vertex minus the sum of the labels of all arcs leaving it. A digraph D is called *antimagic* if it admits an antimagic labeling. The oriented vertex weight of a vertex $v \in V(D)$ is denoted by $w(v)$. An orientation D of a graph G is called an antimagic orientation if the digraph D is antimagic.

D. Hefetz, T. Mutze, and J. Schwartz [3] proved that, for every orientation of the following undirected graphs, stars S_n on $n + 1$ vertices for every $n \neq 2$, wheels W_n on $n + 1$ vertices for every $n \geq 3$ and cliques K_n on n vertices for every $n \neq 3$, there exists an antimagic labeling. They also proved that, given $G = (V, E)$ a $(2d + 1)$ -regular (not necessarily connected) undirected graph with $d \geq 0$, there exists an antimagic orientation of G . They posed the following problem and conjecture.

Question 1.1. [3] *Is every connected directed graph with at least 4 vertices antimagic?*

Conjecture 1.2. [3] *Every connected undirected graph admits an antimagic orientation.*

They observed that the answer to the Question 1.1 is “no”. Indeed, if $G = K_{1,2}$ or K_3 , then G admits an orientation that is not antimagic. Thus, not every directed graph is antimagic. Also they answer to the Conjecture 1.2 is “yes” for

few families of undirected connected graphs, but not for all connected graphs. Therefore, the Conjecture 1.2 is still open.

In this paper we discuss the existence of antimagic labelings for digraphs and we use the concept of hooked Skolem sequence.

Definition 1.3. [2, 4, 5, 6] A *hooked Skolem sequence* of order n is a sequence $S = (s_1, s_2, \dots, s_{2n+1})$ satisfying the following conditions:

- (1) For every $t \in \{1, 2, \dots, n\}$ there exist exactly two elements $s_i, s_j \in S$ such that $s_i = s_j = t$.
- (2) If $s_i = s_j = t$ with $i < j$, then $j - i = t$.
- (3) $s_{2n} = 0$.

Hooked Skolem sequence can also be written as a collection of ordered pairs $\{(a_i, b_i); 1 \leq i \leq n, b_i - a_i = i$ with $\bigcup_{i=1}^n (a_i, b_i) = \{1, 2, 3, \dots, 2n - 1, 2n + 1\}$. For example, when $n = 2$, the hooked Skolem sequence $S = (1, 1, 2, 0, 2)$ can be written as $(1, 2), (3, 5), (0, 4)$. When $n = 3$, the hooked Skolem sequence $S = (1, 1, 2, 3, 2, 0, 3)$ can be written as $(1, 2), (3, 5), (4, 7), (0, 6)$.

Theorem 1.4. [2, 4, 5, 6, 7] A *hooked Skolem sequence of order n exists if and only if $n \equiv 2$ or $3 \pmod{4}$* .

We now describe a method for the construction of a hooked Skolem sequence of order $n > 2$, where $n \equiv 2$ or $3 \pmod{4}$. We use the ordered pair notation for the hooked Skolem sequence.

When $n = 4s + 2$, $s > 0$ a hooked Skolem sequence of order n is given by

$$\left\{ \begin{array}{ll} (r, 4s - r + 2), & 1 \leq r \leq 2s, \\ (4s + r + 3, 8s - r + 4), & 1 \leq r \leq s - 1, \\ (5s + r + 2, 7s - r + 3), & 1 \leq r \leq s - 1, \\ (2s + 1, 6s + 2), (4s + 2, 6s + 3), \\ (4s + 3, 8s + 5), (7s + 3, 7s + 4). \end{array} \right.$$

When $n = 4s - 1$, $s \geq 1$, a hooked Skolem sequence of order n is given by

$$\left\{ \begin{array}{ll} (4s + r, 8s - r - 2), & 1 \leq r \leq 2s - 2, \\ (r, 4s - r - 1), & 1 \leq r \leq s - 2, \\ (s + r + 1, 3s - r), & 1 \leq r \leq s - 2, \\ (s - 1, 3s), (s, s + 1), (2s, 4s - 1), \\ (2s + 1, 6s - 1), (4s, 8s - 1). \end{array} \right.$$

When $s = 1$, skip the ordered pair $(2s, 4s - 1)$.

2. MAIN RESULTS

In this section we prove that some families of digraphs are antimagic, using the concept of hooked Skolem sequences. We use the ordered pair notation for hooked Skolem sequences.

A *non symmetric digraph* G^* is obtained from a symmetric digraph S^* by adding a new vertex v and an arc uv or vu , where $u \in V(S^*)$.

Theorem 2.1. *Let $G = \bigcup_{i=1}^k K_{1,\alpha_i}$ be the union of k stars, where $\alpha_i \geq 1, \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_k$ and $\sum_{i=1}^k \alpha_i \equiv 2$ or $3 \pmod{4}$. Then G^* is antimagic.*

PROOF. Let $V(G^*) = \bigcup_{j=1}^k \{c_j, v_1^j, v_2^j, \dots, v_{\alpha_j}^j, v\}$ and $E(G^*) = \bigcup_{j=1}^k \{c_j v_1^j, c_j v_2^j, \dots, c_j v_{\alpha_j}^j\} \cup \bigcup_{j=1}^k \{v_1^j c_j, v_2^j c_j, \dots, v_{\alpha_j}^j c_j\} \cup \{c_k v\}$. Clearly, $q = |E(G)| = \sum_{i=1}^k \alpha_i \equiv 2$ or $3 \pmod{4}$ and $|E(G^*)| = 2q + 1, q \equiv 2$ or $3 \pmod{4}$. Hence, there exists a hooked Skolem sequence S of order q (See Theorem 1.4). Let $S = \{(a_1, b_1), (a_2, b_2), \dots, (a_q, b_q), (0, b_{q+1})\}$, where $1 \leq a_i \leq 2q + 1, 1 \leq b_i \leq 2q + 1$ and $b_i - a_i = i$, where $1 \leq i \leq q$. Now define $f^* : E(G^*) \rightarrow \{1, 2, \dots, 2q + 1\}$ by

$$\begin{aligned} f^*(c_j v_i^j) &= a_i, \text{ for } 1 \leq j \leq k, 1 \leq i \leq \alpha_j, \\ f^*(v_i^j c_j) &= b_i, \text{ for } 1 \leq j \leq k, 1 \leq i \leq \alpha_j, \\ f^*(c_k v) &= 2q. \end{aligned}$$

Then

$$\begin{aligned} w(c_1) &= \frac{\alpha_1(\alpha_1 + 1)}{2}, \\ w(c_j) &= (\alpha_1 + \alpha_2 + \dots + \alpha_{j-1})\alpha_j + \frac{\alpha_j(\alpha_j + 1)}{2}, \text{ if } 2 \leq j \leq k - 1 \text{ and } 1 \leq i \leq \alpha_j, \\ w(c_k) &= (\alpha_1 + \alpha_2 + \dots + \alpha_{k-1})\alpha_k + \frac{\alpha_k(\alpha_k + 1)}{2} - 2q, \\ w(v_i^1) &= -i, 1 \leq i \leq \alpha_1. \\ \text{Let } x &= \sum_{j=2}^k \alpha_{j-1}. \\ w(v_{x+i}^j) &= - \left[\sum_{j=2}^k \alpha_{j-1} + i \right], 2 \leq j \leq k \text{ and } 1 \leq i \leq \alpha_j, \\ w(v) &= 2q. \end{aligned}$$

Clearly, all the oriented vertex weights are distinct and hence, f^* is an antimagic labeling of G^* .

Example 2.2. Let $G = K_{1,4} \cup K_{1,6}$, so that $\alpha_1 = 4$ and $\alpha_2 = 6$. Clearly, $\alpha_1 + \alpha_2 \equiv 2 \pmod{4}$. A hooked Skolem sequence of order 10 (in ordered pair notation) is given by $S = \{(17, 18), (4, 6), (13, 16), (3, 7), (10, 15), (2, 8), (12, 19), (1, 9), (5, 14), (11, 21), (0, 20)\}$. The corresponding antimagic labeling of the symmetric digraph G^* is given in Figure 1.




FIGURE 1. An antimagic labeling of $(K_{1,4} \cup K_{1,6})^*$.

Theorem 2.3. Let $G = K_{r,s}$ be the complete bipartite graph and $rs \equiv 2$ or $3 \pmod{4}$. Then G^* is antimagic.

PROOF. Let $X(G^*) = \{u_1, u_2, \dots, u_s\}$ and $Y(G^*) = \{v_1, v_2, \dots, v_s, v\}$ be the bipartition of G^* . Then $E(G^*) = \bigcup_{i=1}^r \{u_i v_1, u_i v_2, \dots, u_i v_s\} \cup \bigcup_{i=1}^r \{v_1 u_i, v_2 u_i, \dots, v_s u_i\} \cup \{u_r v\}$. Clearly, $q = |E(G)| = rs \equiv 2$ or $3 \pmod{4}$ and $|E(G^*)| = 2q + 1$. Hence, there exists a hooked Skolem sequence S of order q and let $S = \{(a_1, b_1), (a_2, b_2), \dots, (a_q, b_q), (0, b_{q+1})\}$, where $1 \leq a_i \leq 2q+1, 1 \leq b_i \leq 2q+1$ and $b_i - a_i = i$.

Now we define $f^* : E(G^*) \rightarrow \{1, 2, \dots, 2q + 1\}$ by

$$\begin{aligned} f^*(u_i v_j) &= a_{(i-1)s+j}, \\ f^*(v_j u_i) &= b_{(i-1)s+j}, \\ f^*(u_s v) &= rs. \end{aligned}$$

Since $b_{(i-1)s+j} - a_{(i-1)s+j} = (i-1)s + j$, $1 \leq i \leq r$ and $1 \leq j \leq s$, it follows that $w(u_i) = (i-1)s^2 + \frac{s(s+1)}{2}$, $1 \leq i \leq r-1$, $w(u_r) = (r-1)s^2 + \frac{s(s+1)}{2} - 2rs$, $w(v) = 2rs$ and $w(v_j) = -\left[rj + \frac{rs(r-1)}{2}\right]$. Clearly, all the oriented vertex weights are distinct and hence, f^* is an antimagic labeling of G^* .

Example 2.4. Let $G = K_{2,3}$. Clearly, $q = 6$. A hooked Skolem sequence of order 6 (in ordered pair notation) is given by $S = \{(10, 11), (2, 4), (6, 9), (1, 5), (3, 8), (7, 13), (0, 12)\}$. The corresponding antimagic labeling of the digraph G^* is given in Figure 2.




FIGURE 2. An antimagic labeling of $K_{2,3}^*$.

Theorem 2.5. Let $G = C_p$ be the cycle of order p , where $p \equiv 2$ or $3 \pmod{4}$. Then G^* is antimagic.

PROOF. Let $V(G^*) = \{u_1, u_2, \dots, u_p, v\}$ and $E(G^*) = \{u_1 u_2, u_2 u_3, \dots, u_p u_1\} \cup \{u_2 u_1, u_3 u_2, \dots, u_1 u_p\} \cup \{u_p v\}$. Since $q = p \equiv 2$ or $3 \pmod{4}$, there exists a hooked Skolem sequence S of order q . Let $S = \{(a_1, b_1), \dots, (a_q, b_q), (0, b_{q+1})\}$,

where $1 \leq a_i \leq 2q + 1, 1 \leq b_i \leq 2q + 1$ and $b_i - a_i = i, 1 \leq i \leq q$. We consider two cases.

Case 1. $q \equiv 2(\text{mod } 4)$.

Define $f^* : E(G^*) \rightarrow \{1, 2, \dots, 2q + 1\}$ as follows:

$$\begin{aligned} f(u_i u_{i+1}) &= \begin{cases} b_{2i-1}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p}{2} \\ a_{2i-1}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} \end{cases} \\ f(u_{i+1} u_i) &= \begin{cases} a_{2i-1}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p}{2} \\ b_{2i-1}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} \end{cases} \\ f(u_{\frac{p}{2}+i} u_{\frac{p}{2}+i+1}) &= \begin{cases} a_{p+2-2i}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p}{2} - 1 \\ b_{p+2-2i}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} - 1 \end{cases} \\ f(u_{\frac{p}{2}+i+1} u_{\frac{p}{2}+i}) &= \begin{cases} b_{p+2-2i}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p}{2} - 1 \\ a_{p+2-2i}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} - 1 \end{cases} \\ f(u_p u_1) &= a_2, \quad f(u_1 u_p) = b_2, \quad f(u_p v) = 2p. \end{aligned}$$

Then

$$\begin{aligned} w(u_{i+1}) &= \begin{cases} 4i, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p}{2} - 1 \\ -4i, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} - 1 \end{cases} \\ w(u_{\frac{p}{2}+i}) &= \begin{cases} 2p - 1, & \text{if } i = 1 \\ 2p + 6 - 4i, & \text{if } i \text{ is odd, } 3 \leq i \leq \frac{p}{2} - 1 \\ -(2p + 6 - 4i), & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p}{2} - 1 \end{cases} \\ w(u_1) &= -3, \quad w(u_p) = -2p + 6. \end{aligned}$$

Clearly, all the oriented vertex weights are distinct.

Case 2. $q \equiv 3(\text{mod } 4)$.

$$\begin{aligned} f(u_i u_{i+1}) &= \begin{cases} b_{2i-1}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p+1}{2} \\ a_{2i-1}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} \end{cases} \\ f(u_{i+1} u_i) &= \begin{cases} a_{2i-1}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p+1}{2} \\ b_{2i-1}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} \end{cases} \\ f(u_{\frac{p+1}{2}+i} u_{\frac{p+1}{2}+i+1}) &= \begin{cases} b_{p+1-2i}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p+1}{2} - 2 \\ a_{p+1-2i}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} - 2 \end{cases} \\ f(u_{\frac{p+1}{2}+i+1} u_{\frac{p+1}{2}+i}) &= \begin{cases} a_{p+1-2i}, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p+1}{2} - 2 \\ b_{p+1-2i}, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} - 2 \end{cases} \\ f(u_p u_1) &= a_2, \quad f(u_1 u_p) = b_2, \quad f(u_p v) = 2p. \end{aligned}$$

Then

$$w(u_{i+1}) = \begin{cases} 4i, & \text{if } i \text{ is odd, } 1 \leq i \leq \frac{p+1}{2} - 2 \\ -4i, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} - 2 \end{cases}$$

$$w(u_{\frac{p+1}{2}+i}) = \begin{cases} -(2p-1), & \text{if } i = 1 \\ -(2p+4-4i), & \text{if } i \text{ is odd, } 3 \leq i \leq \frac{p+1}{2} - 2 \\ 2p+4-4i, & \text{if } i \text{ is even, } 2 \leq i \leq \frac{p+1}{2} - 2 \end{cases}$$

$$w(u_1) = -3, \quad w(u_p) = -(2p+6).$$

Clearly, all the oriented vertex weights are distinct and hence, f^* is an antimagic labeling of G^* .

Example 2.6. Let $G = C_{10}$ and C_{11} . A hooked Skolem sequence of order 10 (in ordered pair notation) is given by $S = \{(17, 18), (4, 6), (13, 16), (3, 7), (10, 15), (2, 8), (12, 19), (1, 9), (5, 14), (11, 21), (0, 20)\}$, and a hooked Skolem sequence of order 11 (in order pair notation) is given by $S = \{(3, 4), (16, 18), (5, 8), (15, 19), (6, 11), (14, 20), (2, 9), (13, 21), (1, 10), (7, 17), (12, 23), (0, 22)\}$. The corresponding antimagic labeling of the digraph G^* is given in Figure 3 and 4, respectively.





FIGURE 3. An antimagic labeling of C_{10}^* .

FIGURE 4. An antimagic labeling of C_{11}^* .

3. CONCLUSION AND SCOPE

In this paper we have discussed the existence of antimagic labelings of digraphs. In particular, if G^* is a digraph associated with an undirected graph G , we have used the concept of hooked Skolem sequences to prove the existence of antimagic labelings of G^* for several classes of graphs. This proof technique can be used to prove the existence of antimagic labelings of G^* for other families of graphs.

Acknowledgement. The author would like to thank the referees for their valuable comments.

REFERENCES

- [1] Chartrand, G., and Lesniak, L., *Graphs & Digraphs*, Fourth Edition, Chapman & Hall/CRC, 2005.
- [2] Colbourn, C.J., and Dinitz, J.H., *Handbook of combinatorial designs*, second edition, Chapman & Hall/ CRC, 2006.
- [3] Hefetz, D., Mutze, T., and Schwartz, J., "On Antimagic Directed Graphs", *J. Graph Theory*, **64** (2010), 219-232.
- [4] Linek, V., and Jiang, Z., Hooked k-extended Skolem sequences, *Discrete Math.*, **196**(1-3) (1999), 229-238.
- [5] O'Keefe, E.S., "Verification of a conjecture of Th. Skolem", *Math. Scand.*, **9** (1961), 80-82.
- [6] Shalaby, N., "The existence of near-Skolem sequences and hooked near-Skolem sequences", *Discrete Math.*, **135** (1994), 303-319.
- [7] Shalaby, N., *Skolem sequences: Generalizations and Applications*, Ph. D. thesis, McMaster University, Canada, 1991.

