



Draft Genome Sequence of Terrestrial *Streptomyces* sp. Strain VITNK9, Isolated from Vellore, Tamil Nadu, India, Exhibiting Antagonistic Activity against Fish Pathogens

Subhasish Saha,^a Nabila Mohammad Ishaque,^b Ilia Burgsdorf,^c Markéta Macho,^a Daniela Ewe,^a Laura Steindler,^c Pavel Hrouzek,^a Krishnan Kannabiran,^b  Kumar Saurav^a

^aLaboratory of Algal Biotechnology-Centre Algotech, Institute of Microbiology of the Czech Academy of Sciences, Treboň, Czech Republic

^bDepartment of Biomedical Sciences, School of Biosciences and Technology, VIT University, Vellore, India

^cDepartment of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa, Israel

Subhasish Saha, Nabila Mohammad Ishaque, and Ilia Burgsdorf contributed equally to this work. Author order was determined on the basis of seniority.

ABSTRACT We report the draft genome sequence of *Streptomyces* sp. strain VITNK9, isolated from a soil sample collected in Vellore District (12.9165°N, 79.1325°E), Tamil Nadu, India, with an assembly size of 7,920,076 bp and 72.7% GC content.

Streptomyces spp. have been considered one of the most prolific sources of pharmacologically active compounds for decades, contributing approximately 60% of antibiotics which are in clinical use today. To discover natural products possessing antagonistic activity against fish pathogens, we isolated a new actinomycete strain, *Streptomyces* sp. strain VITNK9, from a terrestrial soil sample from Vellore, Tamil Nadu, India, using actinomycete isolation agar (AIA) plates (1, 2). The strain had a smooth colony surface, an earthy odor, and a spiral spore formation, including aerial mycelia and substrate hyphae penetrating through agar, morphologies typical of *Streptomyces*. The crude extract of the strain exhibited antimicrobial activity against several fish pathogens, implying that this strain can be an important resource of bioactive metabolites (1). Further, to obtain the genomic potential of VITNK9 to synthesize bioactive compounds, its genome was sequenced.

For DNA isolation, *Streptomyces* sp. strain VITNK9 was incubated on ISP1 broth at 30°C and 250 rpm for 4 days. Genomic DNA was isolated using the High Pure PCR template preparation kit (Roche) following the manufacturer's protocol. DNA libraries were prepared using the TruSeq DNA PCR-free kit. DNA shotgun libraries were sequenced on an Illumina platform (101-bp paired-end reads) at Macrogen (South Korea), yielding a total of 6.05 Gbp of sequence with 59,886,702 reads. Sequences were trimmed using Trimmomatic version 0.36 (minlen 100, sliding window 4:20, ILLUMINACLIP:TruSeq3PE-2.fa:2:30:8) (3), and read quality was assessed using FastQC version 0.11.5 (4) prior to *de novo* assembly using SPAdes version 3.13.0 (parameters: `-k 21,33,55,77,91 -careful -only-assembler -cov-cutoff auto`) (5). Scaffolds of ≥ 2 kb were used for further analysis. Completeness and contamination were estimated with CheckM version 1.0.7 (6) based on 460 markers and using lineage workflow. For the phylogenetic analysis, sequences of the closest described type strains were obtained from EzTaxon (<https://www.ezbiocloud.net/>) (7). Open reading frames (ORFs) for the VITNK9 genome were identified and annotated using Rapid Annotations using Subsystems Technology (RAST) (8) with the RASTtk algorithm (<https://rast.nmpdr.org/>) (9).

Genome assembly of VITNK9 yielded 18 scaffolds with an N_{50} value of 1,271,901 bp and with the largest contig being 1,996,233 bp. The genome consists of 7,919,889 bp without ambiguous nucleotides (N) with a GC content of 72.7%, 100% estimated

Citation Saha S, Ishaque NM, Burgsdorf I, Macho M, Ewe D, Steindler L, Hrouzek P, Kannabiran K, Saurav K. 2021. Draft genome sequence of terrestrial *Streptomyces* sp. strain VITNK9, isolated from Vellore, Tamil Nadu, India, exhibiting antagonistic activity against fish pathogens. *Microbiol Resour Announc* 10:e01282-20. <https://doi.org/10.1128/MRA.01282-20>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2021 Saha et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kumar Saurav, sauravverma17@gmail.com.

Received 8 November 2020

Accepted 29 November 2020

Published 7 January 2021

completeness, 0.06% contamination, and 0% heterogeneity. The VITNK9 genome contains 7,148 predicted and 73 RNA genes (68 tRNA genes). Based on EzTaxon analyses of the extracted 16S rRNA gene (876 bp) from VITNK9, the strain showed 84.11% similarity to *Streptomyces rochei* strain NRRL B-2410^T.

The closest neighbors of strain VITNK9 were *Streptomyces coelicolor* A3(2) (score, 500), *Streptomyces avermitilis* MA-4680 (score, 436), *Streptomyces scabiei* 87.22 (score, 428), *Streptomyces griseus* subsp. *griseus* NBRC 13350 (score, 377), and *Saccharopolyspora erythraea* NRRL 2338 (score, 213) detected by RAST annotation (10). The MICs of the ethyl acetate (EA) crude extract of VITNK9 were found to be moderate against the following fish pathogens: *Aeromonas hydrophila* and *Edwardsiella tarda* (0.03 mg ml⁻¹), *Vibrio anguillarum* and *Vibrio harveyi* (0.06 mg ml⁻¹, and *Aeromonas caviae* (0.125 mg ml⁻¹) (1). Furthermore, an antiSMASH version 5.1.2 analysis (11) of the VITNK9 genome revealed 30 putative biosynthetic gene clusters (BGCs). According to the prediction, 12 gene clusters possess >75% similarity to BGCs, 6 gene clusters possess 50 to 75% similarity, 8 gene clusters showed <50% similarity with known BGCs, and 4 gene clusters were detected as completely unknown. The notable BGCs found using antiSMASH prediction were in the ribosomally synthesized and posttranslationally modified peptide (RiPP) class, (lantipeptide SapB [12]), the terpene class (geosmin [13], isorenieratene [14], hopene [15], albaflavenone [16], siderophore desferrioxamine B/E [17], and ectoine [18]), nonribosomal peptide synthetase (coelibactin and coelichelin [15]), and several polyketide synthase (PKS) classes. Default parameters were used for all the tools used except where otherwise noted.

Overall, *Streptomyces* sp. strain VITNK9 possesses a high potential for diverse and bioactive secondary metabolites and can further be considered an important strain to study for bioactive compounds and their biosynthesis in the future.

Data availability. The genome of the *Streptomyces* sp. strain VITNK9 has been deposited at GenBank under accession number no. [JACWAA000000000](https://ncbi.nlm.nih.gov/assembly/JACWAA000000000/), assembly accession no. [ASM1485402v1](https://ncbi.nlm.nih.gov/assembly/ASM1485402v1/), BioProject no. [PRJNA661553](https://ncbi.nlm.nih.gov/bioproject/PRJNA661553/), and BioSample no. [SAMN16063085](https://ncbi.nlm.nih.gov/biosample/SAMN16063085/). The raw reads are available under accession no. [SRR12997718](https://ncbi.nlm.nih.gov/sra/SRR12997718/) in the Sequence Read Archive (SRA).

ACKNOWLEDGMENTS

We sincerely thank the management of the Vellore Institute of Technology for providing the necessary facilities to carry out this research work.

This study was also supported by the Ministry of Education, Youth and Sports of the Czech Republic MSCA IF II project (CZ.02.2.69/0.0/0.0/18_070/0010493), a Technology Agency of the Czech Republic (TAČR)–National Centre of Competence grant (no. TN01000048/03), and Czech Science Foundation (GAČR) project no. 19-17868Y.

REFERENCES

- Ishaque Nabila M, Kannabiran K. 2018. Antagonistic activity of terrestrial *Streptomyces* sp. VITNK9 against Gram negative bacterial pathogens affecting the fish and shellfish in aquaculture. *Rev Biol Mar Oceanogr* 53:171–183. <https://doi.org/10.22370/rbmo.2018.53.2.1291>.
- Ishaque NM, Burgsdorf I, Limlingan Malit JJ, Saha S, Teta R, Ewe D, Kannabiran K, Hrouzek P, Steindler L, Costantino V, Saurav K. 2020. Isolation, genomic and metabolomic characterization of *Streptomyces tendae* VITAKN with quorum sensing inhibitory activity from southern India. *Microorganisms* 8:121. <https://doi.org/10.3390/microorganisms8010121>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Wingett SW, Andrews S. 2018. FastQ Screen: a tool for multi-genome mapping and quality control. *F1000Res* 7:1338. <https://doi.org/10.12688/f1000research.15931.2>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Kim OS, Cho YJ, Lee K, Yoon SH, Kim M, Na H, Park SC, Jeon YS, Lee JH, Yi H, Won S, Chun J. 2012. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol* 62:716–721. <https://doi.org/10.1099/ijs.0.038075-0>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, III, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation

- pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
10. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
 11. Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T. 2019. antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res* 47:W81–W87. <https://doi.org/10.1093/nar/gkz310>.
 12. Kodani S, Hudson ME, Durrant MC, Buttner MJ, Nodwell JR, Willey JM. 2004. The SapB morphogen is a lantibiotic-like peptide derived from the product of the developmental gene *ramS* in *Streptomyces coelicolor*. *Proc Natl Acad Sci U S A* 101:11448–11453. <https://doi.org/10.1073/pnas.0404220101>.
 13. Jiang J, He X, Cane DE. 2007. Biosynthesis of the earthy odorant geosmin by a bifunctional *Streptomyces coelicolor* enzyme. *Nat Chem Biol* 3:711–715. <https://doi.org/10.1038/nchembio.2007.29>.
 14. Iftime D, Kulik A, Hartner T, Rohrer S, Niedermeyer TH, Stegmann E, Weber T, Wohlleben W. 2016. Identification and activation of novel biosynthetic gene clusters by genome mining in the kirromycin producer *Streptomyces collinus* Tu 365. *J Ind Microbiol Biotechnol* 43:277–291. <https://doi.org/10.1007/s10295-015-1685-7>.
 15. Bentley SD, Chater KF, Cerdeno-Tarraga AM, Challis GL, Thomson NR, James KD, Harris DE, Quail MA, Kieser H, Harper D, Bateman A, Brown S, Chandra G, Chen CW, Collins M, Cronin A, Fraser A, Goble A, Hidalgo J, Hornsby T, Howarth S, Huang CH, Kieser T, Larke L, Murphy L, Oliver K, O’Neil S, Rabinowitsch E, Rajandream MA, Rutherford K, Rutter S, Seeger K, Saunders D, Sharp S, Squares R, Squares S, Taylor K, Warren T, Wietzorrek A, Woodward J, Barrell BG, Parkhill J, Hopwood DA. 2002. Complete genome sequence of the model actinomycete *Streptomyces coelicolor* A3(2). *Nature* 417:141–147. <https://doi.org/10.1038/417141a>.
 16. Zhao B, Lin X, Lei L, Lamb DC, Kelly SL, Waterman MR, Cane DE. 2008. Biosynthesis of the sesquiterpene antibiotic albaflavenone in *Streptomyces coelicolor* A3(2). *J Biol Chem* 283:8183–8189. <https://doi.org/10.1074/jbc.M710421200>.
 17. Barona-Gomez F, Wong U, Giannakopoulos AE, Derrick PJ, Challis GL. 2004. Identification of a cluster of genes that directs desferrioxamine biosynthesis in *Streptomyces coelicolor* M145. *J Am Chem Soc* 126:16282–16283. <https://doi.org/10.1021/ja045774k>.
 18. Prabhu J, Schauwecker F, Grammel N, Keller U, Bernhard M. 2004. Functional expression of the ectoine hydroxylase gene (*thpD*) from *Streptomyces chrysomallus* in *Halomonas elongata*. *Appl Environ Microbiol* 70:3130–3132. <https://doi.org/10.1128/aem.70.5.3130-3132.2004>.