



# Genome Sequence of a Moderately Halophilic *Bacillus cereus* Strain, TS2, Isolated from Saltern Sediments

Manoharan Shankar,<sup>a\*</sup> Anbazhagan Mageswari,<sup>a\*</sup> Chandrasekaran Suganthi,<sup>a\*</sup> Paramasamy Gunasekaran,<sup>a\*</sup> Kodiveri M. Gothandam,<sup>a</sup> Sivashanmugam Karthikeyan<sup>a</sup>

<sup>a</sup>School of Bio-Sciences and Technology, Vellore Institute of Technology, Vellore, Tamil Nadu, India

**ABSTRACT** We report the 5.3-Mbp genome sequence of *Bacillus cereus* strain TS2, which was isolated from the sediments of a solar saltern in southern India. Genome analysis of *B. cereus* TS2, a salt-resistant strain, will improve our understanding of how *B. cereus*, a food pathogen, responds to hyperosmotic stress.

*Bacillus cereus* is one of the most frequent causes of food contamination and consequent food poisoning. Owing to its ability to produce enterotoxins, *B. cereus* can cause food poisoning, which manifests as either an emetic or a diarrheal syndrome (1). This is achieved by means of evading the host's defenses by production of various virulence factors that also allow the pathogen to damage the host's intestinal epithelial cells, resulting in diarrhea (2). Since some *B. cereus* strains and spores are capable of surviving normal food preservation methods, including refrigeration (3), use of heat, alkali, or acid (4), and salt treatment (5), it is essential to study the mechanisms by which resistance to these treatments is achieved.

*B. cereus* TS2 was isolated from the sediments of a solar saltern in Tuticorin, Tamil Nadu, India, and was capable of growing in 8% NaCl. The sequence of the *B. cereus* TS2 genome was determined on a HiSeq 2000 sequencing system per the standard Illumina protocol. Paired-end sequence reads (100 bp) were subjected to adapter removal and quality control using the SeqQC v2.2 suite (Genotypic, Bangalore, India). SPAdes v3.1.0 (6) was then used for *de novo* assembly of the high-quality reads into 173 contigs ( $N_{50}$ , 537,946 bp). Using the scaffolding tool SSPACE v2.0 (7), the 173 contigs were organized into 121 scaffolds ( $N_{50}$ , 1,475,409 bp). Finally, using the *B. cereus* Q1 genome (NCBI accession no. NC\_011969) as a reference, gap closure was attempted using GapCloser v1.12 from the SOAPdenovo2 assembly package (8), resulting in 24 gap-joined scaffolds totaling 5,308,102 bp in length with a GC content of 35.3%.

Gene prediction using the Rapid Annotation using Subsystems Technology (RAST) server (9) led to the identification of 5,493 coding sequences. We identified homologs of the GroELS and DnaK heat shock response chaperonin families in *B. cereus* TS2, among other heat shock response-inducible genes. Strain TS2 also harbored the cold shock response proteins CspA and CspD, which allow *Bacillus subtilis* to survive at low food preservation temperatures (10). Homologs of genes previously implicated in Gram-positive salt stress response (11), including the K<sup>+</sup> uptake protein TrkH, osmo-protectant uptake (Opu) class choline/glycine betaine uptake systems, and the L-proline transporter ProP, were identified on the TS2 chromosome. We also identified various virulence factors, including the toxins (enterotoxins A, B, and C, a predicted hemolysin, and cytotoxin K) (12) usually encoded by pathogenic *B. cereus* strains. Additionally, hydrolytic enzymes (phospholipase C, sphingomyelinase C, and collagenase), which act as virulence factors *in vivo*, were detected in *B. cereus* TS2. Homologs of acid resistance determinants (F1F0-ATPase, cyclopropane fatty acyl phospholipid synthase, and amino acid decarboxylases) which enable *B. cereus* to transit through the hostile gastric environment (13) and reach the intestines were also detected. To our

Received 19 June 2018 Accepted 10 July 2018 Published 16 August 2018

**Citation** Shankar M, Mageswari A, Suganthi C, Gunasekaran P, Gothandam KM, Karthikeyan S. 2018. Genome sequence of a moderately halophilic *Bacillus cereus* strain, TS2, isolated from saltern sediments. Microbiol Resour Announc 7:e00873-18. <https://doi.org/10.1128/MRA.00873-18>.

**Editor** Frank J. Stewart, Georgia Institute of Technology

**Copyright** © 2018 Shankar et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sivashanmugam Karthikeyan, [asstdirector.pat@vit.ac.in](mailto:asstdirector.pat@vit.ac.in).

\* Present address: Manoharan Shankar, Department of Bioscience & Bioengineering, Indian Institute of Technology, Jodhpur, Rajasthan, India; Anbazhagan Mageswari, Department of Biotechnology, M.M.E.S Women's Arts and Science College Vellore, Tamil Nadu, India; Chandrasekaran Suganthi, Department of Biotechnology, D.K.M College for Women, Vellore, Tamil Nadu, India; Paramasamy Gunasekaran, Vellore Institute of Technology, Bhopal, Madhya Pradesh, India.

surprise, we also detected resistance mechanisms against zinc, arsenic, cobalt, cadmium, copper, chromium, bile, and antibiotics (streptothricin, tetracycline, aminoglycosides, vancomycin, fosfomycin, fluoroquinolones, and  $\beta$ -lactam antibiotics) encoded on the TS2 genome. We are currently testing and evaluating *B. cereus* TS2 as a potential food contaminant and a pathogen.

**Data availability.** This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [PIYQ00000000](https://doi.org/10.1111/j.1574-6976.2018.00112.x). The version described in this paper is the first version, PIYQ01000000.

## ACKNOWLEDGMENTS

We thank the Vellore Institute of Technology, Vellore, India, for access to research facilities and for providing research support. Technical and bioinformatics support for bacterial genome sequencing from Genotypic Technology, Bangalore, India, is acknowledged.

## REFERENCES

1. Sergeev N, Distler M, Vargas M, Chizhikov V, Herold KE, Rasooly A. 2006. Microarray analysis of *Bacillus cereus* group virulence factors. *J Microbiol Methods* 65:488–502. <https://doi.org/10.1016/j.mimet.2005.09.013>.
2. Stenfors Arnesen LP, Fagerlund A, Granum PE. 2008. From soil to gut: *Bacillus cereus* and its food poisoning toxins. *FEMS Microbiol Rev* 32: 579–606. <https://doi.org/10.1111/j.1574-6976.2008.00112.x>.
3. Guinebretiere M-H, Thompson FL, Sorokin A, Normand P, Dawyndt P, Ehling-Schulz M, Svensson B, Sanchis V, Nguyen-The C, Heyndrickx M, De Vos P. 2008. Ecological diversification in the *Bacillus cereus* group. *Environ Microbiol* 10:851–865. <https://doi.org/10.1111/j.1462-2920.2007.01495.x>.
4. Shaheen R, Svensson B, Andersson MA, Christiansson A, Salkinoja-Salonen M. 2010. Persistence strategies of *Bacillus cereus* spores isolated from dairy silo tanks. *Food Microbiol* 27:347–355. <https://doi.org/10.1016/j.fm.2009.11.004>.
5. Browne N, Dowds BCA. 2001. Heat and salt stress in the food pathogen *Bacillus cereus*. *J Appl Microbiol* 91:1085–1094. <https://doi.org/10.1046/j.1365-2672.2001.01478.x>.
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
7. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
8. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, Tang J, Wu G, Zhang H, Shi Y, Liu Y, Yu C, Wang B, Lu Y, Han C, Cheung DW, Yiu S-M, Peng S, Xiaoqian Z, Liu G, Liao X, Li Y, Yang H, Wang J, Lam T-W, Wang J. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read *de novo* assembler. *Gigascience* 1:18. <https://doi.org/10.1186/2047-217X-1-18>.
9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
10. Graumann P, Wendrich TM, Weber MHW, Schröder K, Marahiel MA. 1997. A family of cold shock proteins in *Bacillus subtilis* is essential for cellular growth and for efficient protein synthesis at optimal and low temperatures. *Mol Microbiol* 25:741–756. <https://doi.org/10.1046/j.1365-2958.1997.5121878.x>.
11. Sleator RD, Hill C. 2002. Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence. *FEMS Microbiol Rev* 26:49–71. <https://doi.org/10.1111/j.1574-6976.2002.tb00598.x>.
12. Clair G, Roussi S, Armengaud J, Duport C. 2010. Expanding the known repertoire of virulence factors produced by *Bacillus cereus* through early secretome profiling in three redox conditions. *Mol Cell Proteomics* 9:1486–1498. <https://doi.org/10.1074/mcp.M000027-MCP201>.
13. Duport C, Jobin M, Schmitt P. 2016. Adaptation in *Bacillus cereus*: from stress to disease. *Front Microbiol* 7:1550. <https://doi.org/10.3389/fmicb.2016.01550>.