

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/277667730>

African Journal of Biotechnology Heterosis expression, interrelationship, direct and indirect effects of component characters on yield in intervarietal crosses of eggplant

Article in AFRICAN JOURNAL OF BIOTECHNOLOGY · December 2013

DOI: 10.5897/AJB2013.12438

CITATIONS

5

READS

92

5 authors, including:



Ramesh Kumar Selvan

vanavarayar institute of agriculture

26 PUBLICATIONS 188 CITATIONS

SEE PROFILE



Arumugam Thangaiah

Tamil Nadu Agricultural University

43 PUBLICATIONS 345 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Bioecology and Management of gall midge in bitter gourd [View project](#)

Full Length Research Paper

Heterosis expression, interrelationship, direct and indirect effects of component characters on yield in intervarietal crosses of eggplant

S. Ramesh Kumar^{1*}, T. Arumugam², C. R. Anandakumar³, S. Balakrishnan² and D. S. Rajavel⁴

¹Department of Crop Improvement, Vanavarayar Institute of Agriculture, Manakkadavu, Pollachi-642103, TNAU, Tamil Nadu, India.

²Department of Horticulture, Agricultural College and Research Institute, TNAU, Madurai-625104, Tamil Nadu, India.

³Director, Centre for Plant Breeding and Genetics, TNAU, Coimbatore-641003, Tamil Nadu, India.

⁴Department of Crop Protection, Agricultural College and Research Institute, Killikulam, TNAU, Tuticorin- 628 252, Tamil Nadu, India.

Accepted 26 August, 2013

Heterosis has been amply exploited in developing hybrids in brinjal. In India, only 17.8% area of brinjal cultivation is under hybrid seed due to lack of appropriate hybrids for specific area and purpose. In the present investigation, 14 parents were selected on the basis of divergence and mated in line (L) × tester (T) design hybrids and parents were raised to measure heterosis for different yield attributes. 40 hybrids resulting from a L × T mating design comprised of 10 lines (female) and 4 testers (males) were studied to determine the magnitude of heterosis and genotypic correlation between yield and yield characters. Expression of superiority over the commercial check occurred in 7 crosses, which ranged from -56.36 (L₃ × T₄) to 34.07% (L₇ × T₂). The hybrid (L₇ × T₂) had good heterosis values for growth and yield and is recommended as the most promising combination for developing high yielding hybrid eggplant varieties. Most crosses involving T₂ as tester parent had significant positive heterosis over the mid-parent and standard variety. There were strong correlations between fruit yield and numbers of branches per plant, average fruit weight and number of fruit per plant. Path analysis indicated that plant height, number of branches per plant, fruit length, fruit pedicel length, number of fruit per plant, average fruit weight and little leaf incidence had direct and positive effects on yield per plant, but negative and direct effects occurred for days to first flowering, fruit circumference, calyx length, shoot borer infestation, fruit borer infestation, ascorbic acid content and total phenol content. Simple selection would be effective for improvement of fruit yield.

Key words: *Solanum melongena*, association analysis, hybrid vigour.

INTRODUCTION

Brinjal (*Solanum melongena* L.), also known as eggplant and an important vegetable crop of India. It is widely cultivated in both temperate and tropical regions of the

globe mainly for its immature fruits. In Tamilnadu (India), the productivity of brinjal is less as compared to national average, owing to use of low yielding cultivars grown for

*Corresponding author. E-mail: rameshamar06@gmail.com or ramesh_amar06@yahoo.co.in.

local preferences and their susceptibility to pests and diseases (Nalini, 2007). The present production is not proportionate with the demand. So, brinjal deserves a deep deliberation for improvement. It should be highly pragmatic by the fact that, India being the centre of origin and diversity of brinjal, it should pave the way for bringing about a kind of plant type, which could enhance its quality and productivity without sacrificing the consumer needs (Shafeeq et al., 2007). Quality and productivity of eggplant can be improved through heterosis breeding (Kakikazi, 1931). The estimation of heterosis for yield and its component characters would be useful to judge the best hybrid combination for exploitation of superior hybrids. Genotypic correlations exist between yield and yield attributing characters are important in breeding. Yield is the end product of many correlated characters. Selection for yield would be more effective when it is based on component characters which are positively correlated. When more number of variables is considered in correlation, the association becomes more complex. Use of path coefficient analysis makes clear direct and indirect associations and identifies the most reliable yield contributing characters (Daliya and Wilson, 2002). The study was under taken to study the extent of heterosis in different crosses and their utilization in future crop improvement.

MATERIALS AND METHODS

The experiment was conducted during June to October (*kharif*) 2010 to 2011 at College Orchard, Agricultural College and Research Institute, Madurai, India, situated at 9°5' N latitude and 78°5' E longitudes at 147 m above MSL. Ten lines were crossed with 4 testers through Line × Tester mating design to derive 40 F₁ hybrids (Table 1). The hybrids and 14 parents were established in a sandy loam soil and arranged in a Randomized Complete Block Design with three replications. Three ploughings were done with cultivator to make the soil fine tilth. Thirty (30)-day-old seedlings raised in the nursery beds were transplanted on ridges with a spacing of 60 × 60 cm. 30 plants were maintained for each hybrid and parent in each replication. Cultural practices were followed uniformly for all hybrids and (TNAU Crop Production Guide, 2005). Observations were recorded from 5 randomly selected plants in each replication. The data recorded for the traits plant height, days to first flowering, number of branches per plant, fruit length, fruit circumference, number of fruit per plant, average fruit weight and fruit yield per plant for estimating heterosis. Selections were made in F₁ hybrids based on fruit shape, color, size and fruit yield per plant. Superior hybrids were selected and selfed. Seed were collected from selfed fruit and stored for further breeding.

The magnitude of heterosis in hybrids was expressed as percent of increase or decrease of a character over mid-parent (d_i), better parent (d_{ii}) and standard hybrid (d_{iii}) and estimated following the formula of Fonseca and Patterson (1968). The significance of magnitude of the relative heterosis, heterobeltiosis and standard heterosis was tested with the formula suggested by Turner (1953). Correlation coefficients were computed using formulae of Johnson et al. (1955). Path coefficients were obtained following the method of Dewey and Lu (1959).

RESULTS AND DISCUSSION

The analysis of variance revealed that parents showed significant differences for all characters (Table 2). Variance due to lines was significant for all traits indicating existence of genetic variability for growth and yield attributes among lines (females) and testers (males). The interaction between lines × testers was also significant for all the growth and yield traits.

Average performance and magnitude of heterosis for different quantitative characters varied (Table 3). In parents, T₂ (66.90 cm) had the shortest plants and L₂ (98.83 cm) was tallest. The hybrid L₁ × T₄ recorded tallest plants height, followed by L₂ × T₁ with positive heterosis over mid, better and standard parents indicating over-dominance. Among the 40 hybrids, 15 had significant positive over the mid-parent for plant height. Three hybrids had significant positive heterosis and one hybrid had significant negative heterosis (L₆ × T₃) over the better parent. Twenty-seven (27) of 40 hybrids had significant positive heterosis over standard variety and the extent of heterosis over the standard variety. Similar findings were reported by Preneetha (2002) and Thangamani (2003).

Earliness is considered an important character in any crop improvement programme, which is manifested in F₁ hybrids and preferred for commercial cultivation when high yield is coupled with earliness. The hybrid L₈ × T₁ had the shortest period of days to first flower and L₈ × T₄ had the longest period. For days to first flowering negative heterosis is desirable. Of 40 hybrids, 8 (L₁ × T₁, L₄ × T₁, L₅ × T₁, L₆ × T₁, L₆ × T₃, L₇ × T₃, L₈ × T₁ and L₁₀ × T₁) had significant heterosis in the desirable direction over the mid-parent. Seventeen (17) hybrids had significant heterobeltiosis in the negative direction (desirable). As many as 29 hybrids had heterosis in the desirable direction over the standard variety; only 2 (L₃ × T₂ and L₈ × T₄) had significant positive heterosis over the standard variety.

Number of branches per plant influences yield. Of 40 crosses, the highest number of branches per plant was for L₆ × T₂. Of 40 hybrids 21 had significant positive heterosis while three hybrids (L₂ × T₁ and L₈ × T₄) exhibited significant negative heterosis. Seventeen hybrids had significant heterosis over the better parent, of which 10 had heterosis in the positive direction and 7 (L₂ × T₁, L₂ × T₃, L₄ × T₃, L₈ × T₄, L₁₀ × T₁, L₁₀ × T₃ and L₁₀ × T₄) had heterosis in the negative direction. In 18 hybrids, there was significant heterosis in the negative direction, where-as three hybrids (L₅ × T₂, L₆ × T₂ and L₇ × T₂) had significant positive heterosis over standard variety. These results agree with findings of Preneetha (2002).

Fruit length is an important character to be considered while selecting eggplant for high yield. The longest fruit was in T₁ followed by L₈ and shortest fruit were in T₂. The hybrid L₆ × T₁ exhibited good performance and heterobeltiosis for fruit length. 21 hybrids had significant negative

Table 1. Features of parents used in study.

Name of local type	Flower bearing	Fruit bearing	Fruit shape	Fruit color	Calyx type	Calyx spininess	Source	Symbol
Lines								
Alavayal Local	Cluster	Cluster	Round	Light purple	Persistent	Non- spiny	Alavayal, Madurai D.t ,Tamil Nadu	L ₁
Sedapatty Local (Green)	Cluster	Cluster	Oval	Purplish green	Persistent	Non- spiny	Sedapatty, Madurai D.t, Tamil Nadu	L ₂
Kariapatty Local	Cluster	Cluster	Round	Green striped	Persistent	Non- spiny	Kariapatty, Vir dhunagar D.t, Tamil Nadu	L ₃
Alagarkovil Local	Cluster	Cluster	Round	Green striped	Persistent	Non- spiny	Alagarkovil, Madurai D.t, Tamil Nadu	L ₄
Palamedu Local	Cluster	Cluster	Round	Light blue	Persistent	Non- spiny	Palamedu, Madurai D.t, Tamil Nadu	L ₅
Melur Local	Cluster	Cluster	Round	Purple	Persistent	Non- spiny	Melur, Madurai D.t, Tamil Nadu	L ₆
Keerikai Local	Cluster	Cluster	Oval	Purplish green	Persistent	Non- spiny	Sempatty, Dindigul D.t, Tamil Nadu	L ₇
Nilakottai Local	Cluster	Cluster	Oblong	Green striped	Persistent	Non- spiny	Nilakottai, Dindigul D.t, Tamil Nadu	L ₈
Singampunari Local	Cluster	Cluster	Round	Purplish green	Persistent	Non- spiny	Singampunari, Sivagangai D.t, Tamil Nadu	L ₉
Sedapatty Local (Blue)	Cluster	Cluster	Round	Purple striped	Persistent	Non- spiny	Sedapatty, Madurai D.t, Tamil Nadu	L ₁₀
Tester								
Annamalai	Cluster	Cluster	Long	Purple	Non persistent	Non- spiny	Vegetable Research Station, Palur, Tamil Nadu	T ₁
KKM 1	Cluster	Cluster	Egg shaped	White	Persistent	Non- spiny	Agricultural College and Research Institute, Tuticorin, Tamil Nadu	T ₂
Punjab Sadabahar	Cluster	Cluster	Long	Purple	Non persistent	Non- spiny	Tamil Nadu agricultural university, Coimbatore	T ₃
EP 65	Cluster	Cluster	Oval	Dark purple	Non persistent	Non- spiny	Vegetable Research Station, Palur, Tamil Nadu	T ₄

heterosis over the mid-parent.

A total of 27 hybrids had significant heterosis over the better parent in desirable direction (negative). Useful heterosis was exhibited by all 40 hybrids over the standard variety. The cross combination exceeding the superior parent is a

valuable character in heterosis breeding (Patil et al., 2001). Similar findings were reported by Kaur et al. (2001).

Fruit circumference contributes to improved yield and parents L₁, L₅, T₃, T₄, and L₉ had the largest fruit. The highest fruit circumference was

recorded in L₆ x T₄ followed by L₁ x T₂ and the lowest was in L₃ x T₃. The observations were positive and significant for 14 hybrids. Positive and negative non-significant heterosis were observed in seven and four hybrids, respectively. Seven hybrids had significant and heterobeltiosis,

Table 2. Analysis of variance for parents and hybrids for vegetative and reproductive characters.

Source	df	PH ^a	DFF	NB/P	FL	FC	NF/P	AFW	FY/P
Hybrids	39	346.5306*	54.8730*	17.4039*	5.4570*	9.9952*	161.9557*	149.5525*	0.6288*
Lines	9	973.0851*	76.5412*	40.2728*	4.9768*	20.0498*	197.8137*	328.0857*	0.9020*
Testers	3	66.6626*	17.7451*	12.1612*	9.7032*	16.6776*	234.2205*	72.4168*	0.4214*
Line x Testers	27	168.7756*	51.7756*	10.3635*	5.1453*	5.9011*	141.9736*	98.6121*	0.5608*
Error	78	85.0412	3.3519	4.3914*	0.0933	0.4478	1.8176	4.0138	0.0100

*Significant at 5% level; ^aPH, plant height (cm); DFF, days to first flowering; NB/P, number of branches per plant; FL, fruit length (cm); FC, fruit circumference (cm); NF/P, number of fruit per plant; AFW, average fruit weight (g) and FY/P, fruit yield per plant (kg).

Table 3. Average performance and magnitude of heterosis for plant height and days to first flowering in eggplant.

Entry	Average value	Plant height (cm)			Average value	Days to first flowering		
		MP	BP	SV		MP	BP	SV
Alavayal Local	93.97				78.51			
Sedapatty Local (Green)	98.83				75.43			
Kariapatty Local	90.60				77.67			
Alagarkovil Local	91.93				78.02			
Palamedu Local	83.53				73.33			
Melur Local	95.20				76.89			
Keerikai Local	96.33				71.28			
Nilakottai Local	88.93				78.63			
Singampunari Local	96.60				73.11			
Sedapatty Local (Blue)	84.93				79.62			
Annamalai	73.17				81.27			
KKM 1	66.90				68.24			
Punjab Sadabahar	79.93				73.32			
EP 65	96.43				71.45			
L ₁ X T ₁	95.47	14.24	1.60	30.48**	73.67	-7.79**	-9.35**	-9.35**
L ₁ X T ₂	102.47	27.39**	9.05	40.05**	77.18	5.19**	-1.69	-5.03**
L ₁ X T ₃	104.77	20.49**	11.49	43.19**	79.28	4.43*	0.98	-2.45
L ₁ X T ₄	125.13	31.44**	29.76**	71.03**	79.60	6.16**	1.39	-2.05
L ₂ X T ₁	124.20	44.42**	25.67**	69.75**	81.13	3.55*	-0.17	-0.17
L ₂ X T ₂	102.13	23.24**	3.33	39.58**	76.13	5.98**	0.92	-6.32**
L ₂ X T ₃	110.20	23.29**	11.50	50.62**	83.64	12.46**	10.88**	2.92
L ₂ X T ₄	101.80	4.27	3.00	39.13**	76.20	3.76*	1.02	-6.24**
L ₃ X T ₁	112.27	37.11**	23.91**	53.44**	82.85	4.26*	1.94	1.94
L ₃ X T ₂	100.50	27.62**	10.93	37.36**	84.83	16.28**	9.22**	4.38*
L ₃ X T ₃	99.22	16.36*	9.51	35.61**	75.16	-0.45	-3.23	-7.52**
L ₃ X T ₄	96.70	3.40	0.28	32.16**	74.48	-0.11	-4.10*	-8.35**
L ₄ X T ₁	100.17	21.34**	8.96	36.90**	70.73	-11.19**	-12.97**	-12.97**
L ₄ X T ₂	91.63	15.38	-0.33	25.24*	71.50	-2.22	-8.35**	-12.02**
L ₄ X T ₃	101.20	17.77*	10.08	38.31**	75.00	-0.89	-3.87*	-7.72**
L ₄ X T ₄	94.20	0.02	-2.32	28.75**	74.67	-0.09	-4.29*	-8.12**
L ₅ X T ₁	86.90	10.91	4.03	18.77	70.89	-8.29**	-12.77**	-12.77**
L ₅ X T ₂	85.53	13.72	2.39	16.90	74.26	4.91**	1.26	-8.63**
L ₅ X T ₃	95.60	16.97*	14.45	30.66**	78.12	6.54**	6.53**	-3.87*
L ₅ X T ₄	83.90	-6.76	-13.00	14.67	72.65	0.36	-0.93	-10.61**
L ₆ X T ₁	91.77	9.01	-3.61	25.42*	74.28	-6.07**	-8.60**	-8.60**
L ₆ X T ₂	85.90	5.98	-9.77	17.40	70.94	-2.24	-7.74**	-12.71**
L ₆ X T ₃	72.33	-17.40*	-24.02**	-1.14	71.87	-4.31*	-6.53**	-11.57**
L ₆ X T ₄	93.90	-2.00	-2.63	28.34**	73.55	-0.83	-4.34*	-9.50**
L ₇ X T ₁	96.73	14.14	0.42	32.21**	78.66	3.13	-3.21	-3.21
L ₇ X T ₂	103.20	26.44**	7.13	41.05**	76.05	9.02**	6.69**	-6.42**
L ₇ X T ₃	106.93	21.33**	11.00	46.15**	69.25	-4.22*	-5.55**	-14.79**
L ₇ X T ₄	97.23	0.88	0.83	32.89**	76.45	7.13**	7.00**	-5.93**
L ₈ X T ₁	93.77	15.69*	5.43	28.15**	68.69	-14.08**	-15.48**	-15.48**
L ₈ X T ₂	86.60	11.14	-2.62	18.36	78.50	6.90**	-0.17	-3.41
L ₈ X T ₃	96.67	14.49	8.70	32.12**	75.46	-0.68	-4.03*	-7.15**
L ₈ X T ₄	96.83	4.48	0.41	32.35**	87.98	17.25**	11.90**	8.26**
L ₉ X T ₁	86.53	1.94	-10.42	18.27	75.51	-2.18	-7.09**	-7.09**

*, **Significant at 5 and 1% level.

Table 3. Contd.

Entry	Average value	Number of branches per plant			Average value	Fruit length (cm)		
		MP	BP	SV		MP	BP	SV
L ₉ X T ₂	93.20	14.01	-3.52	27.38**	73.94	4.62*	1.14	-9.02**
L ₉ X T ₃	87.29	-1.11	-9.64	19.30	77.70	6.13**	5.97**	-4.39*
L ₉ X T ₄	84.33	-12.62	-12.70	15.26	72.22	-0.08	-1.22	-11.14**
L ₁₀ X T ₁	84.25	6.57	-0.81	15.14	75.31	-6.38**	-7.33**	-7.33**
L ₁₀ X T ₂	85.33	12.40	0.47	16.63	76.38	3.32	-4.07*	-6.01**
L ₁₀ X T ₃	87.27	5.86	2.75	19.27	78.54	2.71	-1.36	-3.36
L ₁₀ X T ₄	82.50	-9.02	-14.45	12.76	82.21	8.84**	3.25	1.16
SEd	7.25	6.23	7.20	1.52		1.31	1.51	
Alavayal Local	21.13				7.44			
Sedapatty Local (Green)	22.32				7.54			
Kariapatty Local	16.36				6.93			
Alagarkovil Local	20.49				6.53			
Palamedu Local	20.04				7.87			
Melur Local	21.89				7.14			
Keerikai Local	22.11				7.79			
Nilakottai Local	21.74				10.23			
Singampunari Local	17.88				5.41			
Sedapatty Local (Blue)	21.62				6.92			
Annamalai	24.18				11.61			
KKM 1	15.31				5.25			
Punjab Sadabahar	18.50				6.85			
EP 65	18.01				8.07			
L ₁ X T ₁	26.36	16.36**	9.03	9.03	6.58	-30.74**	-43.28**	-43.28**
L ₁ X T ₂	23.70	27.71**	12.15	-1.99	6.92	9.30**	-6.57*	-40.41**
L ₁ X T ₃	20.02	1.03	-5.25	-17.19**	6.83	-4.16	-7.74*	-41.15**
L ₁ X T ₄	19.63	0.32	-7.08	-18.79**	8.17	5.58*	1.20	-29.61**
L ₂ X T ₁	19.12	-17.74**	-20.90**	-20.90**	7.78	-18.73**	-32.97**	-32.97**
L ₂ X T ₂	19.74	3.11	-11.53	-18.34**	5.86	-8.39**	-22.28**	-49.51**
L ₂ X T ₃	18.81	-7.82	-15.70*	-22.18**	6.63	-7.85**	-12.07**	-42.88**
L ₂ X T ₄	19.60	-2.82	-12.19	-18.94**	8.00	2.48	-0.91	-31.07**
L ₃ X T ₁	22.87	12.83*	-5.40	-5.40	6.16	-33.57**	-46.96**	-46.96**
L ₃ X T ₂	20.71	28.08**	26.58**	-14.33*	6.52	7.03*	-5.92	-43.83**
L ₃ X T ₃	16.80	-3.63	-9.21	-30.51**	6.97	1.16	0.58	-39.95**
L ₃ X T ₄	19.46	13.22	8.03	-19.51**	6.54	-12.77**	-18.95**	-43.62**
L ₄ X T ₁	26.52	18.75**	9.69	9.69	6.34	-30.11**	-45.40**	-45.40**
L ₄ X T ₂	25.50	39.84**	24.45**	5.47	7.34	24.62**	12.46**	-36.76**
L ₄ X T ₃	17.25	-11.54	-15.83*	-28.66**	5.88	-12.14**	-14.21**	-49.37**
L ₄ X T ₄	18.09	-6.05	-11.73	-25.19**	9.84	34.79**	21.88**	-15.22**
L ₅ X T ₁	23.09	4.45	-4.49	-4.49	6.00	-38.41**	-48.33**	-48.33**
L ₅ X T ₂	27.09	50.41**	35.19**	12.04*	6.41	-2.29	-18.52**	-44.77**
L ₅ X T ₃	26.78	38.97**	33.65**	10.77	6.74	-8.40**	-14.32**	-41.93**
L ₅ X T ₄	25.92	36.24**	29.36**	7.21	7.86	-1.38	-2.64	-32.28**
L ₆ X T ₁	23.07	0.16	-4.58	-4.58	4.36	-53.46**	-62.41**	-62.41**
L ₆ X T ₂	28.53	50.67**	30.33**	18.01**	10.15	63.75**	42.09**	-12.55**
L ₆ X T ₃	25.08	24.18**	14.57*	3.74	6.32	-9.72**	-11.57**	-45.58**
L ₆ X T ₄	24.20	21.29**	10.55	0.10	5.44	-28.54**	-32.66**	-53.16**
L ₇ X T ₁	23.69	2.35	-2.03	-2.03	5.58	-42.47**	-51.92**	-51.92**
L ₇ X T ₂	27.53	44.55**	24.51**	13.87*	7.88	20.80**	1.11	-32.11**
L ₇ X T ₃	22.75	12.02*	2.88	-5.91	5.74	-21.60**	-26.35**	-50.55**
L ₇ X T ₄	23.85	18.87**	7.85	-1.36	7.90	-0.46	-2.19	-31.96**
L ₈ X T ₁	26.83	16.88**	10.99	10.99	6.87	-37.10**	-40.84**	-40.84**
L ₈ X T ₂	23.02	22.07**	5.90	-4.77	5.89	-23.90**	-42.41**	-49.25**
L ₈ X T ₃	22.22	10.43	2.21	-8.09	10.70	25.32**	4.63*	-7.81**
L ₈ X T ₄	16.21	-18.45**	-25.44**	-32.95**	8.83	-3.46	-13.62**	-23.89**
L ₉ X T ₁	28.24	34.29**	16.81**	16.81**	5.50	-35.34**	-52.61**	-52.61**
L ₉ X T ₂	19.49	15.14*	9.02	-19.37**	5.69	6.82	5.30	-50.95**
L ₉ X T ₃	22.91	25.94**	23.82**	-5.24	6.78	10.63**	-1.02	-41.59**
L ₉ X T ₄	20.75	15.60*	15.17	-14.19*	6.74	0.00	-16.52**	-41.93**
L ₁₀ X T ₁	15.29	-33.24**	-36.77**	-36.77**	5.91	-36.20**	-49.08**	-49.08**
L ₁₀ X T ₂	19.97	6.25	-7.60	-17.39**	6.30	3.45	-9.01**	-45.75**
L ₁₀ X T ₃	18.83	-6.15	-12.91*	-22.13**	5.12	-25.64**	-26.01**	-55.89**
L ₁₀ X T ₄	18.15	-8.40	-16.04*	-24.93**	5.62	-25.03**	-30.39**	-51.58**
SEd	1.65	1.22		1.41	0.23	0.20		0.23
CD at 1%	4.25				0.60			

*, **Significant at 5 and 1% level.

Table 3. Contd.

Entry	Average value	Average fruit weight (g)			Average value	Fruit yield per plant (kg)		
		MP	BP	SV		MP	BP	SV
CD at 5%	3.23		0.46	CD at 5%	3.23			
Alavayal Local	66.32				1.72			
Sedapatty Local (Green)	56.09				1.86			
Kariapatty Local	41.05				1.27			
Alagarkovil Local	46.01				1.26			
Palamedu Local	81.66				1.91			
Melur Local	55.72				1.75			
Keerikai Local	65.76				1.79			
Nilakottai Local	50.01				1.27			
Singampunari Local	34.94				1.16			
Sedapatty Local (Blue)	54.17				1.37			
Annamalai	55.9				2.12			
KKM 1	42.42				1.46			
Punjab Sadabahar	50.34				1.56			
EP 65	53.38				1.36			
L ₁ X T ₁	69.40	13.57**	4.65	24.15**	2.47	28.65**	16.33**	16.33**
L ₁ X T ₂	54.62	0.47	-17.63**	-2.28	1.35	-15.13**	-21.55**	-36.58**
L ₁ X T ₃	65.78	12.77**	-0.81	17.67**	1.35	-17.60**	-21.36**	-36.42**
L ₁ X T ₄	51.83	-11.14**	-21.84**	-7.27*	1.45	-5.42	-15.34**	-31.55**
L ₂ X T ₁	48.00	-14.28**	-14.42**	-14.13**	2.07	3.77	-2.67	-2.67
L ₂ X T ₂	54.32	10.29**	-3.15	-2.82	1.42	-14.17**	-23.48**	-32.97**
L ₂ X T ₃	57.82	8.64**	3.08	3.43	2.14	24.95**	14.87**	0.63
L ₂ X T ₄	52.24	-1.83	-6.86*	-6.54*	1.56	-2.80	-15.95**	-26.37**
L ₃ X T ₁	43.50	-10.26**	-22.18**	-22.18**	1.45	-14.45**	-31.71**	-31.71**
L ₃ X T ₂	44.44	6.49	4.76	-20.50**	0.99	-27.05**	-31.81**	-53.22**
L ₃ X T ₃	61.99	35.65**	23.13**	10.89**	1.74	23.11**	11.54*	-18.05**
L ₃ X T ₄	47.41	3.76	-5.82	-15.19**	0.93	-29.35**	-31.70**	-56.36**
L ₄ X T ₁	60.91	19.54**	8.97**	8.97**	2.40	41.73**	13.03**	13.03**
L ₄ X T ₂	52.64	19.04**	14.39**	-5.84*	2.32	70.34**	59.04**	9.11*
L ₄ X T ₃	53.38	10.79**	6.03	-4.51	1.49	5.79	-4.27	-29.67**
L ₄ X T ₄	58.18	20.76**	15.57**	4.08	1.92	46.31**	41.28**	-9.73*
L ₅ X T ₁	66.13	-3.85	-19.02**	18.30**	1.71	-15.44**	-19.62**	-19.62**
L ₅ X T ₂	59.64	-3.87	-26.97**	6.68*	2.34	38.87**	22.30**	10.20**
L ₅ X T ₃	56.04	-15.10**	-31.38**	0.24	2.36	36.08**	23.52**	11.30**
L ₅ X T ₄	51.17	-22.47**	-37.34**	-8.47**	2.21	35.17**	15.51**	4.08
L ₆ X T ₁	51.64	-7.47**	-7.62**	-7.62**	1.48	-23.65**	-30.30**	-30.30**
L ₆ X T ₂	55.96	14.05**	0.44	0.11	2.24	39.56**	27.76**	5.49
L ₆ X T ₃	53.48	0.84	-4.02	-4.34	1.35	-18.31**	-22.81**	-36.26**
L ₆ X T ₄	51.00	-3.83	-8.47**	-8.77**	1.68	7.82	-4.37	-21.04**
L ₇ X T ₁	56.55	-7.04**	-14.01**	1.16	1.13	-42.25**	-46.78**	-46.78**
L ₇ X T ₂	69.49	28.46**	5.67*	24.31**	2.85	75.36**	59.03**	34.07**
L ₇ X T ₃	54.50	-6.11*	-17.12**	-2.50	1.43	-14.43**	-19.93**	-32.50**
L ₇ X T ₄	57.67	-0.65	-12.30**	3.17	1.74	10.38*	-2.98	-18.21**
L ₈ X T ₁	42.04	-20.61**	-24.79**	-24.79**	2.26	33.14**	6.28	6.28
L ₈ X T ₂	42.74	-7.51*	-14.53**	-23.54**	1.66	21.91**	13.96*	-21.82**
L ₈ X T ₃	40.84	-18.60**	-18.87**	-26.94**	1.04	-26.18**	-33.12**	-50.86**
L ₈ X T ₄	49.54	-1.27	-1.60	-11.38**	2.06	57.31**	52.09**	-2.83
L ₉ X T ₁	46.66	2.73	-16.53**	-16.53**	2.29	39.15**	7.69*	7.69*
L ₉ X T ₂	46.87	21.17**	10.49**	-16.15**	1.73	31.81**	18.54**	-18.68**
L ₉ X T ₃	56.43	32.34**	12.09**	0.95	1.22	-10.40*	-21.79**	-42.54**
L ₉ X T ₄	50.54	18.54**	0.40	-9.58**	1.42	12.43*	4.42	-33.28**
L ₁₀ X T ₁	57.01	3.59	1.98	1.98	1.12	-35.82**	-47.25**	-47.25**
L ₁₀ X T ₂	52.21	8.11**	-3.61	-6.60*	1.85	30.81**	26.77**	-13.03**
L ₁₀ X T ₃	61.70	18.07**	13.91**	10.38**	1.43	-2.51	-8.55	-32.81**
L ₁₀ X T ₄	56.59	8.29**	4.47	1.23	1.45	6.73	6.34	-31.55**
SEd	1.60	1.39		1.60	0.08	0.07		0.08
CD at 1%	4.15				0.21			
CD at 5%	3.15				0.16			

*, **Significant at 5% and 1% level.

24 had significant and negative heterosis. Thirty-one (31) hybrids had significant and positive heterosis over the

standard variety. Four hybrids had non-significant and negative values. The findings corroborate the results of

Table 4. Genotypic correlation between fruit yield and growth characters in eggplant.

Parameter	Days to first flowering	Number of branches per plant	Fruit yield per plant (kg)
Plant height (cm)	0.454*	-0.125	0.007
Days to first flowering		-0.385*	-0.211
Number of branches per plant			0.598*

*Significant at 5% level.

Nalini (2007).

The number of fruit per plant influences yield. The lowest number of fruit per plant was in L_5 and the highest in T_1 followed by L_9 among parents. Nineteen crosses, $L_1 \times T_1$, $L_2 \times T_1$, $L_2 \times T_3$, $L_4 \times T_1$, $L_4 \times T_2$, $L_4 \times T_4$, $L_5 \times T_2$, $L_5 \times T_3$, $L_5 \times T_4$, $L_6 \times T_2$, $L_6 \times T_4$, $L_7 \times T_2$, $L_7 \times T_4$, $L_8 \times T_1$, $L_8 \times T_2$, $L_8 \times T_4$, $L_9 \times T_1$, $L_9 \times T_2$ and $L_{10} \times T_2$ exhibited positive heterosis in the desirable direction over the mid-parent. Expression of heterosis over the better parent in the positive direction occurred in 14 crosses. Significant heterosis over commercial check was recorded in 33 hybrids with 9 had positive and 23 had negative heterosis. Considerable positive heterosis in different cross combinations were also reported by Chowdhury et al. (2010).

The highest single fruit weight was for $L_7 \times T_2$ followed by $L_1 \times T_1$ and the lowest was in $L_8 \times T_3$. Of 40 hybrids, 17 had significant desirable heterosis over the mid-parent in the positive direction. Only 7 hybrids had significant heterobeltiosis. In 8 hybrids, heterosis was significant and in the positive direction over the standard variety. The heterosis over the better and standard parents was negative for fruit weight. Similar findings reported by Nalinidharwad et al. (2011) and Patil et al. (2001) lend support to the present results. This trend is not a constraint, because smaller sized eggplant fruit are preferred in South India and hybrids with smaller and more fruit could be selected.

The ultimate interest of the breeder is to get high yield. An appreciable amount of heterosis in F_1 s over the mid-parent value occurred for fruit yield per plant. The T_1 parent had the highest yield followed by L_5 and L_2 , while the lowest yield was for L_9 . Fruit yield per plant showed a wide range among hybrids. Among the 40 hybrids, 19 had significant positive heterosis over the mid-parent. The hybrid from the cross $L_7 \times T_2$ had the highest magnitude of heterosis in the positive direction followed by $L_4 \times T_2$, $L_8 \times T_4$ and $L_4 \times T_4$. Most crosses involving T_2 as tester parent had significant, positive, heterosis over the mid-parent. This agrees with Ramesh Kumar et al. (2013).

Association analysis

Genotypic correlation coefficients varied depending on character (Tables 4 to 8). Of 15 characters studied, only

numbers of fruit per plant, number of branches per plant and average fruit weight had significant association with fruit yield at phenotypic and genotypic levels (Prabakaran, 2010).

Fruit yield had significant negative association with fruit borer infestation at both levels. This trait could be considered as an important criterion for selection for fruit yield. Total phenol content had considerable non-significant positive correlation with fruit yield per plant followed by fruit length, fruit circumference and shoot borer infestation. During selection these traits can be considered to improve fruit yield. There were significant, and positive, correlations for plant height with days to first flowering, while days to first flowering significantly, and negatively, correlated with numbers of branches per plant. As days to first flowering decreases, there will be an increase in number of branches per plant. Fruit pedicel length had a significant, and positive, correlation with calyx length. Fruit circumference was significantly, and positively, associated with average fruit and average fruit weight may result in simultaneous improvement of fruit yield per plant (Thangamani and Jansirani, 2012). These characters are highly reliable components of fruit yield and could be utilized as yield indicators during selection. Days to first flowering, fruit pedicel length, calyx length, fruit borer infestation and ascorbic acid, an important quality trait, were negatively associated with fruit yield per plant. Selection for this trait will result in reduction weight. Similarly, the association between shoot borer infestation and little leaf incidence was positive and significant. Selection for this trait will result in reduction of fruit yield, number of branches per plant, number of fruit per plant of fruit yield.

Path coefficient analysis

The path coefficient analysis permits the separation of direct effect from indirect effects through other related traits by partitioning genotypic correlation coefficients. Plant height, number of branches per plant, fruit length, fruit pedicel length, number of fruit per plant, average fruit weight and little leaf incidence had positive direct effect on yield (Table 9).

Direct selection for these characters are likely will bring about an overall improvement in fruit yield per plant. The residual effect determines how causal factors account

Table 5. Genotypic correlation between fruit yield and fruit characters in eggplant.

Parameter	Fruit pedicel length (cm)	Fruit circumference (cm)	Calyx length (cm)	Number of fruits per plant	Average fruit weight (g)	Fruit yield per plant (kg)
Fruit length (cm)	0.101	-0.185	0.181	0.139	0.094	0.200
Fruit pedicel length (cm)	-	0.034	0.248*	-0.185	0.088	-0.124
Fruit circumference (cm)	-	-	-0.011	0.061	0.273*	0.154
Calyx length (cm)	-	-	-	-0.094	0.161	-0.023
Number of fruit per plant	-	-	-	-	-0.148	0.836*
Average fruit weight (g)	-	-	-	-	-	0.380*

*Significant at 5% level.

Table 6. Phenotypic correlation between fruit yield and fruit characters in eggplant.

Parameter	Fruit pedicel length (cm)	Fruit circumference (cm)	Calyx length (cm)	Number of fruits per plant	Average fruit weight (g)	Fruit yield per plant (kg)
Fruit length (cm)	0.084	-0.169	0.147	0.150	0.085	0.205
Fruit pedicel length (cm)	-	0.027	0.247*	-0.186	0.084	-0.125
Fruit circumference (cm)	-	-	-0.019	0.064	0.247*	0.147
Calyx length (cm)	-	-	-	-0.108	0.161	-0.034
Number of fruit per plant	-	-	-	-	-0.140	0.831*
Average fruit weight (g)	-	-	-	-	-	0.381*

*Significant at 5% level.

Table 7. Genotypic correlation between fruit yield, pest and disease incidence and quality characters in eggplant.

Parameter	Fruit borer infestation (%)	Little leaf incidence (%)	Ascorbic acid content (mg/100g)	Total phenols content (mg/100g)	Fruit yield per plant (kg)
Shoot borer infestation (%)	-0.007	0.400*	-0.053	0.141	0.102
Fruit borer infestation (%)	-	-0.048	0.084	0.187	-0.280*
Little leaf incidence (%)	-	-	0.048	0.208	0.043
Ascorbic acid content (mg/100 g)	-	-	-	-0.151	-0.045
Total phenols content (mg/100 g)	-	-	-	-	0.227

*Significant at 5% level.

Table 8. Phenotypic correlation between fruit yield, pest and disease incidence and quality characters in eggplant.

Parameter	Fruit borer infestation (%)	Little leaf incidence (%)	Ascorbic acid content (mg/100 g)	Total phenols content (mg/100 g)	Fruit yield per plant (kg)
Shoot borer infestation (%)	-0.009	0.393*	-0.047	0.137	0.094
Fruit borer infestation (%)		-0.054	0.065	0.168	-0.275*
Little leaf incidence (%)			0.057	0.203	0.047
Ascorbic acid content (mg/100 g)				-0.151	-0.036
Total phenols content (mg/100 g)					0.215

* Significant at 5% level.

Table 9. Direct and indirect effects of different characters on fruit yield in eggplant.

Character	X1 ^a	X2	X3	X4	X5	X6	X7	X8	X9	X10	X11	X12	X13	X14	Fruit yield per plant
X1	0.0072	-0.0123	-0.0022	0.0000	-0.0006	-0.0012	0.0019	0.0429	-0.0058	0.0025	0.0119	0.0061	-0.0040	-0.0055	0.039
X2	0.0027	-0.0328	-0.0130	0.0001	-0.0002	0.0037	0.0012	-0.0581	-0.0455	0.0015	-0.0035	0.0042	0.0005	0.0027	-0.138
X3	-0.0002	0.0073	0.0583	-0.0000	-0.0002	-0.0046	0.0012	0.0370	0.0864	0.0000	-0.0096	0.0041	-0.0039	-0.0045	0.166
X4	0.0008	-0.0021	0.0152	0.0003	-0.0005	0.0025	0.0005	-0.0304	0.1915	0.0025	0.0111	-0.0018	0.0013	-0.0025	0.169
X5	0.0001	-0.0047	-0.0030	0.0011	0.0004	0.0070	-0.0011	0.0384	0.0539	0.0034	0.0084	-0.0042	0.0029	0.0027	0.100
X6	-0.0010	0.0016	-0.0035	0.0001	0.0047	-0.0012	-0.0015	-0.1185	0.0503	0.0034	0.0035	-0.0062	-0.0013	0.0055	-0.062
X7	0.0002	0.0032	0.0071	-0.0002	0.0001	-0.0379	0.0000	0.0898	0.1571	0.0020	0.0064	-0.0002	-0.0000	-0.0001	0.229
X8	-0.0022	0.0065	-0.0115	0.0002	0.0011	0.0004	-0.0064	-0.0532	0.0928	0.0055	-0.0005	-0.0118	0.0013	0.0058	0.030
X9	0.0003	0.0022	0.0025	0.0000	-0.0006	-0.0040	0.0004	0.8516	-0.0525	-0.0004	0.0002	0.0011	-0.0004	-0.0007	0.801*
X10	-0.0000	0.0026	0.0087	0.0001	0.0004	-0.0103	-0.0010	-0.0777	0.5758	0.0024	0.0206	-0.0003	-0.0015	-0.0040	0.509*
X11	-0.0015	0.0041	-0.0004	-0.0003	-0.0013	0.0061	0.0028	0.0275	-0.1143	-0.0124	0.0010	0.0088	0.0010	-0.0024	-0.077
X12	-0.0015	-0.0020	0.0098	-0.0001	-0.0002	0.0042	-0.0000	-0.0042	-0.2072	0.0002	-0.0572	-0.0001	-0.0010	0.0025	-0.253*
X13	0.0019	-0.0060	0.0106	-0.0002	-0.0012	0.0004	0.0033	0.0408	-0.0074	-0.0048	0.0002	0.0229	-0.0003	-0.0044	0.057
X14	0.0020	0.0012	0.0154	-0.0002	0.0004	-0.0001	0.0005	0.0237	0.0615	0.0009	-0.0038	0.0005	-0.0148	0.0035	0.093
X15	0.0021	0.0046	0.0137	-0.0001	-0.0013	-0.0003	0.0019	0.0324	0.1216	-0.0016	0.0075	0.0053	0.0027	-0.0191	0.167

*Significant at 5% level; Residual effect, 0.102; Diagonal values, direct effects; Half diagonal values, indirect effects. ^a X1, Plant height (cm); X2, Days to first flowering; X3, Number of branches per plant; X4, Fruit length (cm); X5, Fruit pedicel length (cm); X6, Fruit circumference (cm); X7, Calyx length (cm); X8, Number of fruit per plant; X9, Average fruit weight (g); X10, Shoot borer infestation (%); X11, Fruit borer infestation (%); X12, Little leaf incidence (%); X13, Ascorbic acid content (mg/100g); and X14, Total phenols content (mg/100g).

for variability of the dependent factor, that is, fruit yield per plant in this study. The residual effect was only 0.102, accounting for 97.50% of variability in fruit yield per plant, was explained by the 15 variables included in the study.

The characters studied by path analysis for yield appear to be appropriate. Plant height, number of branches per plant, fruit length, fruit pedicel length, number of fruit per plant, average fruit weight and little leaf incidence are important characters to

bring about overall improvement in fruit yield per plant. Similar trend of findings were also obtained by Thangamani and Jansirani (2012) for fruit yield per plant, number of branches per plant and average fruit weight in brinjal.

Conclusion

Promising lines are to be used to obtain higher yield, earliness and increased fruit number. Due to high heterosis the importance of non-additive genetic effects in expression can be inferred. The establishment of a population with a wide genetic base, using recurrent selection methods for increasing combining ability, will lead to future new lines which result in hybrids superior to those studied. The hybrids L7xT2, L1xT1, L4 xT1, L5xT3, L5xT2, L8 x T4 and L4 x T2 had the highest values over standard variety heterosis. These hybrids can be utilized for selecting superior desirable segregants in later generations. Correlation and path analysis indicated that more fruit per plant with heavier weight are contributors to improved yield. At the outset comprehensive results obtained from the correlation and path analysis indicated that more number of fruits per plant with more fruit weight are outstanding contributors made by the hybrids for yield per plant. Therefore, while selecting the hybrids due weightage may given to the above said traits for the overall improvement of fruit yield.

ACKNOWLEDGEMENTS

The author wishes to thank research institutes viz., Vegetable Research Station, Palur, Agricultural College and Research Institute, Killikulam, Madurai and TNAU, Coimbatore for providing the germplasms to carry out the present investigation. Dr. V. Premalakshmi, Assistant Professor (Horticulture), Agricultural College and Research Institute, Madurai is duly acknowledged for giving valuable suggestions for this dissertation research.

REFERENCES

- Chowdhury MJ, Ahmad S, Nazim Uddin M, Quamruzzaman AKM, Patwary MMA (2010). Expression of heterosis for productive traits in F1 brinjal (*Solanum melongena* L.) hybrids. *The Agriculturists* 8(2):8-13.
- Daliya MS, WM Wilson (2002). Genetical study for yield and quality in aubergine (*Solanum melongena* L.). *Philipp. Agric.* 125:117-118.
- Dewey DR, Lu KH (1959). A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.* 51:515-518.
- Fonseca S, Patterson FL (1968). Hybrid vigour in a seven parent diallel crosses in common winter wheat (*Triticum aestivum* L.). *Crop Sci.* 8:85-88.
- Johnson WW, Robinson HF, Comstock RE (1955). Genotypic and phenotypic correlation in soybeans and their implications in selection. *Agron. J.* 47:477-482.
- Kakikazi Y (1931). Hybrid vigour in *Solanum melongena* L. *Japanese Agric. Horticult.* 3:371-380.
- Kaur J, Patel JA, Patel, MJ, Acharya RR, Bhanvadia AS (2001). Genetic analysis of earliness and plant stature in brinjal. *Capsicum Eggplant Newslett.* 20:94-97.
- Nalini D (2007). Studies on heterosis and combining ability in brinjal (*Solanum melongena* L.). M.Sc (Hort.) Thesis, University of Agricultural Sciences, Dharwad, India.
- Nalinidharwad S, Patil SA, Salimath PM (2011). Study on genetic diversity and its relation to heterosis in brinjal (*Solanum melongena* L.). *Karnataka J. Agric. Sci.* 24 (2):110-113.
- Patil SD, Warade SD, Badgujar CD (2001). Heterosis studies in brinjal (*Solanum melongena* L.). *J. Soil Crops* 11(1):47- 51.
- Prabakaran S (2010). Evaluation of local types of brinjal (*Solanum melongena* L.). M.Sc., (Hort.) Thesis, Agricultural College and Research Institute, TNAU, Madurai.
- Preneetha S (2002). Breeding for shoot and fruit borer (*Leucinodes orbonalis* G.) resistance in brinjal (*Solanum melongena* L.). Ph.D (Hort.) Thesis, Tamil Nadu Agricultural University, Coimbatore, India.
- Ramesh Kumar S, Arumugam T, Premalakshmi V, Anandakumar CR, Rajavel DS (2013). Out breeding for yield and horticultural attributes in indigenous eggplant germplasm. *African J. Agric. Res.* 8(29):4099-4110.
- Thangamani C (2003). Evaluation of F₁ brinjal (*Solanum melongena* L.) for yield and quality. M.Sc. (Hort.) Thesis, Tamil Nadu Agricultural University, Coimbatore, India.
- Thangamani C, Jansirani P (2012). Correlation and path coefficient analysis studies on yield and attributing characters in brinjal (*Solanum melongena* L.). *Electron J. Plant. Breed.* 3(3):939-944.
- Turner JR (1953). A study on heterosis in upland cotton II. Combining ability and inbreeding effects. *Agronomy J.* 45:487-490.