



In silico identification of genetic variants in glucocerebrosidase (GBA) gene involved in Gaucher's disease using multiple software tools

Madhumathi Manickam, Palaniyandi Ravanan, Pratibha Singh and Priti Talwar*

Apoptosis and Cell Death Research Laboratory, Centre for Biomedical Research, School of Biosciences and Technology, Vellore Institute of Technology University, Vellore, India

Edited by:

Babajan B., King Abdulaziz University, Saudi Arabia

Reviewed by:

Dhananjai M. Rao, Miami University, USA

Indira Shrivastava, University of Pittsburgh, USA

*Correspondence:

Dr. Priti Talwar, Apoptosis and Cell Death Research Laboratory, Centre for Biomedical Research, School of Biosciences and Technology, Vellore Institute of Technology University, Vellore, Tamil Nadu-632 014, India
e-mail: priti.t@vit.ac.in

Gaucher's disease (GD) is an autosomal recessive disorder caused by the deficiency of glucocerebrosidase, a lysosomal enzyme that catalyses the hydrolysis of the glycolipid glucocerebroside to ceramide and glucose. Polymorphisms in GBA gene have been associated with the development of Gaucher disease. We hypothesize that prediction of SNPs using multiple state of the art software tools will help in increasing the confidence in identification of SNPs involved in GD. Enzyme replacement therapy is the only option for GD. Our goal is to use several state of art SNP algorithms to predict/address harmful SNPs using comparative studies. In this study seven different algorithms (SIFT, MutPred, nsSNP Analyzer, PANTHER, PMUT, PROVEAN, and SNPs&GO) were used to predict the harmful polymorphisms. Among the seven programs, SIFT found 47 nsSNPs as deleterious, MutPred found 46 nsSNPs as harmful. nsSNP Analyzer program found 43 out of 47 nsSNPs are disease causing SNPs whereas PANTHER found 32 out of 47 as highly deleterious, 22 out of 47 are classified as pathological mutations by PMUT, 44 out of 47 were predicted to be deleterious by PROVEAN server, all 47 shows the disease related mutations by SNPs&GO. Twenty two nsSNPs were commonly predicted by all the seven different algorithms. The common 22 targeted mutations are F251L, C342G, W312C, P415R, R463C, D127V, A309V, G46E, G202E, P391L, Y363C, Y205C, W378C, I402T, S366R, F397S, Y418C, P401L, G195E, W184R, R48W, and T43R.

Keywords: glucocerebrosidase, SIFT, MutPred, PANTHER, PMUT, PROVEAN, SNPs&GO

INTRODUCTION

Gaucher's disease (GD) is a rare genetic disease in which fatty substances accumulate in cells and certain organs (James et al., 2006). It is a common lysosomal storage disorder and results from an inborn deficiency of the enzyme glucocerebrosidase (also known as acid β -glucosidase). This enzyme is responsible for glucocerebroside (glucosylceramide) degradation. The accumulation of undegraded substrate generally happens because of enzyme deficiency, mainly within cells of the macrophage lineage or monocyte, and it is responsible for the clinical manifestations of the disease (Beutler and Grabowski, 2001). This glucosylceramide degrading enzyme is encoded by a gene named GBA, which is 7.6 kb in length and located in 1q21 locus. Recessive mutation in GBA gene affects both males and females (Horowitz et al., 1989; Zimran et al., 1991; Winfield et al., 1997). GBA protein is 497 amino acids long with the molecular weight of 55.6 KD. GBA enzyme catalyses the breakdown of glucosylceramide, a cell membrane constituent of white blood cells and red blood cells. The macrophages fail to eliminate the waste product and results in accumulation of lipids in fibrils and this turn into Gaucher cells (Aharon et al., 2004). GD can be classified into three classes namely types 1, 2, and 3. In type 1, Glucosylceramide accumulate in visceral organs whereas in type 2 and 3, the accumulation is in the central nervous system (Grabowski, 2008).

The international disease frequency of GD is 200,000 except for areas of the world with large Ashkenazi Jewish populations where 60% of the patients are estimated to be homozygous, which accounts for 75% of disease alleles (Pilar et al., 2012). Almost 300 unique mutations have been reported in the GBA gene, with distribution that spans the entire gene. These include 203 missense mutations, 18 nonsense mutations, 36 small insertions or deletions that lead to frameshift or in-frame alterations, 14 splice junction mutations and 13 complex alleles carrying two or more mutations (Hruska et al., 2008). The single nucleotide variations in the genome that occur at a frequency of more than 1% are referred as single nucleotide polymorphisms (SNPs) and in the human genome, SNPs occur in just about every 3000 base pairs (Cargill et al., 1999).

Nearly 200 mutations in the GBA gene have been described in patients with GD types 1, 2, and 3 (Jmoudiak and Futerman, 2005). L444P mutation was identified in GBA gene in patients with GD types 1, 2, and 3. The L444P substitution is one of the major SNP associated with the GBA gene. D409H, A456P, and V460V mutations were also identified in patients with GD (Tsuji et al., 1987; Latham et al., 1990). Previous findings have shown that, in 60 patients with types 1 and 3, the most common Gaucher mutations identified were N370S, L444P, and R463C (Sidransky et al., 1994). The other mutation E326K had been identified in patients with all three types of GD, but in each instance it was

found on the same allele with another GBA mutation. Also, Park et al. identified the E326K allele in 1.3% of patients with GD and in 0.9% of controls, indicating that it is a polymorphism (Park et al., 2002).

The harmful SNPs for the GBA gene have not been predicted to date *in silico*. Therefore we designed a strategy for analyzing the entire GBA coding region. Different algorithms such as SIFT (Ng and Henikoff, 2001), MutPred (Li et al., 2009), nsSNP Analyzer (Bao et al., 2005), PANTHER (Mi et al., 2012), PMUT (Costa et al., 2002), PROVEAN (Choi et al., 2012), and SNPs&GO (Calabrese et al., 2009) were utilized to predict high-risk nonsynonymous single nucleotide polymorphisms (nsSNPs) in coding regions that are likely to have an effect on the function and structure of the protein.

MATERIALS AND METHODS

DATA SET

SNPs associated with GBA gene were retrieved from the single nucleotide polymorphism database (dbSNP) (<http://www.ncbi.nlm.nih.gov/snp/>), and are commonly referred by their reference sequence IDs (rsID) (Wheeler et al., 2005).

VALIDATION OF TOLERATED AND DELETERIOUS SNPs

The type of genetic mutation that causes a single amino acid substitution (AAS) in a protein sequence is called nsSNP. An nsSNP could potentially influence the function of the protein, subsequently altering the phenotype of carrier. This protocol describes the use of the Sorting Intolerant From Tolerant (SIFT) algorithm (<http://sift.jcvi.org>) for predicting whether an AAS affects protein function. To assess the effect of a substitution, SIFT assumes that important positions in a protein sequence have been conserved throughout evolution and therefore at these positions substitutions may affect protein function. Thus, by using sequence homology, SIFT predicts the effects of all possible substitutions at each position in the protein sequence. The protocol typically takes 5–20 min, depending on the input (Kumar et al., 2009).

PREDICTION OF HARMFUL MUTATIONS

MutPred (<http://mutdb.org/mutpred>) models structural features and functional sites changes between mutant sequences and wild-type sequence. These changes are expressed as probabilities of gain or loss of structure and function. The MutPred output contains a general score (g), i.e., the probability that the AAS is deleterious/disease-associated and top five property scores (p), where p is the P -value that certain structural and functional properties are impacted. Certain combinations of high values of general scores and low values of property scores are referred to as hypotheses (Li et al., 2009).

IDENTIFYING DISEASE-ASSOCIATED nsSNPs

nsSNP Analyzer (<http://snpanalyzer.uthsc.edu>) is a tool to predict whether a nsSNP has a phenotypic effect (disease-associated vs. neutral) using a machine learning method called Random Forest, and extracting structural and evolutionary information from a query nsSNP (Bao et al., 2005).

PREDICTION OF DELETERIOUS nsSNPs

PANTHER (<http://pantherdb.org/tools/csnpscoreForm.jsp>) estimates the likelihood of a particular nsSNP to cause a functional impact on a protein (Thomas et al., 2003). It calculates the substitution position-specific evolutionary conservation (subPSEC) score based on the alignment of evolutionarily related proteins. The subPSEC score is the negative logarithm of the probability ratio of the wild-type and the mutant amino acids at a particular position. The subPSEC scores are values from 0 (neutral) to about -10 (most likely to be deleterious).

PREDICTION OF PATHOLOGICAL MUTATIONS ON PROTEINS

PMUT (<http://mmb2.pcb.ub.es:8080/PMut>) uses a robust methodology to predict disease-associated mutations. PMUT method is based on the use of neural networks (NNs) trained with a large database of neutral mutations (NEMUs) and pathological mutations of mutational hot spots, which are obtained by alanine scanning, massive mutation, and genetically accessible mutations. The final output is displayed as a pathogenicity index ranging from 0 to 1 (indexes > 0.5 single pathological mutations) and a confidence index ranging from 0 (low) to 9 (high) (Costa et al., 2005).

PREDICTING THE FUNCTIONAL EFFECT OF AMINO ACID SUBSTITUTIONS

PROVEAN (Protein Variation Effect Analyzer) (<http://provean.jcvi.org>) is a sequence based predictor that estimates the effect of protein sequence variation on protein function (Choi et al., 2012). It is based on a clustering method where BLAST hits with more than 75% global sequence identity are clustered together and top 30 such clusters from a supporting sequence are averaged within and across clusters to generate the final PROVEAN score. A protein variant is predicted to be “deleterious” if the final score is below a certain threshold (default is -2.5), and is predicted to be “neutral” if the score is above the threshold.

PREDICTION OF DISEASE RELATED MUTATIONS

The SNPs&GO algorithms (<http://snps-and-go.biocomp.unibo.it/snps-and-go/>) predict the impact of protein variations using functional information encoded by Gene Ontology (GO) terms of the three main roots: Molecular function, Biological process, and Cellular component (Calabrese et al., 2009). SNPs&GO is a support vector machine (SVM) based web server to predict disease related mutations from the protein sequence, scoring with accuracy of 82% and Matthews correlation coefficient equal to 0.63. SNPs&GO collects, in a unique framework, information derived from protein sequence, protein sequence profile and protein functions.

RESULTS

nsSNPs FOUND BY SIFT PROGRAM

Protein sequence with mutational position and amino acid residue variants associated with 97 missense nsSNPs were submitted as input to the SIFT server, and the results are shown in **Table 1**. The lower the tolerance index, the higher the functional impact a particular amino acid residue substitution is likely to have and vice versa. Among the 97 nsSNPs analyzed, 47 nsSNPs

Table 1 | Tolerated and deleterious nsSNPs using SIFT.

S. No	rsID	Alleles	Position	AA change	Prediction	Score
1	rs121908314	L/V	371	Leu/Val	Damaging	0.04
2	rs121908313	F/L	251	Phe/Leu	Damaging	0.01
3	rs121908312	K/N	79	Lys/Asn	Tolerated	0.52
4	rs121908311	G/S	377	Gly/Ser	Damaging	0.02
5	rs121908310	V/F	398	Val/Phe	Damaging	0.01
6	rs121908308	R/G	353	Arg/Gly	Tolerated	0.38
7	rs121908307	S/T	364	Ser/Thr	Tolerated	0.12
8	rs121908306	C/G	342	Cys/Gly	Damaging	0.01
9	rs121908305	G/R	325	Gly/Arg	Tolerated	0.44
10	rs121908304	W/C	312	Trp/Cys	Damaging	0.00
11	rs121908303	F/V	216	Phe/Val	Damaging	0.00
12	rs121908302	V/L	15	Val/Leu	Tolerated	0.07
13	rs121908301	G/S	478	Gly/Ser	Tolerated	0.17
14	rs121908300	Y/H	212	Tyr/His	Damaging	0.03
15	rs121908299	P/S	122	Pro/Ser	Tolerated	0.37
16	rs121908298	P/L	289	Pro/Leu	Tolerated	0.48
17	rs121908297	K/Q	157	Lys/Gln	Tolerated	0.06
18	rs121908295	P/R	415	Pro/Arg	Damaging	0.00
19	rs80356773	R/H	496	Arg/His	Tolerated	0.19
20	rs80356772	R/H	463	Arg/His	Tolerated	0.06
21	rs80356771	R/C	463	Arg/Cys	Damaging	0.02
22	rs80356769	V/L	394	Val/Leu	Damaging	0.03
23	rs80356765	A/T	338	Ala/Thr	Tolerated	0.39
24	rs80356763	R/L	131	Arg/Leu	Tolerated	0.24
25	rs80205046	P/L	182	Pro/Leu	Damaging	0.00
26	rs80116658	G/D	265	Gly/Asp	Damaging	0.00
27	rs80020805	M/I	416	Met/Ile	Tolerated	0.42
28	rs79945741	F/L	213	Phe/Leu	Tolerated	0.18
29	rs79796061	D/V	127	Asp/Val	Damaging	0.00
30	rs79696831	R/H	285	Arg/His	Damaging	0.00
31	rs79653797	R/Q	120	Arg/Gln	Damaging	0.00
32	rs79637617	P/L	122	Pro/Leu	Damaging	0.02
33	rs79215220	P/R	266	Pro/Arg	Damaging	0.00
34	rs79185870	F/L	417	Phe/Leu	Damaging	0.01
35	rs78973108	R/Q	257	Arg/Gln	Tolerated	0.05
36	rs78911246	G/V	189	Gly/Val	Damaging	0.02
37	rs78802049	D/E	409	Asp/Glu	Tolerated	0.32
38	rs78769774	R/Q	48	Arg/Gln	Tolerated	0.06
39	rs78715199	D/E	380	Asp/Glu	Damaging	0.00
40	rs78396650	A/V	309	Ala/Val	Damaging	0.00
41	rs78198234	H/R	311	His/Arg	Damaging	0.00
42	rs78188205	A/D	318	Ala/Asp	Tolerated	0.63
43	rs77959976	M/I	123	Met/Ile	Tolerated	1.00
44	rs77834747	I/S	119	Ile/Ser	Tolerated	0.34
45	rs77829017	G/E	46	Gly/Glu	Damaging	0.01
46	rs77738682	N/I	392	Asn/Ile	Damaging	0.00
47	rs77451368	G/E	202	Gly/Glu	Damaging	0.02
48	rs77369218	D/V	409	Asp/Val	Tolerated	0.06
49	rs77321207	Y/C	395	Tyr/Cys	Damaging	0.00
50	rs77284004	D/A	380	Asp/Ala	Damaging	0.00
51	rs77035024	F/L	411	Phe/Leu	Tolerated	0.30
52	rs77019233	N/K	117	Asn/Lys	Tolerated	0.21
53	rs76910485	P/L	391	Pro/Leu	Damaging	0.00

(Continued)

Table 1 | Continued

S. No	rsID	Alleles	Position	AA change	Prediction	Score
54	rs76763715	N/S	370	Asn/Ser	Damaging	0.05
55	rs76763715	N/T	370	Asn/Thr	Damaging	0.04
56	rs76539814	T/I	323	Thr/Ile	Tolerated	0.48
57	rs76228122	Y/C	363	Tyr/Cys	Damaging	0.00
58	rs76026102	Y/C	205	Tyr/Cys	Damaging	0.00
59	rs76014919	W/C	378	Trp/Cys	Damaging	0.00
60	rs75954905	F/L	37	Phe/Leu	Tolerated	0.30
61	rs75671029	D/N	443	Asp/Asn	Tolerated	0.93
62	rs75636769	A/E	190	Ala/Glu	Tolerated	1.00
63	rs75564605	I/T	402	Ile/Thr	Damaging	0.04
64	rs75548401	T/M	369	Thr/Met	Tolerated	0.08
65	rs75528494	S/R	366	Ser/Arg	Damaging	0.03
66	rs75385858	N/T	396	Asn/Thr	Damaging	0.00
67	rs75243000	F/S	397	Phe/Ser	Damaging	0.02
68	rs75090908	D/E	399	Asp/Glu	Tolerated	0.17
69	rs74979486	R/Q	359	Arg/Gln	Tolerated	0.05
70	rs74953658	D/E	24	Asp/Glu	Damaging	0.01
71	rs74752878	Y/C	418	Tyr/Cys	Damaging	0.00
72	rs74731340	S/N	271	Ser/Asn	Tolerated	0.26
73	rs74598136	P/L	401	Pro/Leu	Damaging	0.00
74	rs74500255	F/Y	216	Phe/Tyr	Tolerated	0.34
75	rs74462743	G/E	195	Gly/Glu	Damaging	0.00
76	rs61748906	W/R	184	Trp/Arg	Damaging	0.00
77	rs11558184	R/Q	353	Arg/Gln	Tolerated	0.59
78	rs2230288	E/K	326	Glu/Lys	Tolerated	0.86
79	rs1141820	H/R	60	His/Arg	Tolerated	0.54
80	rs1141818	H/Y	60	His/Tyr	Tolerated	0.09
81	rs1141815	M/T	53	Met/Thr	Tolerated	0.59
82	rs1141814	R/W	48	Arg/Trp	Damaging	0.00
83	rs1141812	R/S	44	Arg/Ser	Tolerated	0.14
84	rs1141811	T/I	43	Thr/Ile	Damaging	0.01
85	rs1141811	T/R	43	Thr/Arg	Damaging	0.02
86	rs1141808	E/K	41	Glu/Lys	Tolerated	0.52
87	rs1141804	S/G	16	Ser/Gly	Tolerated	1.00
88	rs1141802	L/S	15	Leu/Ser	Tolerated	0.63
89	rs1064651	D/H	409	Asp/His	Tolerated	0.05
90	rs1064648	R/H	329	Arg/His	Tolerated	0.17
91	rs1064644	S/P	196	Ser/Pro	Tolerated	0.17
92	rs421016	L/P	444	Leu/Pro	Damaging	0.00
93	rs381737	F/I	213	Phe/Ile	Tolerated	0.18
94	rs381427	V/E	191	Val/Glu	Tolerated	0.16
95	rs381427	V/G	191	Val/Gly	Tolerated	0.16
96	rs368060	A/P	456	Ala/Pro	Tolerated	0.09
97	rs364897	N/S	188	Asn/Ser	Tolerated	0.17

The consensus SNPs are shown in bold.

were identified to be deleterious with a tolerance index score ≤ 0.05 (Kumar et al., 2009). Among 47 deleterious nsSNPs, 25 nsSNPs were found to be highly deleterious.

VALIDATION OF HARMFUL MUTATIONS

The MutPred score is the probability that an AAS is deleterious/disease-associated. A missense mutation with a

MutPred score > 0.5 could be considered as “harmful,” while a MutPred score > 0.75 should be considered a high confidence “harmful” prediction (Li et al., 2009). Among the 47 deleterious nsSNPs, 8 were found to be harmful mutations with a score of > 0.5 and < 0.75 and 38 were found to be high confidence (highly harmful) mutations and 1 nsSNP found to be normal with the score of 0.193 (Table 2).

Table 2 | Prediction of functional effects of nsSNPs using MutPred.

S. No	rsID	Alleles	Position	AA change	MutPred prediction	Score
1	rs121908314	L/V	371	Leu/val	High confidence	0.824
2	rs121908313	F/L	251	Phe/Leu	High confidence	0.778
3	rs121908311	G/S	377	Gly/Ser	Neutral	0.193
4	rs121908310	V/F	298	Val/Phe	High confidence	0.765
5	rs121908306	C/G	342	Cys/Gly	High confidence	0.792
6	rs121908304	W/C	312	Trp/Cys	Harmful mutation	0.735
7	rs121908303	F/V	216	Phe/Val	High confidence	0.879
8	rs121908300	Y/H	212	Tyr/His	High confidence	0.82
9	rs121908295	P/R	415	Pro/Arg	High confidence	0.914
10	rs80356771	R/C	463	Arg/Cys	Harmful mutation	0.664
11	rs80356769	V/L	394	Val/Leu	High confidence	0.794
12	rs80205046	P/L	182	Pro/Leu	High confidence	0.892
13	rs80116658	G/D	265	Gly/Asp	High confidence	0.963
14	rs79796061	D/V	127	Asp/Val	High confidence	0.754
15	rs79696831	R/H	285	Arg/His	High confidence	0.884
16	rs79653797	R/Q	120	Arg/Gln	High confidence	0.902
17	rs79637617	P/L	122	Pro/Leu	High confidence	0.835
18	rs79215220	P/R	166	Pro/Arg	High confidence	0.836
19	rs79185870	F/L	417	Phe/Leu	High confidence	0.905
20	rs78911246	G/V	189	Gly/Val	Harmful mutation	0.713
21	rs78715199	D/E	380	Asp/Glu	High confidence	0.837
22	rs78396650	A/V	309	Ala/Val	High confidence	0.776
23	rs78198234	H/R	311	His/Arg	High confidence	0.873
24	rs77829017	G/E	46	Gly/Glu	High confidence	0.856
25	rs77738682	N/I	392	Asn/Ile	High confidence	0.814
26	rs77451368	G/E	202	Gly/Glu	Harmful mutation	0.676
27	rs77321207	Y/C	304	Tyr/Cys	High confidence	0.909
28	rs77284004	D/A	380	Asp/Ala	High confidence	0.872
29	rs76910485	P/L	391	Pro/Leu	High confidence	0.889
30	rs76763715	N/S	370	Asn/Ser	High confidence	0.876
31	rs76763715	N/T	370	Asn/Thr	High confidence	0.89
32	rs76228122	Y/C	363	Tyr/Cys	High confidence	0.93
33	rs76026102	Y/C	205	Tyr/Cys	High confidence	0.857
34	rs76014919	W/C	378	Trp/Cys	High confidence	0.842
35	rs75564605	I/T	402	IleThr	High confidence	0.838
36	rs75528494	S/R	366	Ser/Arg	Harmful mutation	0.681
37	rs75385858	N/T	396	Asn/Thr	High confidence	0.848
38	rs75243000	F/S	397	Phe/Ser	Harmful mutation	0.724
39	rs74953658	D/E	24	Asp/Glu	High confidence	0.818
40	rs74752878	Y/C	418	Tyr/Cys	High confidence	0.872
41	rs74598136	P/L	401	Pro/Leu	High confidence	0.888
42	rs74462743	G/E	195	Gly/Glu	High confidence	0.859
43	rs61748906	W/R	184	Trp/Arg	High confidence	0.902
44	rs1141814	R/W	48	Arg/Trp	High confidence	0.804
45	rs1141811	T/I	43	Thr/Ile	Harmful mutation	0.504
46	rs1141811	T/R	43	Thr/Arg	Harmful mutation	0.579
47	rs421016	L/P	444	Leu/Pro	High confidence	0.899

The consensus SNPs are shown in bold.

DISEASE-ASSOCIATED nsSNPs

Out of 47 deleterious nsSNPs, 43 were found to be a disease causing nsSNPs and 4 were found to be neutral nsSNPs (Table 3).

VALIDATION BY PANTHER

The protein sequence was given as input and analyzed for the deleterious effect on protein function. The subPSEC scores are values from 0 (neutral) to about -10 (deleterious) (Thomas

Table 3 | The results from nsSNP Analyzer, PMUT, PROVEAN, and SNPs&GO.


S. No	rsID	Allele	Position	AA change	nsSNP Analyzer	PMUT	PROVEAN		SNPs&GO
							Score	Prediction	
1	rs121908314	L/V	371	Leu/val	Neutral	Neutral	-2.331	Neutral	Disease
2	rs121908313	F/L	251	Phe/Leu	Disease	Pathological	-4.567	Deleterious	Disease
3	rs121908311	G/S	377	Gly/Ser	Disease	Neutral	-5.128	Deleterious	Disease
4	rs121908310	V/F	398	Val/Phe	Disease	Neutral	-4.185	Deleterious	Disease
5	rs121908306	C/G	342	Cys/Gly	Disease	Pathological	-11.467	Deleterious	Disease
6	rs121908304	W/C	312	Trp/Cys	Disease	Pathological	-12.258	Deleterious	Disease
7	rs121908303	F/V	216	Phe/Val	Disease	Neutral	-7	Deleterious	Disease
8	rs121908300	Y/H	212	Tyr/His	Disease	Neutral	-4.267	Deleterious	Disease
9	rs121908295	P/R	415	Pro/Arg	Disease	Pathological	-8.793	Deleterious	Disease
10	rs80356771	R/C	463	Arg/Cys	Disease	Pathological	-5.279	Deleterious	Disease
11	rs80356769	V/L	394	Val/Leu	Neutral	Neutral	-2.031	Neutral	Disease
12	rs80205046	P/L	182	Pro/Leu	Disease	Neutral	-9.917	Deleterious	Disease
13	rs80116658	G/D	265	Gly/Asp	Disease	Neutral	-6.442	Deleterious	Disease
14	rs79796061	D/V	127	Asp/Val	Disease	Pathological	-8.625	Deleterious	Disease
15	rs79696831	R/H	285	Arg/His	Disease	Neutral	-4.792	Deleterious	Disease
16	rs79653797	R/Q	120	Arg/Gln	Disease	Neutral	-3.641	Deleterious	Disease
17	rs79637617	P/L	122	Pro/Leu	Disease	Neutral	-9.265	Deleterious	Disease
18	rs79215220	P/R	266	Pro/Arg	Disease	Neutral	-8.275	Deleterious	Disease
19	rs79185870	F/L	417	Phe/Leu	Disease	Neutral	-5.095	Deleterious	Disease
20	rs78911246	G/V	189	Gly/Val	Disease	Neutral	-6.4	Deleterious	Disease
21	rs78715199	D/E	380	Asp/Glu	Neutral	Neutral	-3.797	Deleterious	Disease
22	rs78396650	A/V	309	Ala/Val	Disease	Pathological	-3.533	Deleterious	Disease
23	rs78198234	H/R	311	His/Arg	Disease	Neutral	-7.667	Deleterious	Disease
24	rs77829017	G/E	46	Gly/Glu	Disease	Pathological	-5.925	Deleterious	Disease
25	rs77738682	N/I	392	Asn/Ile	Disease	Neutral	-7.593	Deleterious	Disease
26	rs77451368	G/E	202	Gly/Glu	Disease	Pathological	-5.178	Deleterious	Disease
27	rs77321207	Y/C	304	Tyr/Cys	Disease	Neutral	-8.358	Deleterious	Disease
28	rs77284004	D/A	380	Asp/Ala	Disease	Neutral	-7.593	Deleterious	Disease
29	rs76910485	P/L	391	Pro/Leu	Disease	Pathological	-9.269	Deleterious	Disease
30	rs76763715	N/S	370	Asn/Ser	Neutral	Neutral	-2.128	Neutral	Disease
31	rs76763715	N/T	370	Asn/Thr	Disease	Neutral	-3.062	Deleterious	Disease
32	rs76228122	Y/C	363	Tyr/Cys	Disease	Pathological	-8.492	Deleterious	Disease
33	rs76026102	Y/C	205	Tyr/Cys	Disease	Pathological	-7.552	Deleterious	Disease
34	rs76014919	W/C	378	Trp/Cys	Disease	Pathological	-12.306	Deleterious	Disease
35	rs75564605	I/T	402	IleThr	Disease	Pathological	-4.363	Deleterious	Disease
36	rs75528494	S/R	366	Ser/Arg	Disease	Pathological	-2.806	Deleterious	Disease
37	rs75385858	N/T	396	Asn/Thr	Disease	Neutral	-5.562	Deleterious	Disease
38	rs75243000	F/S	397	Phe/Ser	Disease	Pathological	-4.782	Deleterious	Disease
39	rs74953658	D/E	24	Asp/Glu	Disease	Neutral	-3.037	Deleterious	Disease
40	rs74752878	Y/C	418	Tyr/Cys	Disease	Pathological	-8.526	Deleterious	Disease
41	rs74598136	P/L	401	Pro/Leu	Disease	Pathological	-8.136	Deleterious	Disease
42	rs74462743	G/E	195	Gly/Glu	Disease	Pathological	-7.767	Deleterious	Disease
43	rs61748906	W/R	184	Trp/Arg	Disease	Pathological	-13.028	Deleterious	Disease
44	rs1141814	R/W	48	Arg/Trp	Disease	Pathological	-6.879	Deleterious	Disease
45	rs1141811	T/I	43	Thr/Ile	Disease	Neutral	-3.515	Deleterious	Disease
46	rs1141811	T/R	43	Thr/Arg	Disease	Pathological	-2.557	Deleterious	Disease
47	rs421016	L/P	444	Leu/Pro	Disease	Neutral	-4.995	Deleterious	Disease

The consensus SNPs are shown in bold.

Table 4 | Mutant scores from PANTHER.

S. NO	rsID	Alleles	Position	AA change	subPSEC	P _{deleterious}
1	rs121908314	L/V	371	Leu/val	-3.34802	0.58614
2	rs121908313	F/L	251	Phe/Leu	-2.59088	0.39912
3	rs121908311	G/S	377	Gly/Ser	-5.35062	0.91298
4	rs121908310	V/F	398	Val/Phe	-3.36629	0.59056
5	rs121908306	C/G	342	Cys/Gly	-3.57193	0.63921
6	rs121908304	W/C	312	Trp/Cys	-2.59838	0.40092
7	rs121908303	F/V	216	Phe/Val	-4.88341	0.868
8	rs121908300	Y/H	212	Tyr/His	-5.32716	0.9111
9	rs121908295	P/R	415	Pro/Arg	-4.90228	0.87015
10	rs80356771	R/C	463	Arg/Cys	-4.45218	0.81033
11	rs80356769	V/L	394	Val/Leu	-2.8436	0.46098
12	rs80205046	P/L	182	Pro/Leu	-6.3153	0.96495
13	rs80116658	G/D	265	Gly/Asp	-6.00914	0.95299
14	rs79796061	D/V	127	Asp/Val	-6.29967	0.96442
15	rs79696831	R/H	285	Arg/His	-4.32962	0.79078
16	rs79653797	R/Q	120	Arg/Gln	-4.52062	0.82063
17	rs79637617	P/L	122	Pro/Leu	-4.49053	0.81616
18	rs79215220	P/R	266	Pro/Arg	-6.27743	0.96365
19	rs79185870	F/L	417	Phe/Leu	-4.16977	0.7631
20	rs78911246	G/V	189	Gly/Val	-3.38537	0.59517
21	rs78715199	D/E	380	Asp/Glu	-2.02623	0.27413
22	rs78396650	A/V	309	Ala/Val	-4.2769	0.78192
23	rs78198234	H/R	311	His/Arg	-4.57198	0.82807
24	rs77829017	G/E	46	Gly/Glu	-5.04065	0.885
25	rs77738682	N/I	392	Asn/Ile	-4.02188	0.73534
26	rs77451368	G/E	202	Gly/Glu	-1.32995	0.15842
27	rs77321207	Y/C	304	Tyr/Cys	-6.26737	0.96329
28	rs77284004	D/A	380	Asp/Ala	-2.36947	0.34739
29	rs76910485	P/L	391	Pro/Leu	-6.12534	0.95793
30	rs76763715	N/S	370	Asn/Ser	-2.69603	0.42459
31	rs76763715	N/T	370	Asn/Thr	-1.97735	0.26451
32	rs76228122	Y/C	363	Tyr/Cys	-4.75749	0.85289
33	rs76026102	Y/C	205	Tyr/Cys	-5.89294	0.9475
34	rs76014919	W/C	378	Trp/Cys	-5.31772	0.91033
35	rs75564605	I/T	402	IleThr	-3.78009	0.6857
36	rs75528494	S/R	366	Ser/Arg	-2.07688	0.28432
37	rs75385858	N/T	396	Asn/Thr	-3.61569	0.64924
38	rs75243000	F/S	397	Phe/Ser	-2.88329	0.47086
39	rs74953658	D/E	24	Asp/Glu	-4.17446	0.76395
40	rs74752878	Y/C	418	Tyr/Cys	-6.31864	0.96506
41	rs74598136	P/L	401	Pro/Leu	-2.14888	0.2992
42	rs74462743	G/E	195	Gly/Glu	-4.74669	0.85153
43	rs61748906	W/R	184	Trp/Arg	-3.5793	0.64091
44	rs1141814	R/W	48	Arg/Trp	-7.03366	0.9826
45	rs1141811	T/I	43	Thr/Ile	-4.20869	0.77007
46	rs1141811	T/R	43	Thr/Arg	-4.05221	0.7412
47	rs421016	L/P	444	Leu/Pro	-3.43747	0.60766

The consensus SNPs are shown in bold.



et al., 2003). Out of 47 deleterious nsSNPs, 8 were found to be more than -6 (highly deleterious) and rest were found to be less deleterious. The mutant with a greater $P_{deleterious}$ tends to have more severe destructions in function. It was found that 32 out of 47 deleterious nsSNPs scored greater than 3 and rests were below the damage threshold (Table 4).

FUNCTIONAL IMPACT OF MUTATIONS ON PROTEINS

The functional impact of 47 deleterious nsSNPs in protein of GBA was analyzed using PMUT server. Of the 47 nsSNPs, 22

are classified as pathological, and the remaining were neutral (Table 3).

PROTEIN VARIATION EFFECT ANALYSIS

PROVEAN predicts the effect of the variant on the biological function of the protein based on sequence homology. PROVEAN scores are classified as “deleterious” if below a certain threshold (here -2.5) and “neutral” if above it (Choi et al., 2012). Out of 47 nsSNPs, 44 were predicted to be “deleterious” and 3 were found to be “neutral” (Table 3).

PREDICTION OF DISEASE RELATED MUTATIONS BY SNPs&GO

SNPs&GO is trained and tested with cross-validation procedures in which similar proteins are placed together as a dataset to calculate the *LGO* score derived from the GO data base. All 47 deleterious nsSNPs showed the disease related mutations (Table 3).

DISCUSSION

In the recent years, SNPs have emerged as the new generation molecular markers. The harmful SNPs for the GBA gene were never been predicted to date *in silico*. This study was designed to understand the genetic variations associated with GBA gene. We have predicted the harmful nsSNPs using SIFT, MutPred, nsSNP Analyzer, PANTHER, PMUT, PROVEAN, and SNPs&GO state of the art computational tools. Among 97 nsSNPs, 47 were found to be deleterious with a tolerance index score of ≤ 0.05 found by SIFT program. Among the 47 deleterious nsSNPs, 46 were found to be harmful nsSNPs found by MutPred, 43 were found to be disease causing nsSNPs by nsSNP Analyzer tool, 32 are highly deleterious found by PANTHER program, 22 are classified as pathological mutations by PMUT, 44 were predicted to be deleterious by PROVEAN server while all 47 deleterious nsSNPs showed the disease related mutations by SNPs&GO. Also, we found that SNPs&GO was most successful of all state of the art SNP prediction programs that were used for this comparative study. In this work, we found 22 nsSNPs that are common in all (SIFT, MutPred, nsSNP Analyzer, PANTHER, PMUT, PROVEAN, and SNPs&GO) prediction (Figure 1). These sets of 22 nsSNPs (F251L, C342G, W312C, P415R, R463C, D127V, A309V, G46E, G202E, P391L, Y363C, Y205C, W378C, I402T, S366R, F397S, Y418C, P401L, G195E, W184R, R48W, and T43R) are possibly the main targeted mutation for the GD (Tables 1–4). The previous work has shown that, in 60 patients with types 1 and 3, the most common Gaucher mutations identified were L444P, N370S, and R463C. L444P was the most common mutation in GD types 1, 2, and 3 (Latham et al., 1990; Sidransky et al., 1994). In our analysis, out of 7 methods, 6 methods (Sift, MutPred, PROVEAN, PANTHER, nsSNP Analyzer, and SNPs&GO) shows L444P mutation as damaging, 3 methods shows N370S mutation as damaging and all the 7 methods shows R463C mutation as damaging. D409H, A456P, E326K, and V460V mutations were also identified in patients with GD (Tsuji et al., 1987; Park et al., 2002). In our analysis SIFT result shows D409H, A456P, and E326K mutation is the tolerated mutation. Further studies using these mutations will shed light on the genetic understanding of this major lysosomal storage disease.

AUTHOR CONTRIBUTIONS

Madhumathi Manickam, Priti Talwar, and Palaniyandi Ravanan wrote the main manuscript and analyzed original datasets. Pratibha Singh prepared tables and figure. All authors reviewed the manuscript.

ACKNOWLEDGMENT

The authors greatly acknowledge the financial support and laboratory facilities given by VIT University, Vellore, India to carry out this research work. Additionally, authors greatly acknowledge the critical work of all reviewers on this paper.

REFERENCES

- Aharon, P. J., Rosenbaum, H., and Gershoni, B. R. (2004). Mutations in the glucocerebrosidase gene and Parkinson's disease in Ashkenazi Jews. *N. Engl. J. Med.* 351, 1972–1977. doi: 10.1056/NEJMoa033277
- Bao, L., Zhou, M., and Cui, Y. (2005). nsSNP Analyzer: identifying disease-associated nonsynonymous single nucleotide polymorphisms. *Nucleic Acids Res.* 33, 480–482. doi: 10.1093/nar/gki372
- Beutler, E., and Grabowski, G. A. (2001). “Gaucher disease,” in *Metabolic and Molecular Bases of Inherited Disease, II*, eds C. R. Scriver, A. L. Beaudet, W. S. Sly, and D. Valle (New York, NY: McGraw-Hill), 3635–3668.0
- Calabrese, R., Capriotti, E., Fariselli, P., Martelli, P. L., and Casadio, R. (2009). Functional annotations improve the predictive score of human disease-related mutations in proteins. *Hum. Mutat.* 8, 1237–1244. doi: 10.1002/humu.21047
- Cargill, M., Altshuler, D., Ireland, J., Sklar, P., Ardlie, K., Patil, N., et al. (1999). Characterization of single-nucleotide polymorphisms in coding regions of human genes. *Nat. Genet.* 22, 231–238. doi: 10.1038/10290
- Choi, Y., Sims, G. E., Murphy, S., Miller, J. R., and Chan, A. P. (2012). Predicting the functional effect of amino acid substitutions and indels. *PLoS ONE* 7:e46688. doi: 10.1371/journal.pone.0046688
- Costa, F. C., Gelpi, J. L., Zamakola, L., Parraga, I., Cruz, D. L. X., and Orozco, M. (2005). PMUT: a web-based tool for the annotation of pathological mutations on proteins. *Bioinformatics* 14, 3176–3178. doi: 10.1093/bioinformatics/bti486
- Costa, F. C., Orozco, M., and Cruz, D. L. X. (2002). Characterization of disease-associated single amino acid polymorphisms in terms of sequence and structure properties. *J. Mol. Biol.* 315, 771–786. doi: 10.1006/jmbi.2001.5255
- Grabowski, G. A. (2008). Phenotype, diagnosis, and treatment of Gaucher's disease. *Lancet* 372, 1263–1271. doi: 10.1016/S0140-6736(08)61522-6
- Horowitz, M., Wilder, S., Horowitz, Z., Reiner, O., Gelbart, T., and Beutler, E. (1989). The human glucocerebrosidase gene and pseudogene: structure and evolution. *Genomics* 4, 87–96. doi: 10.1016/0888-7543(89)90319-4
- Hruska, K. S., LaMarca, M. E., Scott, C. R., and Sidransky, E. (2008). Gaucher disease: mutation and polymorphism spectrum in the glucocerebrosidase gene (GBA). *Hum. Mutat.* 29, 567–583. doi: 10.1002/humu.20676
- James, W. D., Berger, T. M., and Elston, D. M. (2006). *Andrew's Diseases of the Skin: Clinical Dermatology*, 10th Edn. Philadelphia, PA: Saunders; Elsevier.
- Jmoudiak, M., and Futerman, A. H. (2005). Gaucher disease: pathological mechanisms and modern management. *Br. J. Haematol.* 129, 178–188. doi: 10.1111/j.1365-2141.2004.05351.x
- Kumar, P., Henikoff, S., and Ng, P. C. (2009). Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat. Protoc.* 4, 1073–1081. doi: 10.1038/nprot.2009.86
- Latham, T., Grabowski, G. A., Theophilus, B. D. M., and Smith, F. I. (1990). Complex alleles of the acid beta-glucosidase gene in Gaucher disease. *Am. J. Hum. Genet.* 47, 79–86.
- Li, B., Krishnan, V. G., Mort, M. E., Xin, F., Kamati, K. K., Cooper, D. N., et al. (2009). Automated inference of molecular mechanisms of disease from amino acid substitutions. *Bioinformatics* 25, 2744–2750. doi: 10.1093/bioinformatics/btp528
- Mi, H., Muruganujan, A., and Thomas, P. D. (2012). PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic Acids Res.* 41, 377–386. doi: 10.1093/nar/gks1118
- Ng, P. C., and Henikoff, S. (2001). Predicting deleterious amino acid substitutions. *Genome Res.* 11, 863–874. doi: 10.1101/gr.176601
- Park, J. K., Tayebi, N., Stubblefield, B. K., LaMarca, M. E., MacKenzie, J. J., Stone, D. L., et al. (2002). The E326K mutation and Gaucher disease: mutation or polymorphism? *Clin. Genet.* 61, 32–34. doi: 10.1034/j.1399-0004.2002.610106.x
- Pilar, G., Pilar, A., Pilar, I., Laura, G., Amparo, C., Lluisa, V., et al. (2012). Mapping the genetic and clinical characteristics of Gaucher disease in the Iberian Peninsula. *Orphanet J. Rare Dis.* 7:17. doi: 10.1186/1750-1172-7-17
- Sidransky, E., Bottler, A., Stubblefield, B., and Ginns, E. I. (1994). DNA mutational analysis of type 1 and type 3 Gaucher patients: how well do mutations predict phenotype? *Hum. Mutat.* 3, 25–28. doi: 10.1002/humu.1380030105
- Thomas, P. D., Campbell, M. J., Kejarawal, A., Mi, H., Karlak, B., Daverman, R., et al. (2003). PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res.* 13, 2129–2141. doi: 10.1101/gr.772403
- Tsuji, S., Choudary, P. V., Martin, B. M., Stubblefield, B. K., Mayor, J. A., Barranger, J. A., et al. (1987). A mutation in the human glucocerebrosidase gene in neuronopathic Gaucher's disease. *N. Engl. J. Med.* 316, 570–575. doi: 10.1056/NEJM198703053161002

- Wheeler, D. L., Barrett, T., Benson, D. A., Bryant, S. H., Canese, K., Chetvernin, V., et al. (2005). Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res.* 34, 173–180. doi: 10.1093/nar/gkj158
- Winfield, S. L., Tayebi, N., Martin, B. M., Ginns, E. I., and Sidransky, E. (1997). Identification of three additional genes contiguous to the glucocerebrosidase locus on chromosome 1q21: implications for Gaucher disease. *Genome Res.* 7, 1020–1026.
- Zimran, A., Gelbart, T., Westwood, B., Grabowski, G. A., and Beutler, E. (1991). High frequency of the Gaucher disease mutation at nucleotide 1226 among Ashkenazi Jews. *Am. J. Hum. Genet.* 49, 855–859.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 14 March 2014; accepted: 06 May 2014; published online: 27 May 2014.

Citation: Manickam M, Ravanan P, Singh P and Talwar P (2014) In silico identification of genetic variants in glucocerebrosidase (GBA) gene involved in Gaucher's disease using multiple software tools. *Front. Genet.* 5:148. doi: 10.3389/fgene.2014.00148

This article was submitted to *Genetic Disorders*, a section of the journal *Frontiers in Genetics*.

Copyright © 2014 Manickam, Ravanan, Singh and Talwar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Copyright of *Frontiers in Genetics* is the property of *Frontiers Media S.A.* and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.